

两参数非线性反应扩散积分微分方程 的内层激波渐近解*

欧阳成¹ 莫嘉琪²

摘要 研究了一类两参数非线性反应扩散积分微分奇摄动问题. 利用奇摄动方法, 构造了问题的外部解、内部激波层、边界层及初始层校正项, 由此得到了问题解的形式渐近展开式. 最后利用积分微分方程的比较定理证明了该问题解的渐近展开式的一致有效性.

关键词 反应扩散, 奇异摄动, 初边值问题

MR (2010) 主题分类 35B25

中图法分类 O175.29

文献标志码 A

文章编号 1000-8314(2017)04-0365-10

1 引言

非线性奇异摄动问题在国内外学术界十分重视^[1-2]. 近年来奇异摄动渐近方法有很大改进, 包括边界层校正法、匹配法、平均法和多重尺度法等等, 许多学者作了很多的工作^[1-8], 利用比较定理和其它方法, 作者等也研究了一些奇异摄动问题^[9-23]. 本文讨论一类具有两参数的非线性积分微分反应扩散系统的奇异摄动初边值问题.

考虑如下形式两参数非线性反应扩散系统初边值问题:

$$\mu \frac{\partial u_i}{\partial t} - \varepsilon^2 Lu_i + Tu_i = f_i(t, x, u_i), \quad 0 < t \leq T_0, \quad x \in \Omega, \quad i = 1, 2, \dots, m, \quad (1.1)$$

$$u_i = g_i(t, x), \quad x \in \partial\Omega, \quad i = 1, 2, \dots, m, \quad (1.2)$$

$$u_i(0, x) = h_i(x), \quad i = 1, 2, \dots, m, \quad (1.3)$$

其中

$$L = \sum_{j,k=1}^n \alpha_{jk}(x) \frac{\partial^2}{\partial x_j \partial x_k}, \quad \sum_{j,k=1}^n \alpha_{jk}(x) \xi_j \xi_k \geq \lambda \sum_{j=1}^n \xi_j^2, \quad \lambda > 0,$$

$$Tu_i = \int_{\Omega} K(t, x, y) u_i dy,$$

且 ε 和 μ 为正的小参数, $x = (x_1, x_2, \dots, x_n) \in \Omega$, Ω 为 \mathbb{R}^n 中的有界区域, T_0 为正常数, $\frac{\partial}{\partial n}$ 表示在边界 $\partial\Omega$ 上的外法向导数. 假设:

本文 2016 年 2 月 16 日收到, 2016 年 12 月 24 日收到修改稿.

¹湖州师范学院理学院, 浙江 湖州 313000. E-mail: oyc@hutc.zj.cn

²安徽师范大学数学系, 安徽 芜湖 241003. E-mail: mojiqqi@mail.ahnu.edu.cn

*本文受到国家自然科学基金 (No. 11202106) 和浙江省自然科学基金项目 (No. LY13A010005) 的资助.

[H₁] 线性算子 L 的系数及 f_i (除 $f_i(t, x_0, u_i(t, x_0))$ 外), g_i, h_i 和 K 在各自的定义域内为充分光滑的函数, 且 $g_i(0, x) = A_i(x)$, $x \in \partial\Omega$, 并存在常数 $N_i > 0$ 和 $M_i > 0$, 使得 $-N_i \leq f_i u_i \leq -M_i$ ($i = 1, 2, \dots, m$);

[H₂] $K(t, x, y) \geq 0$, $\int_{\Omega} K(t, x, y) u_i dy \leq d$, 其中 $d > 0$ 为常数.

2 积分微分方程的比较定理

定义 2.1 若 $\bar{u} = (\bar{u}_1, \bar{u}_2, \dots, \bar{u}_m)$, $\underline{u} = (\underline{u}_1, \underline{u}_2, \dots, \underline{u}_m)$ 是 $(t, x) \in [0, T_0] \times (\Omega + \partial\Omega)$ 上的光滑函数, $\underline{u}_i \leq \bar{u}_i$ ($i = 1, 2, \dots, m$), 满足

$$\begin{aligned} \mu(\underline{u}_i)_t - \varepsilon^2 L\underline{u}_i + T\underline{u}_i - f_i(t, x, \underline{u}_i) &\leq 0 \leq \mu(\bar{u}_i)_t - \varepsilon^2 L\bar{u}_i + T\bar{u}_i - f_i(t, x, \bar{u}_i), \\ \underline{u}_i &\leq g_i(t, x) \leq \bar{u}_i, \quad x \in \partial\Omega, \quad \underline{u}_i \leq A_i(x) \leq \bar{u}_i, \quad t = 0, \end{aligned}$$

则称 \bar{u} 和 \underline{u} 分别为积分微分方程初边值问题 (1.1)–(1.3) 的上解和下解.

定理 2.1 (比较定理) 假设 [H₁], [H₂] 成立, 有两个充分小的正常数 ε_0, μ_0 , $\forall \varepsilon \in (0, \varepsilon_0)$, $\mu \in (0, \mu_0)$, 若初边值问题 (1.1)–(1.3) 有一对上、下解 (\bar{u}, \underline{u}) , 则积分微分方程初边值问题 (1.1)–(1.3) 存在解 $u = (u_1, u_1, \dots, u_m)$, 且 $\underline{u}_i \leq u_i \leq \bar{u}_i$, $(t, x) \in [0, T_0] \times (\Omega + \partial\Omega)$.

证 设 $\bar{u}_i^0 = \bar{u}_i \geq \underline{u}_i^0 = \underline{u}_i$ ($i = 1, 2, \dots, m$) 为两个初始函数, 则能按下列线性系统分别构造出两个迭代序列 $\{\bar{u}_i^k\}, \{\underline{u}_i^k\}$ ($i = 1, 2, \dots, m$):

$$\begin{aligned} \mu(\bar{u}_i^k)_t - \varepsilon^2 L\bar{u}_i^k + T\bar{u}_i^k + \sum_{l=1}^m N_l \bar{u}_i^k &= \sum_{l=1}^m N_l \bar{u}_i^{k-1} + f_i(t, x, \bar{u}_i^{k-1}), \quad x \in \Omega, \\ \bar{u}_i^k &= g_i(t, x), \quad x \in \partial\Omega, \quad \bar{u}_i^k(0, x) = h_i(x), \quad x \in \Omega, \\ \mu(\underline{u}_i^k)_t - \varepsilon^2 L\underline{u}_i^k + T\underline{u}_i^k + \sum_{l=1}^m N_l \underline{u}_i^k &= \sum_{l=1}^m N_l \underline{u}_i^{k-1} + f_i(t, x, \underline{u}_i^{k-1}), \quad x \in \Omega, \\ \underline{u}_i^k &= g_i(t, x), \quad x \in \partial\Omega, \quad \underline{u}_i^k(0, x) = h_i(x), \quad x \in \Omega. \end{aligned}$$

事实上, 由假设, 线性反应扩散微分积分方程组初边值问题的存在唯一性理论知, 由上述两组决定的线性反应扩散微分积分方程组初边值问题均依次地存在相应的解^[3].

令 $w_i = \bar{u}_i^0 - \bar{u}_i^1$, 由假设, 有

$$\begin{aligned} \mu(w_i)_t - \varepsilon^2 Lw_i + Tw_i + \sum_{l=1}^m N_l w_i \\ = \mu(\bar{u}_i)_t - \varepsilon^2 L\bar{u}_i + T\bar{u}_i + f_i(t, x, \bar{u}_i) &\geq 0, \quad x \in \Omega, \\ w_i &= 0, \quad x \in \partial\Omega, \quad w_i(0, x) = 0, \quad x \in \Omega. \end{aligned}$$

于是 $w_i \geq 0$ ^[1,2], 即

$$\bar{u}_i^1 \leq \bar{u}_i^0, \quad x \in \Omega + \partial\Omega, \quad i = 1, 2, \dots, m.$$

同理可得

$$\underline{u}_i^1 \geq \underline{u}_i^0, \quad x \in \Omega + \partial\Omega, \quad i = 1, 2, \dots, m.$$

现证 $\bar{u}_i^1 \geq \underline{u}_i^1$. 设 $\bar{w} = \bar{u}_i^1 - \underline{u}_i^1$,

$$\begin{aligned} & \mu(\bar{w}_i)_t - \varepsilon^2 L\bar{w}_i + T\bar{w}_i + \sum_{l=1}^m N_l \bar{w}_i \\ &= N_i(\bar{u}_i^0 - \underline{u}_i^0) + [f_i(t, x, \bar{u}_i^0) - f_i(t, x, \underline{u}_i^0)] \geq 0, \quad x \in \Omega, \\ & \bar{w}_i = 0, \quad x \in \partial\Omega, \quad \bar{w}_i(0, x) = 0, \quad x \in \Omega, \end{aligned}$$

则 $\bar{w}_i \geq 0$, 即

$$\underline{u}_i^1 \leq \bar{u}_i^1, \quad x \in \Omega + \partial\Omega, \quad i = 1, 2, \dots, m.$$

类似地可得

$$\begin{aligned} \underline{u}_i &= \underline{u}_i^0 \leq \underline{u}_i^1 \leq \dots \leq \underline{u}_i^k \leq \dots \leq \bar{u}_i^k \leq \dots \leq \bar{u}_i^1 \leq \bar{u}_i^0 = \underline{u}_i, \\ 0 \leq t \leq T_0, \quad x \in \Omega + \partial\Omega, \quad i &= 1, 2, \dots, m. \end{aligned}$$

由文 [1-2], 在本文的假设下, 成立

$$\lim_{k \rightarrow \infty} \underline{u}_i^k = u_i, \quad 0 \leq t \leq T_0, \quad x \in \Omega + \partial\Omega, \quad i = 1, 2, \dots, m,$$

且 $u = (u_1, u_2, \dots, u_m)$ 是积分微分方程初边值问题 (1.1)–(1.3) 的唯一解. 定理 2.1 证毕.

3 初边值问题的外部解

积分微分方程初边值问题 (1.1)–(1.3) 的退化系统为

$$Tu_i = f_i(t, x, u_i), \quad 0 < t \leq T_0, \quad x \in \Omega, \quad i = 1, 2, \dots, m. \quad (3.1)$$

由假设知, Fredholm 型积分方程组 (3.1) 有唯一的一组光滑解 $(U_{100}, U_{200}, \dots, U_{m00})$.

设系统的外部解为 $U = (U_1, U_2, \dots, U_m)$, 且

$$U_i(t, x) = \sum_{j,k=0}^{\infty} U_{ijk} \varepsilon^j \mu^k, \quad i = 1, 2, \dots, m. \quad (3.2)$$

将 (3.2) 代入系统 (1.1), 取 $\varepsilon = \mu = 0$ 便为系统 (3.1), 故 (3.2) 中的 $(U_{100}, U_{200}, \dots, U_{m00})$ 就是积分系统 (3.1) 的一组解. 由 (3.1), 对于 $j, k = 0, 1, \dots, j+k \neq 0$, 由 $\varepsilon^j \mu^k$ 的同次幂的系数, 有

$$\begin{aligned} & TU_{ijk}(t, x) \\ &= f_{iu_i}(t, x, U_{i00})U_{ijk} + F_{ijk}, \quad j, k = 0, 1, \dots, j+k \neq 0, \quad i = 1, 2, \dots, m, \end{aligned} \quad (3.3)$$

其中 F_{ijk} 为逐次已知的函数, 其结构从略. 同样地, 由假设, Fredholm 型积分方程组 (3.3) 有一组解 $(U_{1jk}, U_{2jk}, \dots, U_{mjk})$, $j, k = 0, 1, \dots, j+k \neq 0, i = 1, 2, \dots, m$. 由 (3.2), 得到原非线性反应扩散系统初边值问题的外部解 $U = (U_1, U_2, \dots, U_m)$. 但它未必在 $x_0 \in \Omega$ 处连续, 且未必满足原非线性反应扩散系统的边界条件和初始条件 (1.2)–(1.3), 所以尚需构造在 $x_0 \in \Omega$ 附近的内部激波层校正项 $Z = (Z_1, Z_1, \dots, Z_m)$ 和边界层校正项 $V = (V_1, V_1, \dots, V_m)$ 以及初始层校正项 $W = (W_1, W_1, \dots, W_m)$.

4 在 x_0 邻域的内部激波解

在 $x_0 \in \Omega$ 的邻域引入伸长变量 $\sigma = \frac{|x-x_0|}{\varepsilon}$. 设内部激波层校正项 $Z = (Z_1, Z_1, \dots, Z_m)$ 为

$$Z_i = \sum_{j,k=0}^{\infty} z_{ijk} \varepsilon^j \mu^k, \quad i = 1, 2, \dots, m, \quad (4.1)$$

并令

$$u_i = U_i + Z_i, \quad i = 1, 2, \dots, m. \quad (4.2)$$

在 x_0 的邻域上, 将 (4.1)–(4.2) 代入积分微分方程初边值问题的系统 (1.1), 可得

$$\mu \frac{\partial z_i}{\partial t} - \varepsilon^2 Lz_i + Tz_i = f_i(t, x_0 \pm \varepsilon\sigma, U_i + z_i) - f_i(t, x_0 \pm \varepsilon\sigma, z_i), \quad i = 1, 2, \dots, m,$$

并对非线性项按 ε, μ 的幂展开, 对应的 $\varepsilon^j \mu^k$ ($j, k = 0, 1, \dots$) 的各次幂的系数, 有

$$Tz_{i00} = f_i(t, x, z_{i00}), \quad i = 1, 2, \dots, m, \quad (4.3)$$

$$\lim_{x \rightarrow x_0} z_{i00} = U_{i00}(t, x_0), \quad i = 1, 2, \dots, m, \quad (4.4)$$

$$Tz_{ijk} = \bar{F}_{ijk}, \quad j, k = 0, 1, \dots, j+k \neq 0, \quad i = 1, 2, \dots, m, \quad (4.5)$$

$$\lim_{x \rightarrow x_0} z_{ijk} = U_{ijk}(t, x_0), \quad j, k = 0, 1, \dots, j+k \neq 0, \quad i = 1, 2, \dots, m, \quad (4.6)$$

其中 \bar{F}_{ijk} 为逐次已知的函数, 他们的结构在此从略. 由 (4.3)–(4.6) 可依次得到 z_{ijk} ($i = 1, 2, \dots, m$), 并由假设知, 它们具有内部激波层性态

$$z_{ijk} = O\left(\exp\left(-\bar{k}_{ijk} \frac{|x-x_0|}{\varepsilon}\right)\right), \quad j, k = 0, 1, \dots, \quad i = 1, 2, \dots, m, \quad (4.7)$$

其中 \bar{k}_{ijk} 为适当小的正常数.

引入一个充分光滑的分隔函数 $\psi(x)$, 使得

$$\psi(x) = \begin{cases} 1, & 0 \leq |x-x_0| \leq \frac{1}{3}\delta, \\ 0, & \frac{2}{3}\delta \leq |x-x_0|, \end{cases}$$

其中 δ 为足够小的常数. 取 $\bar{z}_{ijk} = \psi z_{ijk}$ ($j, k = 0, 1, \dots, i = 1, 2, \dots, m$). 为了方便起见, 下面仍然用 z_{ijk} 来代替 \bar{z}_{ijk} . 将得到的 z_{ijk} 代入 (4.1), 便得到了在 $x_0 \in \Omega$ 的邻域的激波层校正项 $Z = (Z_1, Z_2, \dots, Z_m)$.

5 边界层校正项

在 Ω 的边界 $\partial\Omega$ 的邻域建立一个局部坐标系 (ρ, ϕ) , 在 Ω 的边界 $\partial\Omega$ 的邻域内的每一点 P 的坐标 ρ ($\rho \leq \rho_0$) 为点 P 到 $\partial\Omega$ 的距离, 这里 ρ_0 为足够小的正常数, 并使在边界 $\partial\Omega$ 上的每一点的内法线在边界 $\partial\Omega$ 的邻域 $0 \leq \rho \leq \rho_0$ 内互不相交. 而 $\phi = (\phi_1, \phi_2, \dots, \phi_{n-1})$ 为在 $\partial\Omega$ 上的一个 $n-1$ 维非奇局部坐标系, 且设点 P 的坐标 ϕ 为通过点 P 的内法线和

边界 $\partial\Omega$ 的交点 Q 的坐标 ϕ 相同. 由此, 在 $\partial\Omega$ 的邻域 $0 \leq \rho \leq \rho_0$ 中, 有

$$L = a_{nn} \frac{\partial^2}{\partial \rho^2} + \sum_{j,i=1}^{n-1} a_{jn}(x) \frac{\partial^2}{\partial \rho \partial \phi_i} + \sum_{j,i=1}^{n-1} a_{ij} \frac{\partial^2}{\partial \phi_i \partial \phi_j} + b_{in} \frac{\partial}{\partial \rho} + \sum_{i=1}^{n-1} b_i \frac{\partial}{\partial \phi_i},$$

其中 $a_{nn}, a_{jn}, a_{ij}, b_{in}, b_i$ 为已知函数, 它们的结构从略.

再在 $0 \leq \rho \leq \rho_0$ 上引入一组多重尺度变量^[1-2]:

$$\zeta = \frac{h(\rho, \phi)}{\varepsilon}, \quad \bar{\rho} = \rho, \quad \phi = \phi, \quad (5.1)$$

其中 $h(\rho, \phi)$ 为适当确定的函数. 不失一般性, 下面仍用 ρ 来代替 $\bar{\rho}$. 于是有

$$L = \frac{1}{\varepsilon^2} K_0 + \frac{1}{\varepsilon} K_1 + K_2, \quad (5.2)$$

其中 $K_0 = a_{nn} h_\rho^2 \frac{\partial^2}{\partial \zeta^2}$, 而 K_1, K_2 的结构在此从略. 设积分微分方程初边值问题 (1.1)–(1.3) 的解 (u_1, u_2, \dots, u_n) 为

$$u_i = U_i + Z_i + V_i, \quad i = 1, 2, \dots, m, \quad (5.3)$$

其中

$$V_i = \sum_{j,k=0}^{\infty} v_{ijk}(t, \rho, \phi) \varepsilon^j \mu^k, \quad i = 1, 2, \dots, m. \quad (5.4)$$

将 (3.2), (4.1), (5.2)–(5.4) 代入 (1.1)–(1.2), 按 ε, μ 展开非线性项, 再由 $\varepsilon^j \mu^k$ ($j, k = 0, 1, \dots$) 同次幂的系数, 得到

$$\begin{aligned} & K_0 v_{i00} - T v_{i00} \\ &= -[f_i(t, \rho, \phi, U_{i00} + z_{i00} + v_{i00}) - f_i(t, \rho, \phi, U_{i00} + z_{i00})], \quad i = 0, 1, \dots, m, \end{aligned} \quad (5.5)$$

$$v_{i00} = g_i(t, 0, \phi) - (U_{i00} + z_{i00}), \quad i = 0, 1, \dots, m \quad (5.6)$$

和

$$\begin{aligned} & K_0 v_{ijk} - T v_{ijk} = -f_{i u_i}(t, \rho, \phi, U_{i00} + z_{i00} + v_{ijk}) v_{ijk} + G_{ijk}, \\ & j, k = 0, 1, \dots, j+k \neq 0, \quad i = 1, 2, \dots, m, \end{aligned} \quad (5.7)$$

$$v_{ijk} = -(U_{ijk} + z_{ijk}), \quad j, k = 0, 1, \dots, j+k \neq 0, \quad i = 1, 2, \dots, m, \quad (5.8)$$

其中 G_{ijk} 为逐次已知的函数, 它们的结构从略.

由 (5.5)–(5.8) 和假设知, 能够依次得到具有衰减性态的解 $v_{ijk}(t, \rho, \phi)$:

$$v_{ijk}(t, \rho, \phi) = O\left(\exp\left(-\widehat{k}_{ijk} \frac{\rho}{\varepsilon}\right)\right), \quad j, k = 0, 1, \dots, \quad i = 1, 2, \dots, m, \quad (5.9)$$

其中 \widehat{k}_{ijk} 为适当小的正常数.

再引入一个充分光滑的分隔函数 $\bar{\psi}(\rho)$, 使得

$$\bar{\psi}(\rho) = \begin{cases} 1, & 0 \leq \rho \leq \frac{1}{3}\delta, \\ 0, & \frac{2}{3}\delta \leq \rho. \end{cases}$$

取 $\bar{v}_{ijk} = \bar{\psi} v_{ijk}$ ($j, k = 0, 1, \dots, i = 1, 2, \dots, m$). 为了方便起见, 下面仍然用 v_{ijk} 来代替 \bar{v}_{ijk} . 由 (5.4), 便得到具有边界层校正性质的函数 $V = (V_1, V_2, \dots, V_m)$.

6 初始层校正项

设

$$u_i = U_i + Z_i + V_i + W_i, \quad i = 1, 2, \dots, m \quad (6.1)$$

和

$$W_i(\tau, x) = \sum_{j,k=0}^{\infty} w_{ijk}(\tau, x) \varepsilon^j \mu^k, \quad i = 1, 2, \dots, m, \quad (6.2)$$

其中 $\tau = \frac{t}{\mu}$ 为伸长变量^[1-2].

将(6.1)–(6.2)代入(1.1), (1.3), 按 ε, μ 展开非线性项, 再由 $\varepsilon^j \mu^k$ ($j, k = 0, 1, \dots$) 同次幂的系数, 得到

$$\begin{aligned} \frac{\partial w_{i00}}{\partial \tau} + T w_{i00} &= f_i(0, x, U_{i00} + z_{i00} + v_{i00} + w_{i00}) \\ &\quad - f_i(0, x, U_{i00} + z_{i00} + v_{i00}), \quad i = 1, 2, \dots, m, \end{aligned} \quad (6.3)$$

$$w_{i00} = h_i(x) - U_{i00} - z_{i00} - v_{i00}, \quad i = 1, 2, \dots, m, \quad (6.4)$$

$$\begin{aligned} \frac{\partial w_{ijk}}{\partial \tau} &= f_{iu_i}(0, x, U_{i00} + Z_{i00} + V_{i00} + W_{i00}) w_{ijk} + \tilde{F}_{ijk}, \\ j, k &= 0, 1, \dots, \quad j + k \neq 0, \quad i = 1, 2, \dots, m, \end{aligned} \quad (6.5)$$

$$w_{ijk} = -(U_{ijk} + z_{ijk} + v_{ijk}), \quad j, k = 0, 1, \dots, \quad j + k \neq 0, \quad i = 1, 2, \dots, m, \quad (6.6)$$

其中 \tilde{F}_{ijk} 为逐次已知的函数, 其结构也从略. 由(6.3)–(6.6), 我们能依次得到 w_{ijk} ($j, k = 0, 1, \dots, i = 0, 1, \dots, m$), 且它们具有初始层性态

$$w_{ijk} = O\left(\exp\left(-\tilde{k}_{ijk} \frac{t}{\mu}\right)\right), \quad j, k = 0, 1, \dots, \quad i = 1, 2, \dots, m, \quad (6.7)$$

其中 \tilde{k}_{ijk} 为适当小的正常数.

再引入一个充分光滑的分隔函数 $\tilde{\psi}(\tau)$, 使得

$$\tilde{\psi}(\tau) = \begin{cases} 1, & 0 \leq \tau \leq \frac{1}{3}\delta, \\ 0, & \frac{2}{3}\delta \leq \tau. \end{cases}$$

取 $\bar{w}_{ijk} = \tilde{\psi} w_{ijk}$ ($j, k = 0, 1, \dots, i = 1, 2, \dots, m$). 为了方便起见, 下面仍然用 w_{ijk} 来代替 \bar{w}_{ijk} . 再将 $w_{ijk}(\tau, x)$ 代入(6.2), 便得到具有初始层校正性质的函数 $W = (W_1, W_2, \dots, W_m)$.

由(6.1), 两参数非线性反应扩散系统初边值问题(1.1)–(1.3)的解 (u_1, u_2, \dots, u_m) 有如下形式的渐近展开式

$$\begin{aligned} u_i &\sim \sum_{j=0}^M \sum_{k=0}^N (U_{ijk} + z_{ijk} + v_{ijk} + w_{ijk}) \varepsilon^j \mu^k + O(\lambda), \\ &\quad i = 1, 2, \dots, m, \quad 0 < \varepsilon, \mu \ll 1, \end{aligned} \quad (6.8)$$

其中 $\lambda = \max(\varepsilon^{M+1} \mu^N, \varepsilon^M \mu^{N+1})$, $0 < \varepsilon, \mu \ll 1$.

7 解的一致有效性

定理 7.1 在假设 $[H_1], [H_2]$ 下, 非线性反应扩散系统初边值问题 (1.1)–(1.3) 存在解 $u = (u_1, u_2, \dots, u_m)$, 并在 $[0, T_0] \times (\Omega + \partial\Omega)$ 上关于小参数 ε, μ 成立一致有效的渐近展开式 (6.8).

证 首先构造辅助函数 α_i, β_i :

$$\alpha_i = Y_i - r_i \lambda, \quad \beta_i = Y_i + r_i \lambda, \quad i = 1, 2, \dots, m, \quad 0 < \varepsilon, \mu \ll 1, \quad (7.1)$$

其中 r_i ($i = 1, 2, \dots, m$) 为足够大的正常数, 它们将在下面决定, 而

$$Y_i = \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N (U_{ijk} + z_{ijk} + v_{ijk} + w_{ijk}) \varepsilon^j \mu^k.$$

显然

$$\alpha_i \leq \beta_i. \quad (7.2)$$

由假设不难看出, 存在正常数 D_{i1} ($i = 1, 2, \dots, m$), 在 $x \in \partial\Omega$ 上成立

$$\begin{aligned} \alpha_i|_{\partial\Omega} &\leq g_i(t, 0, \phi) + \left[\sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N (U_{ijk} + z_{ijk} + v_{ijk} + w_{ijk}) \varepsilon^j \mu^k \right]_{\partial\Omega} - r_i \lambda \\ &\leq g_i(t, x) + (D_{i1} - r_i), \quad x \in \partial\Omega. \end{aligned}$$

选取 $r_i \geq D_{i1}$, 便有

$$\alpha_i \leq g_i(t, x), \quad x \in \partial\Omega, \quad i = 1, 2, \dots, m. \quad (7.3)$$

由 (4.7), (5.9), (6.7), 存在正常数 D_{i2} ($i = 1, 2, \dots, m$), 有

$$\alpha_i|_{t=0} = \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N [U_{ijk} + z_{ijk} + v_{ijk} + w_{ijk}]_{t=0} \varepsilon^j \mu^k - r_i \lambda \leq h_i(x) + (D_{i2} - r_i) \lambda.$$

于是, 当 $r_i \geq D_{i2}$ 时, 有

$$\alpha_i \leq h_i(x), \quad t = 0, \quad i = 1, 2, \dots, m. \quad (7.4)$$

同理可得

$$\beta_i|_{\partial\Omega} \geq g_i(t, x), \quad \beta_i|_{t=0} \geq h_i(x), \quad i = 1, 2, \dots, m. \quad (7.5)$$

现证

$$\mu \frac{\partial \alpha_i}{\partial t} - \varepsilon^2 L \alpha_i + T \alpha_i - f_i(t, x, \alpha_i) \leq 0, \quad 0 < t \leq T_0, \quad x \in \Omega, \quad i = 1, 2, \dots, m, \quad (7.6)$$

$$\mu \frac{\partial \beta_i}{\partial t} - \varepsilon^2 L \beta_i + T \beta_i - f_i(t, x, \beta_i) \geq 0, \quad 0 < t \leq T_0, \quad x \in \Omega, \quad i = 1, 2, \dots, m. \quad (7.7)$$

我们区分如下 3 种情形

(i) 当 $\Omega \setminus (\rho \geq (\frac{2}{3})\delta)$ 时; (ii) 当 $(\frac{1}{3})\delta \leq \rho \leq (\frac{2}{3})\delta$ 时; (iii) 当 $0 \leq \rho \leq (\frac{1}{3})\delta$ 时. 现只证明情形 (iii), 其余的情形类似.

当 $0 \leq \rho \leq (\frac{1}{3})\delta$ 时. 由中值定理和关系式 (4.7), (5.9), (6.7) 对于 ε, μ 足够地小, 存在正常数 D_{i3} ($i = 1, 2, \dots, m$), 使得

$$\begin{aligned} & \mu \frac{\partial \alpha_i}{\partial t} - \varepsilon^2 L \alpha_i + T \alpha_i - f_i(t, x, \alpha_i) \\ = & \mu \frac{\partial Y_i}{\partial t} - \varepsilon^2 L Y_i + T Y_i - f_i(t, x, Y_i) + [f_i(t, x, Y_i) - f_i(t, x, Y_i - r_i \lambda)] \\ \leq & [T U_{i00} - f_i(t, x, U_{i00})] \\ & + \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N [T U_{ijk}(t, x) - f_{iu_i}(t, x, U_{i00}) U_{ijk} - F_{ijk}] \varepsilon^j \mu^k \\ & + [T z_{i00} - f_i(t, x, z_{i00})] + \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N [T z_{ijk} - \bar{F}_{ijk}] \varepsilon_j \mu^k \\ & + [K_0 v_{i00} - T v_{i00} + f_i(t, \rho, \phi, U_{i00} + z_{i00} + v_{i00}) - f_i(t, \rho, \phi, U_{i00} + z_{i00})] \\ & + \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N [K_0 v_{ijk} - T v_{ijk} + [f_i(t, \rho, \phi, U_{i00} + z_{i00} + v_{i00}) \\ & - f_i(t, \rho, \phi, U_{i00} + z_{i00})]] \varepsilon^j \mu^k \\ & + \left[\frac{\partial w_{i00}}{\partial \tau} + T w_{i00} - f_i(0, x, U_{i00} + z_{i00} + v_{i00} + w_{i00}) - f_i(0, x, U_{i00} + z_{i00} + v_{i00}) \right] \\ & + \sum_{j=0}^M \sum_{\substack{k=0 \\ j+k \neq 0}}^N \left[\frac{\partial w_{ijk}}{\partial \tau} - f_{iu_i}(0, x, U_{i00} + Z_{i00} + V_{i00} + W_{i00}) w_{ijk} - \tilde{F}_{ijk} \right] \varepsilon^j \mu^k \\ & + [D_{i3} - (M + N)r_i] \lambda \leq [D_{i3} - (M + N)r_i] \lambda. \end{aligned}$$

最后, 选取 r_i ($i = 1, 2, \dots, m$) 足够大, 使得 $r_i \geq \frac{D_{i3}}{M+N}$, 这时便证明了不等式 (7.6). 同理, 不等式 (7.7) 也成立. 由 (7.2)–(7.7) 及比较定理 (定理 2.1), 得到 $\alpha_i \leq u_i \leq \beta_i$ ($i = 1, 2, \dots, m$), $(t, x) \in [0, T_0 \times [\Omega + \partial\Omega]]$. 于是, 由 (7.1), 有 (6.8). 定理 7.1 证毕.

参 考 文 献

- [1] de Jager E M, Jiang Furu. The theory of singular perturbation [M]. Amsterdam: North-Holland Publishing Co, 1996.
- [2] Barbu L, Morosanu G. Singularly perturbed boundary-value problems [M]. Basel: Birkhauserm Verlag AG, 2007.
- [3] Pao, C V. Nonlinear parabolic and elliptic equations [M]. New York: Plenum Press, 1992.
- [4] Martinez S, Wolanski N. A singular perturbation problem for a quasi-linear operator satisfying the natural condition of Lieberman [J]. *SIAM J Math Anal*, 2009, 41(1): 318–359.

- [5] Kellogg R B, Kopteva N A. Singularly perturbed semilinear reaction-diffusion problem in a polygonal domain [J]. *J Differ Equations*, 2010, 248(1):184–208.
- [6] Tian Canrong, Zhu Peng. Existence and asymptotic behavior of solutions for quasilinear parabolic systems [J]. *Acta Appl Math*, 2012, 121(1):157–173.
- [7] Skrynnikov Y. Solving initial value problem by matching asymptotic expansions [J]. *SIAM J Appl Math*, 2012, 72(1):405–416.
- [8] Samusenko P F. Asymptotic integration of degenerate singularly perturbed systems of parabolic partial differential equations [J]. *J Math Sci*, 2013, 189(5):834–847.
- [9] Mo Jiaqi, Chen Xianfeng. Homotopic mapping method of solitary wave solutions for generalized complex Burgers equation [J]. *Chin Phys B*, 2010, 10(10):100203.
- [10] Mo Jiaqi, Han Xianglin, Chen Songlin. The singularly perturbed nonlocal reaction diffusion system [J]. *Acta Math Sci*, 2002, 22B(4):549–556.
- [11] Mo Jiaqi, Lin Wantao, Wang Hui. Variational iteration solution of a sea-air oscillator model for the ENSO [J]. *Prog Nat Sci*, 2007, 17(2):230–232.
- [12] Mo Jiaqi, Lin Wantao. A class of nonlinear singularly perturbed problems for reaction diffusion equations with boundary perturbation [J]. *Acta Math Appl Sinica*, 2006, 22(1):27–32.
- [13] Mo Jiaqi. A class of singularly perturbed differential-difference reaction diffusion equation [J]. *Adv Math*, 2009, 38(2):227–231.
- [14] Mo Jiaqi. Homotopic mapping solving method for gain fluency of a laser pulse amplifier [J]. *Science in China, Ser G*, 2009, 52(7):1007–1070.
- [15] Mo Jiaqi, Lin Wantao. Asymptotic solution of activator inhibitor systems for nonlinear reaction diffusion equations [J]. *J Sys Sci & Complexity*, 2008, 20(1):119–128.
- [16] Mo Jiaqi. Approximate solution of homotopic mapping to solitary wave for generalized nonlinear KdV system [J]. *Chin Phys Lett*, 2009, 26(1):010204.
- [17] Mo Jiaqi. Singularly perturbed reaction diffusion problem for nonlinear boundary condition with two parameters [J]. *Chin Phys*, 2010, 19(1):010203.
- [18] Mo Jiaqi, Chen Huaijun. The corner layer solution of Robin problem for semilinear equation [J]. *Math Appl*, 2012, 25(1):1–4.
- [19] Mo Jiaqi. Singular perturbation for a class of nonlinear reaction diffusion systems [J]. *Science in China, Ser A*, 1989, 32(11):1306–1315.
- [20] Mo Jiaqi. Homotopic mapping solving method for gain fluency of a laser pulse amplifier [J]. *Science in China, Ser G*, 2009, 59(7):1007–1010.

- [21] 欧阳成, 林万涛, 程荣军, 莫嘉琪. 一类厄尔尼诺海-气时滞振子的渐近解 [J]. 物理学报, 2013, 62(6):060201.
- [22] 欧阳成, 姚静菀, 石兰芳, 莫嘉琪. 一类尘埃等离子体孤波解 [J]. 物理学报, 2014, 63(11):110203.
- [23] Ouyang Cheng, Cheng Lihua, Mo Jiaqi. Solving a class of burning disturbed problem with shock layer [J]. *Chin Phys B*, 2012, 21(5):050203.

Interior Shock Asymptotic Solutions to Nonlinear Reaction Diffusion Integral Differential Equations with Two Parameters

OUYANG Cheng¹ MO Jiaqi²

¹Faculty of Science, Huzhou University, Huzhou 313000, Zhejiang, China.

E-mail: oyc@hutczj.cn

²Department of Mathematics, Anhui Normal University, Wuhu 241003, Anhui,

China. E-mail: mojiaqi@mail.ahnu.edu.cn

Abstract The singular perturbation problem for the nonlinear reaction diffusion integral differential problem with two parameters is considered. By using the singular perturbation method, the outer solution, interior shock layer, boundary layer and initial layer corrective terms are constructed, then the formal asymptotic expansion of solution is obtained. Finally, the uniform validity of asymptotic expansion for solution to this problem is proved by using the comparison theorem for integral differential equation.

Keywords Reaction diffusion, Singular perturbation, Initial boundary value problem

2010 MR Subject Classification 35B25