

ON THE OPTIMAL PARKING PROBLEM***

HE SHENGWU (何声武)* WANG JIAGANG (汪嘉冈)**


(Dedicated to the Tenth Anniversary of CAM)

Abstract

In this paper, the concept of dual predictable projection is used, for the optimal parking problem. A strictly rigorous, simpler treatment is introduced and the optimal stopping rule is also given explicitly.

§1. Introduction and the Main Result

The optimal parking problem, presented by Sakaguchi and Tamaki^[4], is described as follows. A motorist is driving his car along a street toward his destination, and is looking for a parking place. If he finds an unoccupied parking place, he must decide either to park there and to walk the distance to his destination or to continue driving, expecting to find another parking place nearer to his destination. It is assumed that

- 1) The location of the destination T is a random variable, $0 < T < \infty$, with a known distribution function $F(x)$ and finite expectation;
- 2) The unoccupied parking places appear randomly in accordance with a Poisson process with parameter $\lambda > 0$, i. e. if denote the unoccupied  parking places by $0 < T_1 < T_2 < \dots < T_n < \dots$, then $T_1, T_2 - T_1, \dots, T_n - T_{n-1}, \dots$ are i. i. d. and the common distribution is exponential with rate λ ;
- 3) T and $\{T_1, T_2, \dots, T_n, \dots\}$ are independent;
- 4) The speed of walking is 1 and the speed of driving is $\frac{1}{r}$, $0 < r < 1$.

If the car stops at S , then the whole time duration spent to reach the destination is

$$rS + |T - S|.$$

What we observe is the two stochastic processes

$$X_t = \sum_{n=1}^{\infty} 1_{T_n \leq t}, \quad t \geq 0, \quad (1)$$

Manuscript received February 28.

* Department of Mathematical Statistic, East China Normal University, Shanghai, China.

** Institute of Applied Mathematics, East China University of Chemical Technology, Shanghai, China.

*** Projects supported by the National Natural Science Foundation of China.

$$Y_t = 1_{T \leq t}, \quad t \geq 0. \quad (2)$$

i. e. at time t we know if T or T_1, T_2, \dots do appear. Therefore, S should be a stopping time with respect to $(X_t, Y_t)_{t \geq 0}$, and $S = T_N$, where N is random. Obviously, S is a finite stopping time, so-called a stopping rule in [1].

Now the problem is to choose S to minimize the expected time

$$E\{rS + |S - T|\}.$$

Let $C = \inf\{x: F(x) = 1\}$, and

$$\varphi(x) = \frac{1}{1 - F(x)} \int_x^\infty e^{-\lambda(y-x)} F(dy), \quad x \in (0, C).$$

It is easy to see that the function φ is well defined on the interval $(0, C)$, and

$$0 \leq \varphi(x) \leq 1, \quad x \in (0, C). \quad (3)$$

Now our main result can be formulated as follows.

Theorem. Suppose that $\varphi(x)$ is increasing on $(0, C)$, and

$$a = \inf\left\{x: \varphi(x) \geq \frac{1-r}{2}\right\}.$$

Then the optimal stopping rule for the parking problem is

$$S^* = \inf\{T_n: n \geq 1, T_n \geq T \text{ or } T_n \geq a\} \quad (4)$$

We give some simple examples to illustrate the condition assumed in the Theorem.

Example 1. When $F(x)$ is the degenerated distribution function at point $C > 0$, we have

$$\varphi(x) = e^{-\lambda(C-x)}, \quad x \in (0, C),$$

and φ is increasing on $(0, C)$.

Example 2. When $F(x)$ is the uniform distribution function on the interval $(0, C)$, we have

$$\varphi(x) = \frac{1}{\lambda(C-x)} [1 - e^{-\lambda(C-x)}], \quad x \in (0, C).$$

It is easy to verify that φ is increasing on $(0, C)$.

Example 3. When $F(x)$ is the exponential distribution function with rate μ , we have $C = \infty$, and

$$\varphi(x) = \frac{\lambda}{\lambda + \mu}, \quad x \geq 0.$$

By the way, if $C < \infty$, it is not difficult to show

$$\lim_{x \rightarrow C} \varphi(x) = 1. \quad (5)$$

But it $C = \infty$, (5) is not true in general, as indicated in Example 3.

Since the stopping rules we considered have the form $S = T_N$, where N takes integer values, our optimal parking problem essentially is one of discrete time case. And under our assumption about the function φ , it will be reduced to a special monotone case, the optimal stopping rule will be found out directly. We need not

appeal to the general results of the optimal stopping theory, presented in [1]. Our idea of dealing with the problem is somehow different from that in [4]. Even the forms of the results are slightly different. In fact, [4] did not give explicitly the conditions which guarantee the existence of the optimal stopping rule. In our opinion, our treatment is strictly rigorous, simpler and easier to understand. In view of the general theory of stochastic processes, what we use is just the concept of dual predictable projection. Essentially, the idea is the same as in [2].

§ 2. A Special Monotone Case

Let $(\mathcal{F}_n)_{n \geq 1}$ be a discrete time filtration, i. e. an increasing sequence of σ -fields: $\mathcal{F}_1 \subset \mathcal{F}_2 \subset \dots \subset \mathcal{F}_n \subset \dots$, and $(X_n)_{n \geq 1}$ be an adapted sequence of integrable random variables, i. e. for each n , X_n is measurable with respect to \mathcal{F}_n . Set

$$Y_n = E[X_n | \mathcal{F}_{n-1}] - X_{n-1}, \quad n \geq 2.$$

Proposition. Suppose

$$Y_2 \leq Y_3 \leq \dots \leq Y_n \leq \dots \quad (6)$$

and

$$S^* = \inf\{n \geq 1: Y_{n+1} \geq 0\} \quad (7)$$

is finite. Then

$$EX_{S^*} = \inf\{EX_S: S \text{ is a stopping rule}\},$$

i. e. S^* is the optimal stopping rule for $(X_n, \mathcal{F}_n)_{n \geq 1}$.

Proof Let S be a stopping rule. Then

$$\begin{aligned} EX_S &= E \left\{ X_1 + \sum_{k=2}^S (X_k - X_{k-1}) \right\} = E \left\{ X_1 + \sum_{n=2}^{\infty} 1_{S \geq n} \sum_{k=2}^n (X_k - X_{k-1}) \right\} \\ &= E \left\{ X_1 + \sum_{k=2}^{\infty} (X_k - X_{k-1}) 1_{S \geq k} \right\} = E \left\{ X_1 + \sum_{k=2}^{\infty} E[X_k - X_{k-1} | \mathcal{F}_{k-1}] 1_{S \geq k} \right\} \\ &= E \left\{ X_1 + \sum_{k=2}^{\infty} Y_k 1_{S \geq k} \right\} \end{aligned} \quad (8)$$

where we commute the summations to get the third equality and use the fact that $\{S \geq k\} \in \mathcal{F}_{k-1}$, $k \geq 2$, for the fourth one.

Now from (6) and (8) it is easy to see that in order to minimize EX_S for $k \geq 2$ we should make

$$\begin{cases} 1_{S \geq k} = 1, & \text{if } Y_k < 0, \\ 1_{S \geq k} = 0, & \text{if } Y_k \geq 0. \end{cases}$$

Hence, S^* defined by (7) is the optimal stopping rule.

According to [1], the monotone case means that

$$A_n = \{E[X_{n+1} | \mathcal{F}_n] \geq X_1\}, \quad n \geq 1$$

is an increasing sequence of events. Obviously, if the condition (6) is satisfied, i. e. $(Y_n)_{n \geq 2}$ is increasing, so is $A_n = \{Y_{n+1} \geq 0\}$. Hence, the case we consider is just a

special monotone one, in which the stopping rule defined by (7) is always optimal.

On the other hand, in view of the general theory of stochastic processes established by French school, $(X_1 + \sum_{k=2}^{\infty} Y_k)_{n \geq 1}$ is just the dual predictable projection or compensator of $(X_n)_{n \geq 1}$. Compared with [2], it can be clearly realized that the idea of using the concept of dual predictable projection to deal with the optimal stopping problem is the same as in [2]. The Proposition here is also a certain predictable criterion, which is practically useful as we shall see in the following.

§ 3. The Proof of the Main Result

Set

$$\mathcal{F}_t = \sigma\{X_s, Y_s, s \leq t\}, t \geq 0,$$

i. e. $(\mathcal{F}_t)_{t \geq 0}$ is the filtration of observation. Since $(X_t, Y_t)_{t \geq 0}$ is a multivariate point process, from the results in [3] we know that $(\mathcal{F}_t)_{t \geq 0}$ is right continuous and the following Lemma is true.

Lemma 1. For each (\mathcal{F}_t) -stopping time S , we have

$$\mathcal{F}_S = \sigma\{X_{S \wedge t}, Y_{S \wedge t}, t \geq 0\}.$$

We should make decisions at $T_n, n=1, 2, \dots$. What we concern with is the filtration $(\mathcal{F}_{T_n})_{n \geq 1}$ indeed. From Lemma 1 and definitions (1) and (2), the following Lemma can be derived immediately.

Lemma 2. For each $n \geq 1$, we have

$$\mathcal{F}_{T_n} \cap \{T_n < T\} = \sigma\{T_1, \dots, T_n\} \cap \{T_n < T\},$$

$$\mathcal{F}_{T_n} \cap \{T_n \geq T\} = \sigma\{T_1, \dots, T_n, T\} \cap \{T_n \geq T\}.$$

Lemma 3. For every integrable random variable ξ , we have

$$E[\xi | \mathcal{F}_{T_n}] = \frac{E[\xi 1_{T_n < T} | T_1, \dots, T_n]}{E[1_{T_n < T} | T_1, \dots, T_n]} 1_{T_n < T} + E[\xi | T_1, \dots, T_n, T] 1_{T_n \geq T}. \quad (9)$$

Lemma 3 is a direct consequence of Lemma 2. Notice that the event $\{T_n < T\}$ is not measurable with respect to $\{T_1, \dots, T_n\}$, which incurs some complexity in the formula (9). The details are referred to the Appendix.

Now we are ready for the proof of the Theorem. At first, we observe that the objective function has the form

$$\begin{aligned} E\{rS + |S - T|\} &= E\left\{\int_0^S r dt + \int_0^S Y_t dt + \int_S^\infty (1 - Y_t) dt\right\} \\ &= E\left\{\int_0^S (r - 1 + 2Y_t) dt + \int_0^\infty (1 - Y_t) dt\right\} \\ &= E\left\{\int_0^S (r - 1 + 2Y_t) dt\right\} + ET. \end{aligned} \quad (10)$$

Secondly, we calculate

$$Z_n = E \left[\int_{T_n}^{T_{n+1}} (r-1+21_{T \leq t}) dt \mid \mathcal{F}_{T_n} \right], \quad n \geq 1$$

by formula (9). Noting that T , $T_{n+1}-T_n$ and $\{T_1, \dots, T_n\}$ are independent, we have

$$\begin{aligned} Z_{T_n} 1_{T_n > T} &= E \left[\int_{T_n}^{T_{n+1}} (r-1+21_{T \leq t}) dt 1_{T_n > T} \mid T_1, \dots, T_n, T \right] \\ &= E[(r+1)(T_{n+1}-T_n) \mid T_1, \dots, T_n, T] 1_{T_n > T} = \frac{r+1}{\lambda} 1_{T_n > T}, \end{aligned} \quad (11)$$

$$Z_{T_n} 1_{T_n < T} = \frac{E[\{(r-1)(T_{n+1}-T_n) + 2(T_{n+1}-T)^+\} 1_{T_n < T} \mid T_1, \dots, T_n]}{E[1_{T_n < T} \mid T_1, \dots, T_n]} 1_{T_n < T}. \quad (12)$$

On the other hand

$$\begin{aligned} E[(T_{n+1}-T_n) 1_{T_n < T} \mid T_1, \dots, T_n] &= E[1_{T_n < T} E[T_{n+1}-T_n \mid T_1, \dots, T_n, T] \mid T_1, \dots, T_n] \\ &= \frac{1}{\lambda} E[1_{T_n < T} \mid T_1, \dots, T_n] \end{aligned} \quad (13)$$

$$E[1_{T_n < T} \mid T_1, \dots, T_n] = E[1_{x < T} \mid T_1, \dots, T_n] \big|_{x=T_n} = 1 - F(T_n) \quad (14)$$

$$\begin{aligned} E[(T_{n+1}-T)^+ 1_{T_n < T} \mid T_1, \dots, T_n] &= E[T_{n+1}-T_n + z - T)^+ 1_{z < T} \mid T_1, \dots, T_n] \big|_{z=T_n} \\ &= \int_{T_n}^{\infty} F(dy) \int_0^{\infty} (x+z-y)^+ \lambda e^{-\lambda x} dx \big|_{z=T_n} \\ &= \frac{1}{\lambda} \int_{T_n}^{\infty} e^{-\lambda(y-T_n)} F(dy). \end{aligned} \quad (15)$$

Substituting (13), (15) into (12) yields

$$Z_n 1_{T_n < T} = \frac{1}{\lambda} [r-1+2\varphi(T_n)] 1_{T_n < T}. \quad (16)$$

Combining (16) with (11) gives

$$Z_n = \frac{r+1}{\lambda} 1_{T \leq T_n} + \frac{1}{\lambda} [r-1+2\varphi(T_n)] 1_{T_n < T}. \quad (17)$$

Observing that from (3) and our assumption on φ we have

$$\begin{aligned} \{T \leq T_n\} &\subset \{T \leq T_{n+1}\}, \\ r-1+2\varphi(T_n) &\leq r+1, \text{ on } \{T_n < T \leq T_{n+1}\}, \\ T_n \leq T_{n+1} < T \leq C, \varphi(T_n) &\leq \varphi(T_{n+1}), \text{ on } \{T_{n+1} < T\}, \end{aligned}$$

we conclude that

$$Z_n \leq Z_{n+1}, \quad n \geq 1.$$

Since $S = T_N$, by using the Proposition in section 2, we obtain the optimal stopping rule

$$S^* = \inf\{T_n: n \geq 1, Z_n \geq 0\}. \quad (18)$$

From (17) and (18) we know immediately that S^* has the form (4).

Appendix

Let (Ω, \mathcal{F}, P) be a probability space, \mathcal{G} a sub- σ -field of \mathcal{F} , and ξ an integrable random variable. Then for every $A \in \mathcal{F}$ and $G \in \mathcal{G}$, we have

$$\int_{AG} \xi dP = \int_{AG} \frac{E[\xi 1_A | \mathcal{G}]}{E[1_A | \mathcal{G}]} dP. \quad (19)$$

Proof Set $H = \{E[1_A | \mathcal{G}] \neq 0\}$, then $H \in \mathcal{G}$ and

$$P(AH^c) = \int_{H^c} E[1_A | \mathcal{G}] dP = 0,$$

where $H^c = \{E[1_A | \mathcal{G}] = 0\}$ is the complement of H . Hence, the right side of (9) makes sense. Define

$$P_A(F) = P(AF), \quad F \in \mathcal{F}.$$

Then for every $G \in \mathcal{G}$ we have

$$\begin{aligned} P_A(G) &= \int_G E[1_A | \mathcal{G}] dP \\ \int_{AG} \frac{E[\xi 1_A | \mathcal{G}]}{E[1_A | \mathcal{G}]} dP &= \int_{AGH} \frac{E[\xi 1_A | \mathcal{G}]}{E[1_A | \mathcal{G}]} dP = \int_{GH} \frac{E[\xi 1_A | \mathcal{G}]}{E[1_A | \mathcal{G}]} dP_A \\ &= \int_{GH} E[\xi 1_A | \mathcal{G}] dP = \int_{GH} \xi 1_A dP = \int_{AG} \xi dP. \end{aligned}$$

(19) follows.

References

- [1] Chow, Y. S., Robbins, H. & Siegmund, D., Great Expectation: the Theory of Optimal Stopping, Houghton Mifflin Company, Boston, 1971.
- [2] He, S. W., Optimization applications of compensators of Poisson random measures, *Probability in Engineering and Information Science*, **3** (1989), 149—155.
- [3] Jacod, J., Multivariate point processes: predictable projection, Radon-Nikodym derivative, representation of martingales, *Z. Wahrsch. Verw. Geb.*, **31** (1975), 235—253.
- [4] Sakaguchi, M. & Tamaki, M., On the optimal parking problem in which spaces appear randomly, *Bulletin of Informatics and Cybernetics*, **25** (1982), 1—10.