

Global Existence of Strong Solutions of Navier-Stokes-Poisson Equations for One-Dimensional Isentropic Compressible Fluids**

Junping YIN* Zhong TAN*

Abstract The authors prove two global existence results of strong solutions of the isentropic compressible Navier-Stokes-Poisson equations in one-dimensional bounded intervals. The first result shows only the existence. And the second one shows the existence and uniqueness result based on the first result, but the uniqueness requires some compatibility condition. In this paper the initial vacuum is allowed, and T is bounded.

Keywords Global strong solutions, Navier-Stokes-Poisson equations, Existence and uniqueness

2000 MR Subject Classification 35A05, 35Q30

1 Introduction

In this paper, we consider the system:

$$\begin{cases} \rho_t + (\rho u)_x = 0, & \text{in } (0, T) \times \Omega, \\ (\rho u)_t + (\rho u^2)_x + \rho \Phi_x - \lambda u_{xx} + p_x = \rho f, & \text{in } (0, T) \times \Omega, \\ \Phi_{xx} = 4\pi g \left(\rho - \frac{1}{|\Omega|} \int_{\Omega} \rho dx \right), & \text{in } (0, T) \times \Omega, \end{cases}$$

where $p = a\rho^\gamma$ ($a > 0$, $\gamma > 1$), $\lambda > 0$. In this paper, we only consider that Ω is a one-dimensional bounded interval. For simplicity we only consider $\Omega = (0, 1)$, $T < +\infty$. The initial and boundary conditions are

$$\rho|_{t=0} = \rho_0(x) \geq 0, \quad u|_{t=0} = u_0, \quad \forall x \in (0, 1), \quad (1.4)$$

$$u(0, t) = u(1, t) = 0, \quad \Phi(0, t) = \Phi(1, t) = 0, \quad \forall t > 0. \quad (1.5)$$

For the vacuum case, in [1], Takayuki Kobayashi and Takashi Suzuki proved the existence of weak solution to Navier-Stokes-Poisson equations. Their methods are similar to Feireisl's methods (see [2]). But as for Navier-Stokes-Poisson systems, the results of strong solutions are few. However, as for the existence, uniqueness, or other virtues of the strong solutions of Navier-Stokes equations, we may refer to [3–10].

Manuscript received July 4, 2006. Revised November 8, 2007. Published online June 24, 2008.

*Department of Mathematics, Xiamen University, Xiamen 361005, Fujian, China.

E-mail: yinjp829829@126.com

**Project supported by the National Natural Science Foundation of China (No. 10531020) and the Program of 985 Innovation Engineering on Information in Xiamen University (2004–2007) and the New Century Excellent Talents in Xiamen University.

Remark 1.1 The problem about the radially solutions of the Navier-Stokes-Poisson equations is worthy to consider.

1.1 Main results

Theorem 1.1 Assume that the initial conditions and f satisfy

$$\rho_0 \in H^1(0, 1), \quad u_0 \in H_0^1(0, 1), \quad f \in L_{\text{loc}}^2(0, \infty; L^{\frac{2\gamma}{\gamma-1}}(0, 1)). \quad (1.6)$$

Then there is a global strong solution (ρ, u, Φ) of (1.1)–(1.5), such that for all $T \in (0, \infty)$, we have

$$\begin{aligned} \rho &\in L^\infty(0, T; H^1(0, 1)), \quad \rho_t \in L^\infty(0, T; L^2(0, 1)), \\ \Phi &\in L^\infty(0, T; H^3(0, 1)), \quad \Phi_t \in L^\infty(0, T; H^2(0, 1)), \\ u &\in L^\infty(0, T; H_0^1(0, 1)), \quad (\rho u)_t, u_{xx} \in L^2(0, T; L^2(0, 1)). \end{aligned}$$

Theorem 1.2 Assume that the initial conditions and f satisfy

$$\begin{aligned} \rho_0 &\in H^1(0, 1), \quad u_0 \in H_0^1(0, 1) \cap H^2(0, 1), \\ f &\in L_{\text{loc}}^2(0, \infty; L^{\frac{2\gamma}{\gamma-1}}(0, 1)), \quad f_x, f_t \in L_{\text{loc}}^2(0, \infty; L^2(0, 1)) \end{aligned} \quad (1.7)$$

and compatibility condition

$$\lambda(u_0)_{xx} - (a\rho_0^\gamma)_x = \rho^{\frac{1}{2}}g \quad \text{for some } g \in L^2(0, 1). \quad (1.8)$$

Then there is a unique strong solution (ρ, u, Φ) , such that for all $T \in (0, \infty)$, we have

$$\begin{aligned} \rho &\in L^\infty(0, T; H^1(0, 1)), \quad u \in L^\infty(0, T; H^2(0, 1)), \quad \Phi \in L^\infty(0, T; H^3(0, 1)), \\ \rho_t, \sqrt{\rho}u_t &\in L^\infty(0, T; L^2(0, 1)), \quad u_t, G_x \in L^2(0, T; H^1(0, 1)), \quad \Phi_t \in L^\infty(0, T; H^2(0, 1)). \end{aligned}$$

Remark 1.2 When $G = \lambda u_x - p$ is an effective flux, we can easily get $\rho \in C([0, T] \times (0, 1))$, $u \in C([0, T]; H_0^1(0, 1))$, $\Phi \in C([0, T]; H^2(0, 1))$, and combining the equations (1.1)–(1.3) and the effective viscous flux, we may obtain other regularity.

2 A priori Estimates for Smooth Solutions

To get the existence of strong solutions, obviously, we require some more regular estimates. So we provide that (ρ, u, Φ) is a smooth solution of (1.1)–(1.5), $\rho > 0$, and $T \in (0, \infty)$ is some fixed time. Moreover we may let $m_0 := \int_0^1 \rho_0(x)dx$ be initial mass and $m_0 > 0$. To simplify, we let $\lambda \equiv 1$. In fact, as we can deal with approximate system, we only consider initial nonvacuum. Combining the classical results of (1.3) with our correlated uniform estimates, we may get the existence of strong solutions of our system. To prove uniqueness, we use the classical method.

Lemma 2.1

$$\sup_{0 \leq t \leq T} \int_0^1 (\rho u^2 + p)dx + \int_0^T \int_0^1 u_x^2 dx dt \leq C, \quad (2.1)$$

where C is dependent on $|\rho_0|_{H^1(0, 1)}$, $|u_0|_{H_0^1(0, 1)}$ and $|f|_{L^2(0, T; L^{\frac{2\gamma}{\gamma-1}}(0, 1))}$, but is independent of the lower bound of ρ_0 .

Proof Firstly, we introduce the energy formula

$$\begin{aligned} E(t) &= \int_0^1 \left(\frac{1}{2} \rho |u|^2 + \frac{a}{\gamma-1} \rho^\gamma \right) dx - \frac{1}{8\pi g} \int_0^1 |\Phi_x|^2 dx, \\ E(0) &= \int_0^1 \left(\frac{1}{2} \rho_0 |u_0|^2 + \frac{a}{\gamma-1} \rho_0^\gamma \right) dx - \frac{1}{8\pi g} \int_0^1 |\Phi_{0x}|^2 dx, \end{aligned}$$

where $E(0)$ is the initial energy. It follows from (1.1) and (1.2) that

$$\rho u_t + (\rho u) u_x + \rho \Phi_x - \lambda u_{xx} + P_x = \rho f.$$

Multiplying this equation by u , integrating (by parts) over $(0, 1)$, and combining the equations (1.1) and (1.3), we can deal with each term as follows:

$$\begin{aligned} \int_0^1 \rho u_t u dx + \int_0^1 (\rho u) u_x u dx &= \frac{1}{2} \frac{d}{dt} \int_0^1 \rho u^2 dx, \\ \int_0^1 \rho \Phi_x u dx &= - \int_0^1 (\rho u)_x \Phi dx = \int_0^1 \rho_t \Phi dx = \frac{1}{4\pi g} \int_0^1 \Phi_{xxt} \Phi dx = - \frac{1}{8\pi g} \frac{d}{dt} \int_0^1 \Phi_x^2 dx, \\ \int_0^1 a\gamma\rho^{\gamma-1} \rho_x u dx &= \int_0^1 a\gamma\rho^{\gamma-1} (-\rho_t - \rho u_x) dx = - \frac{d}{dt} \int_0^1 a\rho^\gamma dx + \gamma \int_0^1 P_x u dx. \end{aligned}$$

Then

$$\int_0^1 P_x u dx = \frac{a}{\gamma-1} \frac{d}{dt} \int_0^1 \rho^\gamma dx.$$

Combining these estimates, we can conclude that

$$\frac{dE(t)}{dt} + \lambda |u_x|_{L^2(0,1)}^2 \leq \int_0^1 \rho u f dx dt. \quad (2.2)$$

We deal with $\int_0^1 |\Phi_x|^2 dx$ of $E(t)$. Multiplying (1.3) by Φ and integrating over $(0, 1)$, we get

$$\int_0^1 \Phi_{xx} \Phi dx = 4\pi g \left(\int_0^1 \rho \Phi dx - m_0 \int_0^1 \Phi dx \right) \quad (2.3)$$

and

$$\begin{aligned} 4\pi g \left(\int_0^1 \rho |\Phi| dx - m_0 \int_0^1 \Phi dx \right) &\leq 8\pi g m_0 |\Phi|_{L^\infty(0,1)} \leq 8\pi g m_0 |\Phi_x|_{L^2(0,1)} \\ &\leq \frac{1}{2} |\Phi_x|_{L^2(0,1)}^2 + 32\pi^2 g^2 m_0^2. \end{aligned}$$

Consequently,

$$\int_0^1 \Phi_x^2 dx \leq C \left(\int_0^1 \rho dx \right)^2 \leq C(m_0). \quad (2.4)$$

Integrating (2.2) over $(0, t)$, we have

$$E(t) + \lambda \int_0^t |u_x|^2 ds \leq E(0) + \int_0^t \int_0^1 \rho |u| |f| dx ds. \quad (2.5)$$

Combining (2.4) and the form of $E(t)$, we obtain

$$\begin{aligned}
& \frac{1}{2}|\sqrt{\rho}u(t)|_{L^2(0,1)}^2 + \frac{a}{4(\gamma-1)}|\rho|_{L^\gamma(0,1)}^\gamma + \lambda \int_0^t u_x^2 dx \\
& \leq C + \int_0^t \int_0^1 \rho |u| |f| dx ds \\
& \leq C + C \int_0^t |f|_{L^{\frac{2\gamma}{\gamma-1}}(0,1)} |\rho|_{L^\gamma(0,1)}^{\frac{1}{2}} |\sqrt{\rho}u|_{L^2(0,1)} ds \\
& \leq C + C \int_0^t |f|_{L^{\frac{2\gamma}{\gamma-1}}(0,1)} (1 + |\rho|_{L^\gamma(0,1)}^\gamma + |\sqrt{\rho}u|_{L^2(0,1)}^2) ds \\
& \leq C + C \int_0^t |f|_{L^{\frac{2\gamma}{\gamma-1}}(0,1)} ds + C \int_0^t |f|_{L^{\frac{2\gamma}{\gamma-1}}(0,1)} (|\rho|_{L^\gamma(0,1)}^\gamma + |\sqrt{\rho}u|_{L^2(0,1)}^2) ds.
\end{aligned}$$

Using Gronwall's inequality, we get

$$\sup_{0 \leq t \leq T} (|\sqrt{\rho}u|_{L^2(0,1)}^2 + |\rho|_{L^\gamma(0,1)}^\gamma) + \lambda \int_0^T \int_0^1 u_x^2 dx dt \leq C(m_0, f), \quad (2.6)$$

where C is independent of the lower bound of ρ_0 .

Lemma 2.2

$$\sup_{0 \leq t \leq T} |\rho(t)|_{L^\infty(0,1)} \leq C, \quad (2.7)$$

where C is dependent on the initial mass m_0 , $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$ and $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$, but is independent of the lower bound of ρ_0 .

Proof Consider Lagrangian flow $X = X(t, x)$ of u and define

$$\begin{cases} \frac{\partial X}{\partial t} = u(t, X(t, x)), \\ X(0, x) = x \in [0, 1]. \end{cases}$$

Then we only require to prove $\rho(t, X(t, x)) \leq C$, for any $(t, x) \in (0, T] \times (0, 1)$. Let $t_0 \in (0, T]$ be any fixed time. Combining $C^{-1} \leq \int_0^1 \rho_0(x) dx = m_0 \leq C$, the conservative mass, and the means of the Lagrangian flow $X = X(t, x)$, proving by contradiction, we can easily find some $x_1 \in (0, 1)$, such that

$$C^{-1} \leq \rho_0(x_1) \quad \text{and} \quad \rho(t_0, X(t_0, x_1)) \leq C. \quad (2.8)$$

Furthermore, if we take some subset $(a, b) \subset (0, 1)$ such that $C^{-1} \leq \rho_0(x)$, then combining the means of X and the conservative mass, we get

$$\int_{X(t_0, a)}^{X(t_0, b)} \rho(t_0, x) dx = \int_a^b \rho_0(x) dx \leq C.$$

Next, we prove that for any $x_2 \in (0, 1)$, $\rho(t_0, X(t_0, x_2)) \leq C$ holds. Let $X_j(t) := X(t, x_j)$, $j = 1, 2$, and $L(t) = \log \rho(t, X_2(t)) - \log \rho(t, X_1(t))$. Then using (1.1)–(1.3), we get

$$\begin{aligned}
\frac{dL}{dt} &= \frac{1}{\rho(t, X_2(t))} \left(\rho_t(t, X_2(t)) + \rho_x \frac{dX_2}{dt} \right) - \frac{1}{\rho(t, X_1(t))} \left(\rho_t(t, X_1(t)) + \rho_x \frac{dX_1}{dt} \right) \\
&= \frac{1}{\rho(t, X_2(t))} (-\rho_x(t, X_2)u(t, X_2) - \rho(t, X_2)u_x(t, X_2) + \rho_x u(t, X_2))
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\rho(t, X_1(t))}(-\rho_x(t, X_1)u(t, X_1) - \rho(t, X_1)u_x(t, X_1) + \rho_x u(t, X_1)) \\
& = -(u_x(t, X_2) - u_x(t, X_1)) = -\int_{X_1(t)}^{X_2(t)} u_{xx} dx \\
& = -\int_{X_1(t)}^{X_2(t)} [(\rho u)_t + (\rho u^2)_x + p_x + \rho \Phi_x - \rho f] dx \\
& = -\int_{X_1(t)}^{X_2(t)} [-(\rho u)_x u + \rho u_t + (\rho u)_x u + \rho u u_x + p_x + \rho \Phi_x - \rho f] dx \\
& = -\int_{X_1(t)}^{X_2(t)} (\rho u_t + \rho u u_x + p_x + \rho \Phi_x - \rho f) dx. \tag{2.9}
\end{aligned}$$

Let $U(t) = \int_{X_1(t)}^{X_2(t)} \rho u(t, x) dx$. Then

$$\begin{aligned}
\frac{dU(t)}{dt} &= \rho u(t, X_2(t)) \frac{dX_2}{dt} - \rho u(t, X_1(t)) \frac{dX_1}{dt} + \int_{X_1(t)}^{X_2(t)} (\rho u(t, x))_t dx \\
&= \rho u^2(t, X_2(t)) - \rho u^2(t, X_1(t)) + \int_{X_1(t)}^{X_2(t)} [-(\rho u)_x u + \rho u_t] dx \\
&= \int_{X_1(t)}^{X_2(t)} \rho u(t, x) u_x(t, x) dx + \int_{X_1(t)}^{X_2(t)} \rho u_t(t, x) dx. \tag{2.10}
\end{aligned}$$

Substituting (2.10) into (2.9), we have

$$\frac{dL(t)}{dt} + \frac{dU(t)}{dt} = - \left(\int_{X_1(t)}^{X_2(t)} p_x dx + \int_{X_1(t)}^{X_2(t)} \rho \Phi_x dx - \int_{X_1(t)}^{X_2(t)} \rho f dx \right). \tag{2.11}$$

Let $\alpha(t) = \frac{p(\rho(t, X_2)) - p(\rho(t, X_1))}{L(t)}$. We easily find $\alpha(t) > 0$. Thus (2.11) becomes

$$\frac{dL(t)}{dt} + \frac{dU(t)}{dt} = -\alpha(L + U) + \alpha U - \int_{X_1(t)}^{X_2(t)} (\rho \Phi_x - \rho f) dx. \tag{2.12}$$

Using the theorem of ODE, we get

$$L(t) + U(t) = e^{-\int_0^t \alpha(s) ds} (L(0) + U(0)) + \int_0^t e^{-\int_s^t \alpha(\tau) d\tau} (\alpha(s) U(s) - \Psi(s)) ds,$$

where $\Psi(s) = \int_{X_2(s)}^{X_1(s)} (\rho \Phi_x - \rho f) dx$. Combining $L(0) \leq C$ ($|\rho_0(x)|_{L^\infty(0,1)} \leq C$), we have

$$L(t) \leq C + |U(0)| + |U(t)| + \int_0^t e^{-\int_s^t \alpha(\tau) d\tau} (\alpha(s) |U(s)| + |\Psi(s)|) ds.$$

Particularly, we let $t = t_0$. Thus

$$L(t_0) = C + |U(0)| + \sup_{0 \leq t \leq t_0} |U(t)| + \int_0^{t_0} |\Psi(s)| ds. \tag{2.13}$$

We deal with the latter terms as follows:

$$\begin{aligned}
|U(0)| &\leq \int_{x_1}^{x_2} \rho_0 |u(0, x)| dx \leq C, \\
|U(t)| &\leq \int_{X_1(t)}^{X_2(t)} \rho |u(t, x)| dx \leq \left(\int_{X_1(t)}^{X_2(t)} \rho dx \right)^{\frac{1}{2}} \left(\int_{X_1(t)}^{X_2(t)} \rho u^2 dx \right)^{\frac{1}{2}}. \tag{2.14}
\end{aligned}$$

Consequently,

$$\sup_{0 \leq t \leq t_0} |U(t)| \leq \sup_{0 \leq t \leq t_0} \left(\int_0^1 \rho dx \right)^{\frac{1}{2}} \left(\int_0^1 \rho u^2 dx \right)^{\frac{1}{2}}.$$

Combining (2.6), we get

$$\begin{aligned} & \sup_{0 \leq t \leq t_0} |U(t)| \leq C, \\ & \int_0^t \int_{X_1(s)}^{X_2(s)} \rho |f| dx ds \leq C \int_0^t \left(\int_0^1 \rho^\gamma dx \right)^{\frac{1}{\gamma}} \left(\int_0^1 |f|^{\frac{2\gamma}{\gamma-1}} dx \right)^{\frac{\gamma-1}{2\gamma}} ds \leq C(T), \\ & \int_0^t \int_{X_1(s)}^{X_2(s)} \rho |\Phi_x| dx ds \leq C \int_0^t \left(\int_0^1 \rho^\gamma dx \right)^{\frac{1}{\gamma}} \left(\int_0^1 |\Phi_x|^{\frac{2\gamma}{\gamma-1}} dx \right)^{\frac{\gamma-1}{2\gamma}} ds. \end{aligned} \quad (2.15)$$

Combining imbedded theorem and the estimates of Poisson equation, we get

$$\int_0^t \int_{X_1(s)}^{X_2(s)} \rho |\Phi_x| dx ds \leq C \int_0^t \int_0^1 |\Phi_{xx}| dx ds \leq C(m_0, T). \quad (2.16)$$

From (2.13)–(2.16), we have $L(t_0) \leq C$. Then

$$\log \rho(t_0, X(t_0, x_2)) = \log \rho(t_0, X(t_0, x_1)) + L(t_0) \leq C. \quad (2.17)$$

Because t_0, x_2 are arbitrary, Lemma 2.2 is proved.

To get the higher estimates, the effective viscous flux is very important. In the proof below, usually we will use the regularity of $G_x = u_{xx} - p_x$.

Lemma 2.3

$$\sup_{0 \leq t \leq T} (|u|_{L^\infty(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T |\sqrt{\rho} u_t|_{L^2(0,1)}^2 dt \leq C, \quad (2.18)$$

where C is dependent on $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$ and $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$, but is independent of the lower bound of ρ_0 .

Proof Multiplying (1.2) by u_t and integrating over $(0, 1)$, we get

$$\int_0^1 \rho u_t^2 dx + \int_0^1 \rho u u_x u_t dx + \int_0^1 \rho \Phi_x u_t dx - \lambda \int_0^1 u_{xx} u_t dx = \int_0^1 \rho f u_t dx - \int_0^1 p_x u_t dx.$$

Thus

$$\int_0^1 \rho u_t^2 dx + \frac{d}{dt} \int_0^1 \frac{1}{2} u_x^2 dx \leq C \left(\int_0^1 \rho u^2 u_x^2 dx + \int_0^1 \Phi_x^2 dx + \int_0^1 p u_{xt} dx + \int_0^1 \rho |f|^2 dx \right). \quad (2.19)$$

Next, we deal with each of the above terms as follows:

$$\begin{aligned} \int_0^1 p u_{xt} dx &= \frac{d}{dt} \int_0^1 p u_x dx - \int_0^1 p_t u_{xt} dx = \frac{d}{dt} \int_0^1 p u_x dx - \int_0^1 a \gamma \rho^{\gamma-1} \rho_t u_{xt} dx \\ &= \frac{d}{dt} \int_0^1 p u_x dx - \int_0^1 p u u_{xx} dx + (\gamma-1) \int_0^1 p u_x^2 dx \\ &= \frac{d}{dt} \int_0^1 p u_x dx - \int_0^1 p u (G_x + p_x) dx + (\gamma-1) \int_0^1 p (G + p)^2 dx, \end{aligned} \quad (2.20)$$

where

$$\begin{aligned} - \int_0^1 p u p_x dx &= \frac{-1}{2(2\gamma-1)} \frac{d}{dt} \int_0^1 p^2 dx; \\ (\gamma-1) \int_0^1 p u_x^2 dx &= (\gamma-1) \int_0^1 p(G+p)^2 dx \\ &= (\gamma-1) \int_0^1 p G^2 dx + 4(\gamma-1) \int_0^1 p p_x u dx - (\gamma-1) \int_0^1 p^3 dx. \end{aligned}$$

Then, (2.20) becomes

$$\begin{aligned} \int_0^1 p u_{xt} dx &= \frac{d}{dt} \int_0^1 p u_x dx - \frac{d}{dt} \int_0^1 \frac{4\gamma-3}{2(2\gamma-1)} p^2 dx \\ &\quad + (\gamma-1) \int_0^1 p(G^2 - p^2) dx - \int_0^1 p u G_x dx. \end{aligned} \quad (2.21)$$

Substituting (2.21) into (2.19) and integrating over $(0, t)$, we get

$$\begin{aligned} &\int_0^t \int_0^1 \rho u_t^2 dx ds + \int_0^1 \frac{1}{2} u_x^2(t) dx - \int_0^1 \frac{1}{2} u_x^2(0) dx \\ &\leq C + \int_0^t \int_0^1 (\rho u^2 u_x^2 + p G^2 + p|u||G_x|) dx ds + \int_0^t \int_0^1 (\Phi_x^2 + \rho f^2) dx ds \\ &\quad + \int_0^1 (p u_x)(t) dx - \int_0^1 (p u_x)(0) dx - \int_0^1 \frac{4\gamma-3}{2(2\gamma-1)} p^2(t) dx \\ &\quad + \int_0^1 \frac{4\gamma-3}{2(2\gamma-1)} p^2(0) dx - (\gamma-1) \int_0^t \int_0^1 p^3 dx ds \\ &\leq C + \int_0^t \int_0^1 (\rho u^2 u_x^2 + p G^2 + p|u||G_x|) dx ds. \end{aligned} \quad (2.22)$$

Now, we deal with each of the right terms of (2.22) as follows:

$$\begin{aligned} \int_0^t \int_0^1 \rho u^2 u_x^2 dx ds &\leq \int_0^t |\rho|_{L^\infty(0,1)} |u|_{L^\infty(0,1)}^2 |u_x|_{L^2(0,1)}^2 ds \leq \int_0^t |u_x|_{L^2(0,1)}^4 ds, \\ \int_0^t \int_0^1 p G^2 dx ds &\leq 2 \int_0^t \int_0^1 p(u_x^2 + p^2) dx ds \leq C, \\ \int_0^t \int_0^1 p|u||G_x| dx ds &\leq C \int_0^t |\rho|_{L^\infty(0,1)}^{\gamma-\frac{1}{2}} |\sqrt{\rho} u|_{L^2(0,1)} |G_x|_{L^2(0,1)} ds \leq C \int_0^t |G_x|_{L^2(0,1)} ds, \end{aligned} \quad (2.23)$$

and

$$G_x = u_{xx} - p_x = \rho u_t + \rho u u_x + \rho \Phi_x - \rho f.$$

Consequently,

$$|G_x|_{L^2(0,1)} \leq C(|\sqrt{\rho} u_t|_{L^2(0,1)} + |u u_x|_{L^2(0,1)} + |\Phi_x|_{L^2(0,1)} + |\rho f|_{L^2(0,1)}).$$

Then (2.23) becomes

$$\begin{aligned} \int_0^t \int_0^1 p|u||G_x| dx ds &\leq C \int_0^t (|\sqrt{\rho} u_t|_{L^2(0,1)} + |u u_x|_{L^2(0,1)} + |\Phi_x|_{L^2(0,1)} + |\rho f|_{L^2(0,1)}) ds \\ &\leq C + C \int_0^t |u_x|_{L^2(0,1)}^4 ds + \frac{1}{2} \int_0^t \int_0^1 \rho u_t^2 dx ds. \end{aligned}$$

Combining the above estimates, we get

$$\int_0^t (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_x|_{L^2(0,1)}^2) ds \leq C + C \int_0^t |u_x|_{L^2(0,1)}^4 ds.$$

Using Gronwall's inequality, we obtain

$$\int_0^t (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_x|_{L^2(0,1)}^2) ds \leq C.$$

Combining $u|_{\partial\Omega} = 0$ and embedding theorem, we have

$$\sup_{0 \leq t \leq T} (|u|_{L^\infty(0,1)} + |u_x|_{L^2(0,1)}^2) + \int_0^t |\sqrt{\rho} u_t|_{L^2(0,1)}^2 dt \leq C.$$

Lemma 2.4

$$\int_0^T (|u_x|_{L^\infty(0,1)}^2 + |G_x|_{L^2(0,1)}^2) dt \leq C, \quad (2.24)$$

where C is dependent on $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$ and $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$, but is independent of the lower bound of ρ_0 .

Proof

$$|u_x|_{L^\infty(0,1)}^2 \leq 2(|G|_{L^\infty(0,1)}^2 + |p|_{L^\infty(0,1)}^2) \leq 2(|G|_{L^2(0,1)}^2 + |G_x|_{L^2(0,1)}^2 + |p|_{L^\infty(0,1)}^2),$$

and

$$\begin{aligned} \int_0^T |G|_{L^2(0,1)}^2 dt &\leq 2 \int_0^T \int_0^1 |u_x|^2 dx dt + \int_0^T \int_0^1 |p|^2 dx dt \leq C, \\ \int_0^T |G_x|_{L^2(0,1)}^2 dt &\leq C \int_0^T (C + |\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u u_x|_{L^2(0,1)}^2 |\sqrt{\rho} f|_{L^2(0,1)}^2) ds \\ &\leq C + C \int_0^T |u_x|_{L^2(0,1)}^2 ds \leq C. \end{aligned}$$

From the above estimates, we have

$$\sup_{0 \leq t \leq T} (|\rho|_{L^\infty(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_x|_{L^\infty(0,1)}^2 + |G_x|_{L^2(0,1)}^2) dt \leq C, \quad (2.25)$$

where C is dependent on $|\rho|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$ and $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$.

Lemma 2.5

$$\sup_{0 \leq t \leq T} |\rho_x|_{L^2(0,1)} \leq C(T), \quad (2.26)$$

where C is dependent on $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$, $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$ and T , but is independent of the lower bound of ρ_0 .

Proof Differentiating (1.1) with respect to x gives

$$(\rho_x)_t + (\rho u)_{xx} = 0.$$

Multiplying it by ρ_x and integrating over $(0, 1)$, we get

$$\begin{aligned} & \int_0^1 (\rho_x)_t \rho_x dx + \int_0^1 (\rho u)_{xx} \rho_x dx = 0, \\ & \frac{d}{dt} \frac{1}{2} \int_0^1 (\rho_x)^2 dx + \int_0^1 [(\rho_x u) + \rho u_x]_x \rho_x dx = 0, \\ & \int_0^1 (\rho_x u)_x \rho_x dx = \int_0^1 (\rho_{xx} u + \rho_x u_x) \rho_x dx = \int_0^1 \rho_{xx} \rho_x u dx + \int_0^1 \rho_x^2 u_x dx. \end{aligned} \quad (2.27)$$

However

$$\int_0^1 \rho_{xx} \rho_x u dx = - \int_0^1 \rho_x (\rho_x u)_x dx = - \int_0^1 \rho_x \rho_{xx} u dx - \int_0^1 \rho_x^2 u_x dx.$$

Thus

$$\int_0^1 \rho_{xx} \rho_x u dx = - \frac{1}{2} \int_0^1 \rho_x^2 u_x dx,$$

but

$$\begin{aligned} \int_0^1 \rho_x^2 u_x dx & \leq |u_x|_{L^\infty(0,1)} \int_0^1 \rho_x^2 dx, \\ \int_0^1 (\rho u_x)_x \rho_x dx & = \int_0^1 \rho_x^2 u_x dx + \int_0^1 \rho u_{xx} \rho_x dx \\ & \leq |u_x|_{L^\infty(0,1)} \int_0^1 \rho_x^2 dx + \int_0^1 \rho (G_x + p_x) \rho_x dx \\ & \leq C \left(|u_x|_{L^\infty(0,1)} \int_0^1 \rho_x^2 dx + \int_0^1 G_x^2 dx \int_0^1 \rho_x^2 dx + \int_0^1 a\gamma \rho^\gamma \rho_x^2 dx \right) \\ & \leq C \left(|u_x|_{L^\infty(0,1)} \int_0^1 \rho_x^2 dx + \int_0^1 G_x^2 dx \int_0^1 \rho_x^2 dx + \int_0^1 \rho_x^2 dx \right). \end{aligned} \quad (2.28)$$

Combining (2.27) and the above estimates, we get

$$\frac{d}{dt} \frac{1}{2} \int_0^1 (\rho_x)^2 dx \leq C \left(|u_x|_{L^\infty(0,1)} \int_0^1 \rho_x^2 dx + \int_0^1 G_x^2 dx \int_0^1 \rho_x^2 dx + \int_0^1 \rho_x^2 dx \right).$$

From the above lemmas and using Gronwall's inequality, we have

$$\sup_{0 \leq t \leq T} |\rho_x|_{L^2(0,1)} \leq C(T).$$

Lemma 2.6

$$\sup_{0 \leq t \leq T} (|\rho|_{H^1(0,1)} + |\rho_t|_{L^2(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|(\rho u)_t|_{L^2(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2) ds \leq C(T),$$

where C is dependent on $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$, $|f|_{L^2(0,T; L^{\frac{2\gamma}{\gamma-1}}(0,1))}$ and T , but is independent of the lower bound of ρ_0 .

Proof From equation (1.1), we have $\rho_t^2 = [-(\rho_x u + \rho u_x)]^2$. Integrating it over $(0, 1)$, we get

$$\begin{aligned} \int_0^1 \rho_t^2 dx & = \int_0^1 \rho_x^2 u^2 dx + 2 \int_0^1 \rho_x \rho u u_x dx + \int_0^1 \rho^2 u_x^2 dx \\ & \leq |u|_{L^\infty(0,1)}^2 \int_0^1 \rho_x^2 dx + 2|\rho|_{L^\infty(0,1)} |u|_{L^\infty(0,1)} |\rho_x|_{L^2(0,1)} |u_x|_{L^2(0,1)}^2 + |\rho|_{L^\infty(0,1)}^2 \int_0^1 u_x^2 dx. \end{aligned}$$

Thus

$$\begin{aligned} \sup_{0 \leq t \leq T} \int_0^1 \rho_t^2 dx &\leq \sup_{0 \leq t \leq T} |u|_{L^\infty(0,1)}^2 \sup_{0 \leq t \leq T} \int_0^1 \rho_x^2 dx \\ &+ 2 \sup_{0 \leq t \leq T} |\rho|_{L^\infty(0,1)} \sup_{0 \leq t \leq T} |u|_{L^\infty(0,1)} \sup_{0 \leq t \leq T} |\rho_x|_{L^2(0,1)} \sup_{0 \leq t \leq T} |u_x|_{L^2(0,1)} \\ &+ \sup_{0 \leq t \leq T} |\rho|_{L^\infty(0,1)}^2 \sup_{0 \leq t \leq T} |u_x|_{L^2(0,1)}^2 \leq C. \end{aligned} \quad (2.29)$$

Consequently

$$\begin{aligned} \int_0^T \int_0^1 (\rho u)_t^2 dx dt &\leq 2 \int_0^T \int_0^1 (\rho_t u)^2 dx dt + 2 \int_0^T \int_0^1 (\rho u_t)^2 dx dt \\ &\leq 2 \sup_{0 \leq t \leq T} |u|_{L^\infty(0,1)}^2 \int_0^T \int_0^1 (\rho_t)^2 dx dt \\ &+ 2 |\rho|^{\frac{1}{2}}|_{L^\infty(0,T) \times (0,1)} \int_0^T \int_0^1 (\sqrt{\rho} u_t)^2 dx dt \leq C. \end{aligned} \quad (2.30)$$

Combining the momentum equation and Poisson equation, we have

$$u_{xx} = (\rho u)_t + (\rho u^2)_x + p_x + \rho \Phi_x - \rho f. \quad (2.31)$$

Using (2.30) and $(\rho u_t) \in L^2((0, T) \times (0, 1))$, we can easily get $p_x \in L^2((0, T) \times (0, 1))$, and

$$[(\rho u^2)_x]^2 = (\rho_x u + 2\rho u u_x)^2 \leq 2(\rho_x^2 u^4 + 4\rho^2 u^2 u_x^2).$$

Combining $|u|_{L^\infty((0,T) \times (0,1))} \leq C$, $|\rho|_{L^\infty((0,T) \times (0,1))} \leq C$, $|\rho_x|_{L^\infty(0,T;L^2(0,1))} \leq C$ and $|u_x|_{L^\infty(0,T;L^2(0,1))} \leq C$, we see that $(\rho u^2)_x \in L^2(0, T) \times (0, 1)$. In fact, we have

$$\begin{aligned} &(\rho u^2)_x \in L^\infty(0, T; L^2(0, 1)), \\ &\int_0^T \int_0^1 |\rho \Phi_x|^2 dx dt \leq |\rho|_{L^\infty((0,T) \times (0,1))}^2 \int_0^T \int_0^1 |\Phi_x|^2 dx dt \leq C, \\ &\rho f \in L^2((0, T) \times (0, 1)). \end{aligned}$$

Combining the above estimates and (2.31), we have

$$\sup_{0 \leq t \leq T} (|\rho|_{H^1(0,1)} + |\rho_t|_{L^2(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|(\rho u)_t|_{L^2(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2) dt \leq C(T). \quad (2.32)$$

To prove Theorem 1.2, we must deal with the following estimate.

Lemma 2.7

$$\begin{aligned} &|\sqrt{\rho} u_t(t)|_{L^2(0,1)}^2 + \int_\tau^t |u_{tx}|_{L^2(0,1)}^2 ds \\ &\leq C + |\sqrt{\rho} u_t(\tau)|_{L^2(0,1)}^2 + C \int_0^t |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_t|_{L^2(0,1)}^2 ds, \end{aligned} \quad (2.33)$$

where C is dependent on $|\rho_0|_{H^1(0,1)}$, $|u_0|_{H_0^1(0,1)}$, $|f|_{L^2(0,T;L^{\frac{2\gamma}{\gamma-1}}(0,1))}$, $|f_x|_{L^2(0,T;L^2(0,1))}$ and $|f_t|_{L^2(0,T;L^2(0,1))}$, but is independent of the lower bound of ρ_0 .

Proof Combining the momentum equation (1.2) and the mass equation (1.1), we can easily get

$$\rho u_t + \rho uu_x + \rho\Phi_x + p_x - u_{xx} = \rho f.$$

Differentiating it with respect to time, we have

$$\rho u_{tt} + \rho_t u_t + (\rho u)u_{xt} + \rho_t uu_x + \rho u_t u_x + \rho_t\Phi_x + \rho\Phi_{xt} + p_{xt} - u_{xxt} = \rho_t f + \rho f_t,$$

that is,

$$\rho u_{tt} + \rho uu_{xt} + p_{xt} - u_{xxt} = -\rho_t(u_t + uu_x + \Phi_x - f) - \rho u_t u_x - \rho\Phi_{xt} + \rho f_t.$$

Multiplying it by u_t and integrating over $(0, 1)$, we have

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} \rho u_t^2 dx - \int_0^1 \frac{1}{2} \rho_t u_t^2 dx + \int_0^1 \frac{1}{2} (\rho uu_t^2)_x dx \\ & - \int_0^1 \frac{1}{2} (\rho u)_x u_t^2 dx - \int_0^1 p_t u_{xt} dx + \int_0^1 u_{xt}^2 dx \\ & = \int_0^1 (\rho u)_x (u_t^2 + uu_x u_t + \Phi_x u_t - fu_t) dx - \int_0^1 \rho u_t^2 u_x dx - \int_0^1 \rho\Phi_{tx} u_t dx + \int_0^1 \rho f_t u_t dx. \end{aligned}$$

From equation (1.1), we have

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} \rho u_t^2 dx + \int_0^1 u_{xt}^2 dx - \int_0^1 p_t u_{xt} dx \\ & = - \int_0^1 \rho u (u_t^2 + uu_x u_t + \Phi_x u_t - fu_t)_x dx \\ & - \int_0^1 \rho u_t^2 u_x dx - \int_0^1 \rho\Phi_{tx} u_t dx + \int_0^1 \rho f_t u_t dx. \end{aligned} \tag{2.34}$$

We deal with each term of (2.34) as follows:

$$\begin{aligned} - \int_0^1 p_t u_{xt} dx &= \int_0^1 a\gamma\rho^{\gamma-1}(\rho u)_x u_{xt} dx = \int_0^1 (p_x u + \gamma p u_x) u_{tx} dx \\ &= \frac{d}{dt} \int_0^1 \frac{\gamma}{2} p u_x^2 dx - \int_0^1 \frac{\gamma}{2} p_t u_x^2 dx + \int_0^1 p_x u u_{tx} dx \\ &= \frac{d}{dt} \int_0^1 \frac{\gamma}{2} p u_x^2 dx + \int_0^1 p_x u u_{tx} dx + \frac{\gamma}{2} \int_0^1 (p_x u + \gamma p u_x) u_x^2 dx \\ &= \frac{d}{dt} \int_0^1 \frac{\gamma}{2} p u_x^2 dx + \int_0^1 p_x u u_{tx} dx + \frac{\gamma}{2} \int_0^1 \gamma p u_x^3 dx \\ &\quad - \frac{\gamma}{2} \int_0^1 p u_x^3 dx - \frac{\gamma}{2} \int_0^1 p u (u_x^2)_x dx \\ &= \frac{d}{dt} \int_0^1 \frac{\gamma}{2} p u_x^2 dx + \int_0^1 p_x u u_{tx} dx + \frac{\gamma}{2} \int_0^1 (-p u (u_x^2)_x + (\gamma - 1) p u_x^3) dx. \end{aligned}$$

Substituting it into (2.34), we have

$$\begin{aligned}
& \frac{d}{dt} \int_0^1 \left(\frac{1}{2} \rho u_t^2 + \frac{\gamma}{2} p u_x^2 \right) dx + \int_0^1 u_{xt}^2 dx \\
& \leq \int_0^1 |p_x| |u| |u_{xt}| dx + \gamma \int_0^1 p |u_x| |u| |u_{xx}| dx + \frac{\gamma}{2} \int_0^1 (\gamma - 1) p |u_x|^3 dx \\
& \quad + 2 \int_0^1 \rho |u_t| |u| |u_{xt}| dx + \int_0^1 \rho |u_t| |u| |u_x|^2 dx + \int_0^1 \rho |u_t| |u|^2 |u_{xx}| dx + \int_0^1 \rho |u_t| |u|^2 |u_x| dx \\
& \quad - \int_0^1 \rho u (\Phi_x u_t)_x dx + \int_0^1 \rho |f_x| |u| |u_t| dx + \int_0^1 \rho |f| |u| |u_{xt}| dx \\
& \quad + \int_0^1 \rho |u_t|^2 |u_x| dx - \int_0^1 \rho \Phi_{xt} u_t dx + \int_0^1 \rho |f_t| |u_t| dx \equiv \sum_{j=1}^{13} I_j. \tag{2.35}
\end{aligned}$$

Next, we deal with I_1 – I_{13} :

$$\begin{aligned}
I_1 &= \int_0^1 |p_x| |u| |u_{xt}| dx \leq \varepsilon |u_{xt}|_{L^2(0,1)}^2 + C(\varepsilon) |p_x|_{L^2(0,1)}^2; \\
I_2 &= \gamma \int_0^1 p |u_x| |u| |u_{xx}| dx \leq C \int_0^1 |u_x| |u_{xx}| dx \leq C(|u_x|_{L^\infty(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2); \\
I_3 &= \frac{\gamma}{2} \int_0^1 (\gamma - 1) p |u_x|^3 dx \leq C |u_x|_{L^2(0,1)} |u_x|_{L^\infty(0,1)}^2 \leq C |u_x|_{L^\infty(0,1)}^2; \\
I_4 &= 2 \int_0^1 \rho |u_t| |u| |u_{xt}| dx \leq C |\sqrt{\rho} u_t|_{L^2(0,1)} |u_{xt}|_{L^2(0,1)} \leq \varepsilon |u_{xt}|_{L^2(0,1)}^2 + C(\varepsilon) |\sqrt{\rho} u_t|_{L^2(0,1)}^2; \\
I_5 &= \int_0^1 \rho |u_t| |u| |u_x|^2 dx \leq C |\sqrt{\rho} u_t|_{L^2(0,1)} |u_x|_{L^\infty(0,1)} |u_x|_{L^2(0,1)} \\
&\leq C |\sqrt{\rho} u_t|_{L^2(0,1)}^2 + C |u_x|_{L^\infty(0,1)}^2; \\
I_6 &= \int_0^1 \rho |u_t| |u|^2 |u_{xx}| dx \leq C |\sqrt{\rho} u_t|_{L^2(0,1)} |u_{xx}|_{L^2(0,1)} \leq C (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2); \\
I_7 &= \int_0^1 \rho |u_t| |u|^2 |u_x| dx \leq C (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_x|_{L^2(0,1)}^2); \\
I_8 &= \left| \int_0^1 \rho u (\Phi_x u_t)_x dx \right| \leq \int_0^1 \rho |u| |\Phi_x| |u_{xt}| dx + \int_0^1 \rho |u| |\Phi_{xx}| |u_t| dx \\
&\leq C |\Phi_x|_{L^2(0,1)}^2 + \varepsilon |u_{xt}|_{L^2(0,1)}^2 + C |\sqrt{\rho} u_t|_{L^2(0,1)}^2; \\
I_9 &= \int_0^1 \rho |f_x| |u| |u_t| dx \leq C |\sqrt{\rho} u_t|_{L^2(0,1)}^2 + C |f_x|_{L^2(0,1)}^2; \\
I_{10} &= \int_0^1 \rho |f| |u| |u_{xt}| dx \leq C |f|_{L^2(0,1)}^2 + \varepsilon |u_{xt}|_{L^2(0,1)}^2; \\
I_{11} &= \int_0^1 \rho |u_t|^2 |u_x| dx \leq C |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_t|_{L^2(0,1)}^2; \\
I_{12} &= \left| \int_0^1 \rho \Phi_{xt} u_t dx \right| \leq C |\sqrt{\rho} u_t|_{L^2(0,1)}^2 + C \left(\int_0^1 |\Phi_{xt}|^2 dx \right).
\end{aligned}$$

We deal with the estimate of Φ_{xt} .

Differentiating (1.3) with respect to time, multiplying it by Φ_t and integrating over $(0, 1)$,

we get

$$\int_0^1 \Phi_{xxt} \Phi_t dx = 4\pi g \int_0^1 \rho_t \Phi_t dx.$$

Then

$$\int_0^1 |\Phi_{xt}|^2 dx \leq C |\rho_t|_{L^2(0,1)} |\Phi_t|_{L^2(0,1)} \leq C |\rho_t|_{L^2(0,1)}^2 + \varepsilon |\Phi_t|_{L^2(0,1)}^2.$$

Thus

$$\int_0^1 |\Phi_{xt}|^2 dx \leq C |\rho_t|_{L^2(0,1)}^2.$$

Consequently, I_{12} becomes

$$\begin{aligned} I_{12} &\leq C(|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |\rho_t|_{L^2(0,1)}^2), \\ I_{13} &= \int_0^1 \rho |f_t| |u_t| dx \leq C |f_t|_{L^2(0,1)}^2 + C |\sqrt{\rho} u_t|_{L^2(0,1)}^2. \end{aligned}$$

From the estimates of I_1 – I_{13} , and (2.35), we obtain

$$\begin{aligned} \frac{d}{dt} \int_0^1 \left(\frac{1}{2} \rho u_t^2 + \frac{\gamma}{2} p u_x^2 \right) dx + \int_0^1 u_{xt}^2 dx &\leq C(1 + |\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |\rho_t|_{L^2(0,1)}^2 + |u_x|_{L^2(0,1)}^2 \\ &\quad + |u_x|_{L^\infty(0,1)}^2 + |f_t|_{L^2(0,1)}^2 + |f_x|_{L^2(0,1)}^2 + |p_x|_{L^2(0,1)}^2) \\ &\quad + C |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_t|_{L^2(0,1)}^2. \end{aligned} \quad (2.36)$$

Integrating it over $(\tau, t) \subset (0, T)$, we conclude that

$$\begin{aligned} &\int_0^1 \rho u_t^2(t) dx + \int_\tau^t \int_0^1 u_{xt}^2 dx ds \\ &\leq C + C \left(\int_0^1 \rho u_t^2(\tau) dx + \int_0^1 p u_x^2(\tau) dx \right) + C \int_\tau^t |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_s|_{L^2(0,1)}^2 ds. \end{aligned}$$

Using Lemma 2.3, we get

$$\begin{aligned} &\int_0^1 \rho u_t^2(t) dx + \int_\tau^t \int_0^1 u_{xt}^2 dx ds \\ &\leq C + C |\sqrt{\rho} u_t(\tau)|_{L^2(0,1)}^2 + C \int_\tau^t |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_s|_{L^2(0,1)}^2 ds. \end{aligned} \quad (2.37)$$

Combining the momentum equation (1.2) and compatibility condition, we have

$$\rho u_t + \rho u u_x + \rho \Phi_x + p_x - u_{xx} = \rho f.$$

Thus

$$\rho^{\frac{1}{2}} u_t = -(\rho^{\frac{1}{2}} u u_x + \rho^{\frac{1}{2}} \Phi_x + \rho^{-\frac{1}{2}} p_x - \rho^{-\frac{1}{2}} u_{xx}) + \rho^{\frac{1}{2}} f.$$

Consequently

$$|\sqrt{\rho} u_t(\tau)|_{L^2(0,1)}^2 \leq C(|\sqrt{\rho} f|_{L^2(0,1)}^2 + |\sqrt{\rho} \Phi_x|_{L^2(0,1)}^2 + |\sqrt{\rho} u u_x|_{L^2(0,1)}^2 + |\rho^{-\frac{1}{2}} p_x - \rho^{-\frac{1}{2}} u_{xx}|_{L^2(0,1)}^2).$$

Combining the above estimates $f_t \in L^2(0, 1; L^2(0, 1))$, $\sup_{0 \leq t \leq T} |f|_{L^2(0,1)}^2 \leq C$, $|\Phi_x|_{L^2(0,1)}^2 \leq C |\rho|_{L^\gamma(0,1)}^\gamma + C$, $|\rho^{\frac{1}{2}} u u_x|_{L^2(0,1)}^2 \leq C |u_x|_{L^2(0,1)}^2$ and $\sup_{0 \leq t \leq T} |u_x|_{L^2(0,1)}^2 \leq C$, we obtain

$$|\sqrt{\rho} u_t(\tau)|_{L^2(0,1)}^2 \leq C + C |\rho^{-\frac{1}{2}} p_x(\tau) - \rho^{-\frac{1}{2}} u_{xx}(\tau)|_{L^2(0,1)}^2.$$

Substituting it into (2.37) and letting $\tau \rightarrow 0$, we get

$$\int_0^1 \frac{1}{2} \rho u_t^2(t) dx + \int_0^t \int_0^1 u_{xt}^2 dx ds \leq C + C \int_0^t |u_x|_{L^\infty(0,1)} |\sqrt{\rho} u_s|_{L^2(0,1)}^2 ds.$$

Using Gronwall's inequality and $\int_0^T |u_x|_{L^\infty(0,1)} dt \leq C$, we get

$$\int_0^1 \rho u_t^2(t) dx + \int_0^t \int_0^1 u_{xt}^2 dx ds \leq C. \quad (2.38)$$

Next, we construct the approximate systems to deal with the existence.

3 Proof of the Existence

Our method that constructed approximate systems is similar to that in [9]. We take a semi-discrete Galerkin scheme. We take our basic function space as $X = H_0^1(0, 1) \cap H^2(0, 1)$ and the finite-dimensional subspaces as $X^m = \text{span}\{\varphi^1, \varphi^2, \dots, \varphi^m\} \subset X \cap C^2([0, 1])$. Here φ^m is the m th eigenfunction of the strongly elliptic operator $A = -\frac{\partial^2}{\partial x^2}$ defined on X .

Let ρ_0 , u_0 and f satisfy the hypotheses of Theorem 1.1 or Theorem 1.2. Assume for the moment that $\rho_0^\delta \in C^1([0, 1])$ and $\rho_0^\delta \geq \delta$ in $(0, 1)$ (for some constant $\delta > 0$). We may construct an approximate solution for any $v \in X^m$, $\varphi \in C^2([0, 1])$

$$\begin{cases} \int_\Omega (\rho^m u_t^m + \rho^m u^m \cdot u_x^m + A u^m + p_x^m + \rho^m \Phi_x^m) \cdot v dx = \int_\Omega \rho^m f^\delta \cdot v dx, \\ \int_\Omega \rho_t^m \varphi dx + \int_\Omega (\rho^m u^m)_x \varphi dx = 0, \\ \int_\Omega \Phi_{xx}^m \varphi dx = 4\pi g \int_\Omega \left(\rho^m - \frac{m_0}{|\Omega|}\right) \varphi dx, \end{cases}$$

where $f^\delta \in C^1((0, T) \times (0, 1))$ and $f^\delta \rightarrow f$ in $L^2(0, T; L^{\frac{2\gamma}{\gamma-1}}(0, 1))$. The initial and boundary conditions are

$$\begin{aligned} u_0^m &\equiv \sum_{k=1}^m (u_0, \varphi^k)_{L^2(\Omega)} \varphi^k \quad \text{and} \quad \rho^m(0) = \rho_0^\delta > \delta, \quad \rho^\delta(0) < |\rho_0|_{L^\infty} + 1, \\ |\rho_0^\delta - \rho_0|_{H^1(0,1)} &\rightarrow 0, \quad u^m(0, x) = u^m(1, x) = 0, \quad \Phi^m(0, x) = \Phi^m(1, x) = 0. \end{aligned}$$

Under the hypotheses of Theorem 1.1, similarly, for any fixed $\delta > 0$, we may get the similar estimate of Lemmas 2.1–2.6.

$$\begin{aligned} \sup_{0 \leq t \leq T} (|\rho_\delta^m|_{L^\infty(0,1)} + |u_{x\delta}^m|_{L^2(0,1)}) + \int_0^T (|\sqrt{\rho_\delta^m} u_{\delta t}^m|_{L^2(0,1)}^2 + |u_{x\delta}^m|_{L^2(0,1)}^2 + |G_{x\delta}^m|_{L^2(0,1)}^2) dt &\leq C(T), \\ \sup_{0 \leq t \leq T} (|\rho_\delta^m|_{H^1(0,1)} + |\rho_{\delta t}^m|_{L^2(0,1)} + |u_{\delta x}^m|_{L^2(0,1)}) + \int_0^T (|(\rho_\delta^m u_\delta^m)_t|_{L^2(0,1)}^2 + |u_{x\delta}^m|_{L^2(0,1)}^2) dt &\leq C(T). \end{aligned}$$

Combining the course of estimates and the initial condition of approximate system, we can easily deduce that C is dependent on T , ρ_0 , u_0 and f . Moreover, because the constants C of Lemmas 2.1–2.6 are independent of the lower bound of ρ_0 . Here, $C(T)$ does not depend on δ and m (for any $m \geq M$, M is dependent on the approximate velocity of initial condition). Thus,

we can deduce from the two above estimates that (ρ^m, u^m, Φ^m) converges, up to an extraction of subsequences, to some limit $(\rho_\delta, u_\delta, \Phi_\delta)$ in the obvious weak sense, and there are estimates:

$$\begin{aligned} \sup_{0 \leq t \leq T} (|\rho_\delta|_{L^\infty(0,1)} + |u_{x\delta}|_{L^2(0,1)}) + \int_0^T (|\sqrt{\rho_\delta} u_{\delta t}|_{L^2(0,1)}^2 + |u_{x\delta}|_{L^2(0,1)}^2 + |G_{x\delta}|_{L^2(0,1)}^2) dt &\leq C(T), \\ \sup_{0 \leq t \leq T} (|\rho_\delta|_{H^1(0,1)} + |\rho_{\delta t}|_{L^2(0,1)} + |u_{\delta x}|_{L^2(0,1)}) + \int_0^T (|(\rho_\delta u_\delta)_t|_{L^2(0,1)}^2 + |u_{xx\delta}|_{L^2(0,1)}^2) dt &\leq C(T). \end{aligned}$$

Because $C(T)$ is independent of δ , when $\delta \rightarrow 0$, we can deduce that $(\rho_\delta, u_\delta, \Phi_\delta)$ converges, up to an extraction of subsequences, to some limit (ρ, u, Φ) in weak sense, and

$$\begin{aligned} \sup_{0 \leq t \leq T} (|\rho|_{L^\infty(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|\sqrt{\rho} u_t|_{L^2(0,1)}^2 + |u_x|_{L^2(0,1)}^2 + |G_x|_{L^2(0,1)}^2) dt &\leq C(T), \\ \sup_{0 \leq t \leq T} (|\rho|_{H^1(0,1)} + |\rho_t|_{L^2(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|(\rho u)_t|_{L^2(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2) dt &\leq C(T). \end{aligned}$$

From the L^p -strong estimates of Poisson equation, we can easily get the regularity in Theorem 1.1.

As for Theorem 1.2, we can deal with it similarly, but the initial and outer power conditions are

$$\begin{aligned} \delta \leq \rho^\delta(0) \leq |\rho_0|_{L^\infty} + 1, \quad \rho_0^\delta \in C^2([0, 1]), \quad g^\delta \in C_c^2(0, 1), \quad |g_0^\delta - g|_{L^2(0,1)} \rightarrow 0, \\ f^\delta \in C_c^2((0, T) \times (0, 1)), \quad |\rho_0^\delta - \rho_0|_{H^1(0,1)} \rightarrow 0, \quad |(f^\delta, f_x^\delta, f_t^\delta) - (f, f_x, f_t)|_{L_{loc}^2(0, T; L^2(0,1))} \rightarrow 0. \end{aligned}$$

Because we have compatibility condition, we let $u_0^\delta \in C^3[0, 1]$ be the solution of the following elliptic equation

$$u_{0xx}^\delta = (p^\delta)_x + \rho_0^{\frac{1}{2}} g^\delta, \quad 0 < x < 1, \quad u_0^\delta(0) = u_0^\delta(1) = 0.$$

Combining the classical stableness results of the elliptic equation and the compatibility condition of Theorem 1.2, we deduce that $u_0^\delta \rightarrow u_0$ in $H^2(0, 1)$, and u_0 satisfies the compatibility of Theorem 1.2.

For any fixed $\delta > 0$, similarly, we may get the similar results with Lemmas 2.1–2.7, and we have estimates

$$\begin{aligned} \sup_{0 \leq t \leq T} (|\rho^\delta|_{H^1(0,1)} + |\rho_t^\delta|_{L^2(0,1)} + |u_x^\delta|_{L^2(0,1)}) + \int_0^T (|u_x^\delta|_{L^\infty(0,1)}^2 + |u_{xx}^\delta|_{L^2(0,1)}^2) dt &\leq C(T), \\ |\sqrt{\rho^\delta} u_t^\delta|_{L^2(0,1)}^2 + \int_0^T |u_{xt}^\delta|_{L^2(0,1)}^2 dt &\leq C(T), \end{aligned}$$

where C is independent of δ . Thus when $\delta \rightarrow 0$, we can deduce from the above estimates that $(\rho_\delta, u_\delta, \Phi_\delta)$ converges to some limit (ρ, u, Φ) , and we have estimates

$$\begin{aligned} \sup_{0 \leq t \leq T} (|\rho|_{H^1(0,1)} + |\rho_t|_{L^2(0,1)} + |u_x|_{L^2(0,1)}) + \int_0^T (|u_x|_{L^\infty(0,1)}^2 + |u_{xx}|_{L^2(0,1)}^2) dt &\leq C(T), \\ |\sqrt{\rho} u_t(t)|_{L^2(0,1)}^2 + \int_0^T |u_{xt}|_{L^2(0,1)}^2 dt &\leq C(T). \end{aligned}$$

From L^p -strong estimates of Poisson equation, we can easily get the regularity in Theorem 1.2.

4 Proof of the Uniqueness

Let (ρ, u, Φ) and $(\bar{\rho}, \bar{u}, \bar{\Phi})$ be two solutions that satisfy the same initial condition. Then combining (1.1) and (1.2), we have

$$\rho u_t + \rho uu_x + \rho \Phi_x + p_x - u_{xx} = \rho f, \quad \bar{\rho} \bar{u}_t + \bar{\rho} \bar{u} \bar{u}_x + \bar{\rho} \bar{\Phi}_x + \bar{p}_x - \bar{u}_{xx} = \bar{\rho} f.$$

Thus

$$\begin{aligned} & \rho(u - \bar{u})_t + \rho u(u - \bar{u})_x - (u - \bar{u})_{xx} \\ &= (\rho - \bar{\rho})(f - \bar{u}_t - \bar{u} \bar{u}_x - \bar{\Phi}_x) - (p - \bar{p})_x - \rho(\Phi - \bar{\Phi})_x - \rho(u - \bar{u})\bar{u}_x. \end{aligned} \quad (4.1)$$

Multiplying it by $(u - \bar{u})$ and integrating over $(0, 1)$, we have

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} \rho(u - \bar{u})^2 dx + \frac{1}{2} \int_0^1 (\rho u)_x(u - \bar{u})^2 dx + \int_0^1 \rho u(u - \bar{u})_x(u - \bar{u}) dx + \int_0^1 (u - \bar{u})_x^2 dx \\ & \leq \int_0^1 |\rho - \bar{\rho}| |f - \bar{u}_t - \bar{u} \bar{u}_x - \bar{\Phi}_x| |u - \bar{u}| dx + \int_0^1 |p - \bar{p}| |(u - \bar{u})_x| dx \\ & \quad + \int_0^1 \rho |(\Phi - \bar{\Phi})_x| |u - \bar{u}| dx + \int_0^1 \rho |u - \bar{u}|^2 |\bar{u}_x| dx, \end{aligned}$$

that is,

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} \rho(u - \bar{u})^2 dx + \int_0^1 (u - \bar{u})_x^2 dx \\ & \leq |\rho - \bar{\rho}|_{L^2(0,1)} |f - \bar{u}_t - \bar{u} \bar{u}_x - \bar{\Phi}_x|_{L^2(0,1)} |u - \bar{u}|_{L^\infty(0,1)} + |p - \bar{p}|_{L^2(0,1)} |(u - \bar{u})_x|_{L^2(0,1)} \\ & \quad + C |(\Phi - \bar{\Phi})_x|_{L^2(0,1)} |u - \bar{u}|_{L^2(0,1)} + |\sqrt{\rho}| |u - \bar{u}|_{L^2(0,1)}^2 |\bar{u}_x|_{L^\infty(0,1)} \\ & \leq \varepsilon |(u - \bar{u})_x|_{L^2(0,1)}^2 + |\rho - \bar{\rho}|_{L^2(0,1)}^2 (C + C |\bar{u}_t|_{L^2(0,1)}^2) \\ & \quad + |p - \bar{p}|_{L^2(0,1)}^2 + |\sqrt{\rho}| |u - \bar{u}|_{L^2(0,1)}^2 |\bar{u}_x|_{L^\infty(0,1)}. \end{aligned}$$

Consequently,

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} \rho(u - \bar{u})^2 dx + \int_0^1 (u - \bar{u})_x^2 dx \\ & \leq |\rho - \bar{\rho}|_{L^2(0,1)} (C + C |\bar{u}_t|_{L^2(0,1)}^2) + |p - \bar{p}|_{L^2(0,1)}^2 + |\sqrt{\rho}| |u - \bar{u}|_{L^2(0,1)}^2 |\bar{u}_x|_{L^\infty(0,1)}. \end{aligned} \quad (4.2)$$

Moreover, from the conservative mass equation, we have

$$\rho_t + \rho_x u + \rho u_x = 0, \quad \bar{\rho}_t + \bar{\rho}_x \bar{u} + \bar{\rho} \bar{u}_x = 0.$$

Then

$$(\rho - \bar{\rho})_t + (\rho - \bar{\rho})_x u + \bar{\rho}_x u - \bar{\rho}_x \bar{u} + (\rho - \bar{\rho}) u_x + \bar{\rho} u_x - \bar{\rho} \bar{u}_x = 0,$$

that is,

$$(\rho - \bar{\rho})_t + (\rho - \bar{\rho})_x u + \bar{\rho}_x (u - \bar{u}) + (\rho - \bar{\rho}) u_x + \bar{\rho} (u_x - \bar{u}_x) = 0.$$

Multiplying it by $(\rho - \bar{\rho})$ and integrating over $(0, 1)$, we get

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2} (\rho - \bar{\rho})^2 dx - \int_0^1 \frac{1}{2} (\rho - \bar{\rho})^2 u_x dx + \int_0^1 \bar{\rho}_x (u - \bar{u}) (\rho - \bar{\rho}) dx \\ & \quad + \int_0^1 (\rho - \bar{\rho})^2 u_x dx + \int_0^1 \bar{\rho} (u - \bar{u})_x (\rho - \bar{\rho}) dx = 0. \end{aligned}$$

Thus

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2}(\rho - \bar{\rho})^2 dx \\ & \leq C \left(\int_0^1 (\rho - \bar{\rho})^2 |u_x| dx + \int_0^1 |\bar{\rho}_x| |(u - \bar{u})| |(\rho - \bar{\rho})| dx + \int_0^1 \bar{\rho} |(u - \bar{u})_x| |(\rho - \bar{\rho})| dx \right) \end{aligned}$$

that is,

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2}(\rho - \bar{\rho})^2 dx \\ & \leq C \left(|u_x|_{L^\infty(0,1)} \int_0^1 (\rho - \bar{\rho})^2 dx + \left(\int_0^1 \rho_x^2 dx \right)^{\frac{1}{2}} |u - \bar{u}|_{L^\infty(0,1)} \right) \left(\int_0^1 (\rho - \bar{\rho})^2 dx \right)^{\frac{1}{2}} \\ & \quad + \left(\int_0^1 |(u - \bar{u})_x|^2 dx \right)^{\frac{1}{2}} \left(\int_0^1 (\rho - \bar{\rho})^2 dx \right)^{\frac{1}{2}} \\ & \leq C \left(|u_x|_{L^\infty(0,1)} + C(\varepsilon) + C(\varepsilon) \int_0^1 \rho_x^2 dx \right) \int_0^1 (\rho - \bar{\rho})^2 dx + \varepsilon |(u - \bar{u})_x|_{L^2(0,1)}^2. \end{aligned} \quad (4.3)$$

Moreover, multiplying (1.1) by $a\gamma\rho^{\gamma-1}$, we get

$$p_t + p_x u + \gamma p u_x = 0, \quad \bar{p}_t + \bar{p}_x \bar{u} + \gamma \bar{p} \bar{u}_x = 0.$$

Similarly, we get

$$(p - \bar{p})_t + (p - \bar{p})_x u + \bar{p}_x u - \bar{p}_x \bar{u} + \gamma(p - \bar{p})u_x + \gamma \bar{p} u_x - \gamma \bar{p} \bar{u}_x = 0,$$

that is,

$$(p - \bar{p})_t + (p - \bar{p})_x u + \bar{p}_x(u - \bar{u}) + \gamma(p - \bar{p})u_x + \gamma \bar{p}(u - \bar{u})_x = 0.$$

Multiplying it by $(p - \bar{p})$ and integrating over $(0, 1)$, we get

$$\begin{aligned} & \frac{d}{dt} \int_0^1 \frac{1}{2}(p - \bar{p})^2 dx - \int_0^1 \frac{1}{2}(p - \bar{p})^2 dx + \int_0^1 \bar{p}(u - \bar{u})(p - \bar{p}) dx \\ & + \gamma \int_0^1 (p - \bar{p})^2 u_x dx + \gamma \int_0^1 \bar{p}(u - \bar{u})_x(p - \bar{p}) dx = 0. \end{aligned}$$

Then

$$\begin{aligned} & \frac{d}{dt} \int_0^1 (p - \bar{p})^2 dx \leq C |u_x|_{L^\infty(0,1)} |p - \bar{p}|_{L^2(0,1)}^2 + |\bar{p}_x|_{L^2(0,1)} |u - \bar{u}|_{L^\infty(0,1)} |p - \bar{p}|_{L^2(0,1)} \\ & \quad + \varepsilon |(u - \bar{u})_x|_{L^2(0,1)}^2 + C(\varepsilon) |p - \bar{p}|_{L^2(0,1)}^2 \\ & \leq C(|u_x|_{L^\infty(0,1)} + |\bar{p}_x|_{L^2(0,1)} + 1) |p - \bar{p}|_{L^2(0,1)}^2 + 2\varepsilon |(u - \bar{u})_x|_{L^2(0,1)}^2. \end{aligned} \quad (4.4)$$

From (4.2)–(4.4), we obtain

$$\begin{aligned} & \frac{d}{dt} \int_0^1 (\rho(u - \bar{u})^2 + (\rho - \bar{\rho})^2 + (p - \bar{p})^2) dx + \int_0^1 (u - \bar{u})_x^2 dx \\ & \leq C(1 + |\bar{u}_t|_{L^2(0,1)}^2 + |\bar{u}_x|_{L^\infty(0,1)} + |\rho_x|_{L^2(0,1)}^2 + |u_x|_{L^\infty(0,1)} + |\bar{p}_x|_{L^2(0,1)}^2) \\ & \quad \cdot (|\sqrt{\rho}(u - \bar{u})|_{L^2(0,1)}^2 + |\rho - \bar{\rho}|_{L^2(0,1)}^2 + |p - \bar{p}|_{L^2(0,1)}^2). \end{aligned}$$

Using Gronwall's inequality and the above regularity of strong solution, we have

$$|\sqrt{\rho}(u - \bar{u})|_{L^2(0,1)}^2 + |\rho - \bar{\rho}|_{L^2(0,1)}^2 + |p - \bar{p}|_{L^2(0,1)}^2 = 0,$$

that is, $u = \bar{u}$, $\rho = \bar{\rho}$ in $L^2(0, 1)$. From the classical theorems of Poisson equation, we get $|\Phi - \bar{\Phi}|_{W^{2,2}(0,1)} = 0$. Finally, we get the uniqueness.

Acknowledgement The authors thank the referee for his (her) useful comments.

References

- [1] Kobayashi, T. and Suzuki, T., Weak solutions to the Navier-Stokes-Poisson equation, preprint.
- [2] Feireisl, E., Novotný, A. and Petzeltová, H., On the existence of globally defined weak solution to the Navier-Stokes equations, *J. Math. Fluid Mech.*, **3**, 2001, 358–392.
- [3] Beirão da Veiga, H., Long time behavior for one-dimensional motion of a general barotropic viscous fluid, *Arch. Rat. Mech. Anal.*, **108**, 1989, 141–160.
- [4] Kazhikov, A. V., Stabilization of solutions of an initial-boundary-value problem for the equations of motion of a barotropic viscous fluid, *Diff. Eqs.*, **15**, 1979, 463–467.
- [5] Solonnikov, V. A. and Kazhikov, A. V., Existence theorem for the equations of motion of a compressible viscous fluid, *Ann. Rev. Fluid Mech.*, **13**, 1981, 79–95.
- [6] Straškraka, I. and Valli, A., Asymptotic behavior of the density for one-dimensional Navier-Stokes equations, *Manuscripta Math.*, **62**, 1988, 401–416.
- [7] Matsumura, A. and Yanagi, S., Uniform boundedness of the solutions for a one dimensional isentropic model system of compressible viscous gas, *Comm. Math. Phys.*, **175**, 1996, 259–274.
- [8] Jiang, S. and Zhang, P., On spherically symmetric solutions of the compressible isentropic Navier-Stokes equations, *Comm. Math. Phys.*, **215**, 2001, 559–581.
- [9] Choe, H. J. and Kim, H., Strong solutions of the Navier-Stokes equations for isentropic compressible fluids, *J. Diff. Eqs.*, **190**, 2003, 504–523.
- [10] Choe, H. J. and Kim, H., Global existence of the radially symmetric solutions of the radially symmetric solutions of the Navier-Stokes equation for the isentropic compressible fluids, *Math. Meth. Appl. Sci.*, **28**, 2005, 1–28.
- [11] Lions, P. L., Mathematical topics in fluids mechanics, Vol. 2, Oxford Lecture Series in Mathematics and Its Applications, Vol. 10, Clarendon Press, Oxford, 1998.
- [12] Xin, Z., Blow up of smooth solutions to the compressible Navier-Stokes equation with compact density, *Comm. Pure Appl. Math.*, **51**, 1998, 229–240.
- [13] Hoff, D., Global solutions of the Navier-Stokes equations for multidimensional compressible flow with discontinuous initial data, *J. Diff. Eqs.*, **120**, 1995, 215–254.
- [14] Hoff, D. and Smoller, J., Non-formation of vacuum states for compressible Navier-Stokes equations, *Comm. Math. Phys.*, **216**, 2001, 255–276.
- [15] Liu, T. P., Xin, Z. and Yang, T., Vacuum states of compressible flow, *Discrete and Continuous Dyn. Systems*, **4**, 1998, 1–32.
- [16] Luo, T., Xin, Z. and Yang, T., Interface behavior of compressible Navier-Stokes equations with vacuum, *SIAM J. Math. Anal.*, **31**(6), 2000, 1175–1191.
- [17] Yang, T., Yao, Z. and Zhu, C., Compressible Navier-Stokes equations with density-dependent viscosity and vacuum, *Comm. PDE*, **26**(5-6), 2001, 965–981.
- [18] Okada, M. and Makino, T., Free boundary problem for the equation of spherically symmetric motion of viscous gas, *Japan J. Indust. Appl. Math.*, **10**, 1993, 219–235.
- [19] Matsusue-Necasova, S., Okada, M. and Makino, T., Free boundary problem for the equation of spherically symmetric motion of viscous gas, (*The Third*) *Japan J. Indust. Appl. Math.*, **14**, 1997, 199–213.