

# Gradient Estimates for the Heat Kernels in Higher Dimensional Heisenberg Groups\*\*

Bin QIAN\*

**Abstract** The author obtains sharp gradient estimates for the heat kernels in two kinds of higher dimensional Heisenberg groups — the non-isotropic Heisenberg group and the Heisenberg type group  $\mathbb{H}_{n,m}$ . The method used here relies on the positive property of the Bakry-Émery curvature  $\Gamma_2$  on the radial functions and some associated semigroup technics.

**Keywords** Gradient estimates,  $\Gamma_2$  curvature, Heat kernels, Sublaplace, Heisenberg group

**2000 MR Subject Classification** 58J35, 43A80

## 1 Introduction

The study of properties of second-order self-adjoint differential operators often depends on Gaussian bounds for the corresponding heat kernels (gradient estimates or not), that is, the kernel of the semigroup generated by these operators. In the past few years, kinds of estimates for the heat kernels were obtained, see [3, 5, 7–9, 13–16, 18].

In the study of the long (or small) time behavior (e.g., gradient estimates, ergodicity, etc.) of simple linear parabolic evolution equations, one often uses lower bounds on the Ricci curvature associated to the generator of the heat kernel (see, for example, [1, 12, 18] and the references therein). But this method fails in general hypoelliptic evolution equations, since the Ricci ( $\Gamma_2$ -) curvature in even the simplest example of the Heisenberg group can not be bounded below as explained, e.g., in [2, 11]. Nevertheless, in the Heisenberg group (type, or nonisotropic) case, many properties of the elliptic case remain true. In this paper, we focus on the gradient estimates for the heat kernels in higher dimensional Heisenberg groups.

Let us recall first some basic facts.

**The elliptic case** Let  $M$  be a complete Riemannian manifold of dimension  $n$  and let  $\mathcal{L} := \Delta + \nabla h$ , where  $\Delta$  is the Laplace-Beltrami operator. For  $t \geq 0$ , denote by  $P_t$  the heat semigroup generated by  $\mathcal{L}$  (that is formally  $P_t = \exp(t\mathcal{L})$ ), and let  $H(t, x, y)$  be the heat kernel.

In [8], Engoulatov obtained the following gradient estimates for the heat kernels on Riemannian manifolds.

**Theorem 1.1** *Let  $M$  be a complete Riemannian manifold of dimension  $n$  with Ricci curvature bounded from below, and  $\text{Ric}(M) \geq -\rho$ ,  $\rho \geq 0$ .*

---

Manuscript received June 3, 2009. Revised September 22, 2009. Published online April 20, 2010.

\*Department of Mathematics and Statistics, Changshu Institute of Technology, Changshu 215500, Jiangsu, China; Institut de Mathématiques de Toulouse, Université de Toulouse, CNRS 5219, France.

E-mail: binqiancn@yahoo.com.cn

\*\*Project supported by China Scholarship Council (No. 2007U13020).

(i) Suppose a non-collapsing condition is satisfied on  $M$ , namely, there exist  $t_0 > 0$  and  $\nu_0 > 0$ , such that for any  $x \in M$ , the volume of the geodesic ball of radius  $t_0$  centered at  $x$  is not too small, i.e.  $\text{Vol}(B_x(t_0)) \geq \nu_0$ . Then there exist two constants  $C(\rho, n, \nu_0, t_0)$  and  $\bar{C}(t_0) > 0$ , such that

$$|\nabla \log H(t, x, y)| \leq C(\rho, n, \nu_0, t_0) \left( \frac{d(x, y)}{t} + \frac{1}{\sqrt{t}} \right)$$

uniformly on  $(0, \bar{C}(t_0)] \times M \times M$ , where  $d(x, y)$  is the Riemannian distance between  $x$  and  $y$ .

(ii) Suppose that  $M$  has a diameter bounded by  $D$ . Then there exists a constant  $C(\rho, n)$ , such that

$$|\nabla \log H(t, x, y)| \leq C(\rho, n) \left( \frac{D}{t} + \frac{1}{\sqrt{t}} + \rho\sqrt{t} \right)$$

uniformly on  $(0, \infty) \times M \times M$ .

**Heisenberg group case** The Heisenberg group can be seen as the Euclidean space  $\mathbb{R}^3$  with a group structure  $\circ$ , which is defined, for  $\vec{x} = (x, y, z)$  and  $\vec{y} = (x', y', z') \in \mathbb{R}^3$ , by

$$\vec{x} \circ \vec{y} = \left( x + x', y + y', z + z' + \frac{1}{2}(xy' - x'y) \right).$$

The left invariant vector fields which are given by

$$\begin{aligned} X(f) &= \lim_{\varepsilon \rightarrow 0} \frac{f(\vec{x} \circ (\varepsilon, 0, 0)) - f(\vec{x})}{\varepsilon} = \left( \partial_x - \frac{y}{2} \partial_z \right) f, \\ Y(f) &= \lim_{\varepsilon \rightarrow 0} \frac{f(\vec{x} \circ (0, \varepsilon, 0)) - f(\vec{x})}{\varepsilon} = \left( \partial_y + \frac{x}{2} \partial_z \right) f, \\ Z(f) &= \lim_{\varepsilon \rightarrow 0} \frac{f(\vec{x} \circ (0, 0, \varepsilon)) - f(\vec{x})}{\varepsilon} = \partial_z f. \end{aligned}$$

The sublaplace is defined by

$$\Delta_{\mathbb{H}} = X^2 + Y^2.$$

Let  $P_t$  be the associated semigroup generated by  $\Delta_{\mathbb{H}}$ , and  $p_t(\vec{x}, \vec{y})$  be the heat kernel of  $P_t$ . Though the  $\Gamma_2$  curvature can not be bounded below, explained in [2, 11]. Recently, Li has shown the following sharp gradient estimates for the heat kernel in the Heisenberg group (see [13]).

**Theorem 1.2** *There exists a constant  $C > 0$  such that for  $t > 0$  and  $g = (x, y, z) \in \mathbb{H}$ ,*

$$|\nabla \log p_t(g)| \leq \frac{Cd(g)}{t}, \tag{1.1}$$

where  $d(g)$  denotes the Carnot-Carathéodory distance between the origin  $o = (0, 0, 0)$  and  $g$ , and  $\nabla f := (Xf, Yf)$ ,  $|\nabla f|^2 := (Xf)^2 + (Yf)^2$ .

Note that in [13–15], Li obtained the precisely upper and lower bounds for the heat kernel  $p_t$  in the Heisenberg (type) group. And the expansion for the heat kernels, Li inequality (i.e.  $|\nabla P_t f| \leq CP_t |\nabla f|$  for some  $C > 0$ ), were also obtained. In the present paper, we will use another method to proof the sharp gradient estimates for the heat kernels. This method relies on the positive property of the Bakry-Émery curvature and some basic facts on groups (see [3, 4]). Our method is more applicable in variant models than the one in [13–15].

In the following, we will study two high dimensional Heisenberg groups — non-isotropic Heisenberg group  $\mathbb{G}$ , and Heisenberg type group  $\mathbb{H}_{n,m}$ . We derive the sharp gradient estimates for the associated heat kernels (see Propositions 2.2 and 3.1).

## 2 Non-isotropic Heisenberg Group

We study the  $(2n + 1)$ -dimensional Heisenberg group  $\mathbb{G}$  (here we use the same symbol as in [5, Subsection 3.4]), normalized as follows. We equip  $\mathbb{R}^{2n} \times \mathbb{R}$  with the group law

$$(x, z) \circ (x', z') = \left( x + x', z + z' + 2 \sum_{j=1}^n a_j [x_{2j} x'_{2j-1} - x_{2j-1} x'_{2j}] \right), \quad x, x' \in \mathbb{R}^{2n}, z, z' \in \mathbb{R},$$

where

$$0 < a_1 \leq a_2 \leq \dots \leq a_l \leq a_{l+1} = \dots = a_n.$$

If  $a_i$  is equal to each other ( $a_1 = a_n$ ), it reduces to the isotropic model. In the isotropic model, the optimal gradient estimates for the heat kernels were obtained by Li [14, 15]. In the non-isotropic case ( $a_1 \neq a_n$ ), although the method used there is not applicable, we will derive the sharp gradient estimate for the heat kernel in this context.

### 2.1 Some basic facts

The left invariant vector fields in  $\mathbb{G}$  are

$$X_{2j-1} = \partial_{2j-1} + 2a_j x_{2j} \partial_z, \quad X_{2j} = \partial_{2j} - 2a_j x_{2j-1} \partial_z, \quad Z = \partial_z,$$

where  $\partial_i = \partial_{x_i}$ . We have the following properties:

- (i) For  $1 \leq j \leq n$ ,  $[X_{2j-1}, X_{2j}] = -4a_j Z$ ,  $[X_{2j-1}, Z] = [X_{2j}, Z] = 0$ ;
- (ii) For  $1 \leq i, j \leq n$ ,  $i \neq j$ ,  $[X_{2i-1}, X_{2j-1}] = [X_{2i-1}, X_{2j}] = [X_{2i}, X_{2j-1}] = [X_{2i}, X_{2j}] = 0$ .

The sublaplace is defined by

$$\Delta = \sum_{j=1}^{2n} X_j^2.$$

Denote the semigroup  $P_t = e^{t\Delta}$ . The heat kernel  $p_t(x, z)$  (density of the semigroup  $P_t$ ) is given by

$$p_t(x, z) = \frac{1}{2(4\pi t)^{n+1}} \int_{-\infty}^{\infty} e^{-\frac{f(x, z, \tau)}{4t}} V(\tau) d\tau,$$

where

$$f(x, z, \tau) = -i\tau z + \sum_{j=1}^n a_j \tau \coth(a_j \tau) r_j^2, \quad V(\tau) = \prod_{j=1}^n \frac{a_j \tau}{\sinh(a_j \tau)},$$

where  $i = \sqrt{-1}$ ,  $r_j^2 = x_{2j-1}^2 + x_{2j}^2$  (see [5]). We have the time scaling property  $p_t(x, z) = t^{-n-1} p_1(\frac{x}{\sqrt{t}}, \frac{z}{t})$ . So it is enough to get the estimates for  $p_1$ .

Set  $x = (\tilde{x}, \hat{x})$  and  $\hat{x} = (x_{2l+1}, x_{2l}, \dots, x_{2n})$ , then the Carnot-Carathéodory distance  $d(x, z)$  from  $(x, z)$  to the origin  $(0, 0)$  is

$$d^2(x, z) = \begin{cases} \sum_{j=1}^n \frac{(2a_j \theta_c)^2 r_j^2}{2 \sin^2(2a_j \theta_c)}, & \text{if } \hat{x} \neq 0, \\ \sum_{j=1}^n \frac{(2a_j \theta_c)^2 r_j^2}{2 \sin^2(2a_j \theta_c)}, & \text{if } |z| < \sum_{j=1}^n a_j \mu \left( \frac{a_j \pi}{a_n} \right) r_j^2, \text{ and } \tilde{x} \neq 0, \hat{x} = 0, \\ \frac{\pi}{a_n} \left( |z| + \sum_{j=1}^n a_j \cot \left( \frac{a_j \pi}{a_n} \right) \right) r_j^2, & \text{if } |z| \geq \sum_{j=1}^n a_j \mu \left( \frac{a_j \pi}{a_n} \right) r_j^2, \text{ and } \tilde{x} \neq 0, \hat{x} = 0, \end{cases}$$

where  $\mu(\phi) = \frac{\phi}{\sin^2 \phi} - \cot \phi$  and  $\theta_c$  is the unique solution of  $|z| = \sum_{j=1}^n a_j \mu(2a_j \theta) r_j^2$  in the interval  $[0, \frac{\pi}{2a_n})$  (see [5, Theorems 3.18, 3.24 and 3.52]). In the first two cases, we clearly have  $d(x, z) \geq |x| \left( |x|^2 := \sum_{j=1}^{2n} x_j^2 \right)$ . In the third case, since  $d(x, z) \simeq |x| + |z|^{\frac{1}{2}}$  (see [5, Theorem 3.52]). (Because  $d(x, z)$  is continuous in this case and  $d(x, z)(|x| + |z|^{\frac{1}{2}})^{-1}$  is homogeneous of degree 0.) In all, we have

$$d(x, z) \geq c|x| \tag{2.1}$$

for some positive constant  $c$ .

Denote

$$\Gamma(f, g) = \frac{1}{2}(\Delta(fg) - g\Delta f - f\Delta g), \quad \Gamma_2(f, f) := \frac{1}{2}(\Delta\Gamma(f, f) - 2\Gamma(f, \Delta f)).$$

Through calculation, we have

$$\begin{aligned} \Gamma(f, g) &= \sum_{i=1}^{2n} (X_i f)(X_i g), \\ \Gamma_2(f, f) &= \sum_{i,j=1}^{2n} (X_j X_i f)^2 + 8 \sum_{k=1}^n a_k (X_{2k-1} f(X_{2k} Z f) - X_{2k} f(X_{2k-1} Z f)). \end{aligned}$$

The mixed term  $\sum_{k=1}^n a_k (X_{2k-1} f(X_{2k} Z f) - X_{2k} f(X_{2k-1} Z f))$  prevents the existence of any constant  $\rho$  such that the curvature dimensional condition  $CD(\rho, \infty)$  holds. Thus the method used in the elliptic context is not applicable.

For any radial function  $f = f(r_1, r_2, \dots, r_n, z)$ , the sublaplace  $\Delta$  has the following form

$$\begin{aligned} \Delta f &= \sum_{i,j=1}^n f_{ij} \Gamma(r_i, r_j) + f_{zz} \Gamma(z, z) + \sum_{i=1}^n f_{iz} \Gamma(r_i, z) + \sum_{i=1}^n f_i \Delta r_i + f_z \Delta z \\ &= \sum_{i=1}^n f_{ii} + 4 \sum_{i=1}^n a_i^2 r_i^2 f_{zz} + \sum_{i=1}^n \frac{1}{r_i} f_i \\ &:= \mathcal{L}f, \end{aligned}$$

where  $\mathcal{L}$  is defined by

$$\mathcal{L} = \sum_{i=1}^n \partial_{ii} + 4 \sum_{i=1}^n a_i^2 r_i^2 \partial_{zz} + \sum_{i=1}^n \frac{1}{r_i} \partial_i.$$

Hence, for radial functions  $f = f(r_1, r_2, \dots, r_n, z)$  and  $g = g(r_1, r_2, \dots, r_n, z)$ ,

$$\begin{aligned} \Gamma(f, g) &:= \frac{1}{2}(\Delta(fg) - f\Delta g - g\Delta f) \\ &= \frac{1}{2}(\mathcal{L}(fg) - f\mathcal{L}g - g\mathcal{L}f) \\ &= \Gamma^{\mathcal{L}}(f, g) \\ &= \sum_{i=1}^n (f_i g_i + 4a_i^2 r_i^2 f_z g_z). \end{aligned}$$

Moreover, through calculation, we have

$$\begin{aligned}
 \Gamma_2(f, f) &:= \frac{1}{2}(\Delta\Gamma(f, f) - 2\Gamma(f, \Delta f)) \\
 &= \frac{1}{2}(\mathcal{L}\Gamma^{\mathcal{L}}(f, f) - 2\Gamma^{\mathcal{L}}(f, \mathcal{L}f)) \\
 &= \Gamma_2^{\mathcal{L}}(f, f) \\
 &= \sum_{i=1}^n f_{ii}^2 + \left(4a_i^2 r_i^2 f_{zz} - \frac{f_z}{r_i}\right)^2 + 8a_i^2 (f_z + r_i f_{zi})^2 \\
 &\quad + \sum_{i \neq j} f_{ij}^2 + 4a_i^2 r_i^2 f_{jz}^2 + 4a_j^2 r_j^2 f_{iz}^2 + 16a_i^2 a_j^2 r_i^2 r_j^2 f_{zz}^2 \\
 &\geq 0.
 \end{aligned} \tag{2.2}$$

Note that the heat kernel  $p_t$  depends only on  $r_1, r_2, \dots, r_n, z$ . So, in particular, we have

$$\Gamma_2(p_t, p_t) \geq 0, \quad \forall t \geq 0.$$

### 2.2 Gradient estimates for the heat kernels

Let us firstly state the Li-Yau type inequality in the non-isotropic Heisenberg group  $\mathbb{G}$ .

**Proposition 2.1** *There exist positive constants  $C_1, C_2, C_3$  such that for any positive function  $f : \mathbb{G} \rightarrow \mathbb{R}^+$ , if  $u = \log P_t f$ , we have*

$$\partial_t u \geq C_1 \Gamma(u) + C_2 t |Zu|^2 - \frac{C_3}{t}. \tag{2.3}$$

**Proof** Following [3], for fixed  $t > 0$ , let  $u_s = P_{t-s} f$ ,  $g_s = \log u_s$ , and

$$\Phi_1(s) = P_s(u_s \Gamma(g_s)), \quad \Phi_2(s) = P_s(u_s (Zg_s)^2).$$

It follows that

$$\Phi_1'(s) = 2P_s(u_s \Gamma_2(g_s, g_s)), \quad \Phi_2'(s) = 2P_s(u_s \Gamma(Zg_s)).$$

Note that

$$\begin{aligned}
 \Gamma_2(f, f) &= \sum_{i,j=1}^{2n} (X_j X_i f)^2 + 8 \sum_{k=1}^n a_k ((X_{2k-1} f)(X_{2k} Zf) - (X_{2k} f)(X_{2k-1} Zf)) \\
 &\stackrel{(*)}{\geq} \frac{1}{2n} (\Delta f)^2 + 8 \sum_{k=1}^n a_k^2 (Zf)^2 + 8 \sum_{k=1}^n a_k ((X_{2k-1} f)(X_{2k} Zf) - (X_{2k} f)(X_{2k-1} Zf)) \\
 &\geq \frac{1}{2n} (\Delta f)^2 + 8 \sum_{k=1}^n a_k^2 (Zf)^2 - 4 \sum_{k=1}^n a_k \left( \frac{1}{\lambda} |\nabla_k f|^2 + \lambda |\nabla_k (Zf)|^2 \right) \\
 &\geq \frac{1}{2n} (\Delta f)^2 + 8 \sum_{k=1}^n a_k^2 (Zf)^2 - 4a_n \left( \frac{1}{\lambda} \Gamma(f) + \lambda \Gamma(Zf) \right),
 \end{aligned}$$

where  $\lambda > 0$  and  $|\nabla_k(f)|^2 := (X_{2k-1} f)^2 + (X_{2k} f)^2$ . Here we use the assumption that for

$1 \leq k \leq n$ ,  $0 < a_k \leq a_n$ , and inequality (\*) follows from

$$\begin{aligned} \sum_{i,j=1}^{2n} (X_j X_i f)^2 &= \sum_{i=1}^{2n} (X_i^2 f)^2 + \sum_{1 \leq i < j \leq 2n} ((X_j X_i f)^2 + (X_i X_j f)^2) \\ &\geq \frac{1}{2n} (\Delta f)^2 + \frac{1}{2} \sum_{1 \leq i < j \leq 2n} (X_i X_j f - X_j X_i f)^2 \\ &= \frac{1}{2n} (\Delta f)^2 + 8 \sum_{k=1}^n a_k^2 (Zf)^2. \end{aligned}$$

Here we use the basic property (i) and (ii) in Subsection 2.1. Since for any  $\gamma \in \mathbb{R}$ ,

$$(\Delta u_s)^2 \geq 2\gamma \Delta u_s - \gamma^2, \quad \Delta g_s = \frac{\Delta u_s}{u_s} - \Gamma(g_s),$$

it yields

$$\Phi'_1(s) \geq -\left(\frac{2\gamma}{n} + \frac{8a_n}{\lambda}\right) \Phi_1(s) + 16 \sum_{k=1}^n a_k^2 \Phi_2(s) - 4\lambda a_n \Phi'_2(s) + \frac{2\gamma}{n} \Delta P_t f - \frac{\gamma^2}{n} P_t f.$$

Let  $a, b$  be two positive functions defined on  $[0, t)$ , with  $b$  decreasing. We have

$$\begin{aligned} (a(s)\Phi_1(s) + b(s)\Phi_2(s))' &\geq \left(a' - \frac{8a_n a}{\lambda} - \frac{2a\gamma}{n}\right) \Phi_1(s) + \left(16 \sum_{k=1}^n a_k^2 a + b'\right) \Phi_2(s) \\ &\quad + (b - 4\lambda a_n a) \Phi'_2(s) + \frac{2\gamma a}{n} \Delta P_t f - \frac{\gamma^2 a}{n} P_t f. \end{aligned}$$

Choose

$$a = -\frac{b'}{16 \sum_{k=1}^n a_k^2}, \quad \lambda = -\frac{4 \sum_{k=1}^n a_k^2 b}{a_n b'}, \quad \gamma = \frac{nb''}{2b'} + \frac{na_n^2 b'}{\sum_{k=1}^n a_k^2 b},$$

such that the coefficients of  $\Phi_1(s)$ ,  $\Phi_2(s)$  and  $\Phi'_2(s)$  in the right-hand side reduce to zero. Then choose  $b(s) = (t - s)^\alpha$  for some  $\alpha > 2$ . Integrating the above differential inequality from 0 to  $t$ , the desired result follows.

As a consequence, we have the following Harnack inequality. There exist positive constants  $A_1$  and  $A_2$ , such that for  $t_2 > t_1 > 0$  and  $g_1, g_2 \in \mathbb{G}$ ,

$$\frac{p_{t_1}(g_1)}{p_{t_2}(g_2)} \leq \left(\frac{t_2}{t_1}\right)^{A_1} e^{A_2 \frac{d^2(g_1, g_2)}{t_2 - t_1}} \tag{2.4}$$

(see [16]).

**Proposition 2.2** *There exists a constant  $C > 0$  such that for  $t > 0$  and  $g = (x, z) \in \mathbb{G}$ ,*

$$\sqrt{\Gamma(\log p_t)(g)} \leq \frac{Cd(g)}{t}, \tag{2.5}$$

where  $d(g)$  denotes the Carnot-Carathéodory distance between 0 and  $g$ .

**Proof** Following [4], for  $0 < s < t$ , let  $\Phi(s) = P_s(p_{t-s} \log p_{t-s})$ . We have

$$\Phi'(s) = P_s(p_{t-s} \Gamma(\log p_{t-s})), \quad \Phi''(s) = 2P_s(p_{t-s} \Gamma_2(\log p_{t-s})).$$

By the fact that  $\Gamma_2(p_t, p_t) \geq 0$  (here we use the positive property of  $\Gamma_2$  on the radial functions), we have that for all  $t \geq 0$ ,  $\Phi''$  is positive, whence  $\Phi'$  is non-decreasing. Thus

$$\int_0^{\frac{t}{2}} \Phi'(s) \, ds \geq \frac{t}{2} \Phi'(0).$$

That is

$$p_t \Gamma(\log p_t) \leq \frac{2}{t} (P_{\frac{t}{2}}(p_{\frac{t}{2}} \log p_{\frac{t}{2}}) - p_t \log p_t).$$

The right-hand side can be bounded by applying the above Harnack inequality (2.4) and the basic fact  $p_{\frac{t}{2}}(g) \leq p_{\frac{t}{2}}(0)$  for all  $g = (x, z) \in \mathbb{G}$ . We have

$$\sqrt{\Gamma(\log p_t)(g)} \leq C \left( \frac{d(g)}{t} + \frac{1}{\sqrt{t}} \right).$$

In particular,

$$\sqrt{\Gamma(\log p_1)(g)} \leq C(d(g) + 1).$$

If  $d(g) \geq 1$ , it is trivial to get the desired result. If  $d(g) \leq 1$ , by (2.1), we have  $|x| \leq \frac{1}{c}$ ,  $g = (x, z)$ .

As in [14], denote

$$W_{1,j} = \int_{\mathbb{R}} \frac{a_j \tau \cosh(a_j \tau)}{\sinh(a_j \tau)} \cdot \prod_{k=1}^n \frac{a_k \tau}{\sinh(a_k \tau)} \, d\tau, \quad W_{2,j} = \int_{\mathbb{R}} a_j \tau \cdot \prod_{k=1}^n \frac{a_k \tau}{\sinh(a_k \tau)} \, d\tau.$$

We have

$$\Gamma(p_1) \leq C \sum_{j=1}^n r_j^2 (|W_{1,j}| + |W_{2,j}|)^2,$$

where  $C$  depends on the dimension  $n$ . From the expression of  $W_{1,j}$  and  $W_{2,j}$ , we can easily obtain that  $|W_{i,j}|_{i=1,2, j=1, \dots, n}$  are bounded. So  $\Gamma(p_1) \leq C_1 |x|^2 \leq C_2 d(g)$  for some positive constants  $C_1$  and  $C_2$ . By the classic estimates on the heat kernel,  $p_1 \simeq 1$ , on  $|d(g)| \leq 1$  (see, for example, [18, Theorems IV 4.2 and IV 4.3]). Thus

$$\Gamma(\log p_1)(g) \leq C d(g).$$

Therefore, we complete the proof by the time scaling property.

### 3 Heisenberg-Type Group $\mathbb{H}_{n,m}$

Heisenberg type group  $\mathbb{H}_{n,m}$  is a nature generalization of Heisenberg group  $\mathbb{H}$ . Let us firstly recall the definition of Heisenberg type groups. It can be seen as  $\mathbb{R}^{n+m}$  with a multiplying law  $\circ$  (refer to [6, 10, 15] and references therein). The multiplier  $\circ$  is defined by

$$(x, t) \circ (\xi, \tau) = \left( \begin{array}{ll} x_j + \xi_j, & j = 1, \dots, n, \\ t_j + \tau_j + \frac{1}{2} \langle x, U^{(j)} \xi \rangle, & j = 1, \dots, m, \end{array} \right), \quad x, \xi \in \mathbb{R}^n, \quad t, \tau \in \mathbb{R}^m,$$

where the matrices  $U^{(1)}, \dots, U^{(m)}$  have the following properties:

(1)  $U^{(j)}$  is an  $n \times n$  anti-symmetric and orthogonal matrix (skew matrix) for every  $j = 1, 2, \dots, m$ ;

(2)  $U^{(i)}U^{(j)} + U^{(j)}U^{(i)} = 0$  for every  $i, j \in \{1, \dots, m\}$  with  $i \neq j$ ,

and  $m, n$  satisfy  $m < \rho(n) = 8p + 2^q$ , where  $n = (2a + 1)2^{4p+q}$  for  $a, p \in \mathbb{N}$  and  $0 \leq q < 3$ . Note that  $n$  is even, and clearly we have  $n \geq \rho(n) \geq m + 1$ . The left invariant vector fields are

$$X_j = \frac{\partial}{\partial x_j} + \frac{1}{2} \sum_{k=1}^m \sum_{i=1}^n x_i U_{i,j}^{(k)} \frac{\partial}{\partial t_k}. \tag{3.1}$$

By using the anti-symmetric of  $U^{(j)}$ , we obtain

$$[X_i, X_j] = \sum_{k=1}^m U_{i,j}^{(k)} \frac{\partial}{\partial t_k} := Z_{ij} \quad \text{for every } i, j \in \{1, \dots, n\}.$$

Also, we have

$$Z_{ij} + Z_{ji} = 0.$$

This yields the commutative relations

$$[X_i, Z_{jk}] = 0 \quad \text{for every } i, j, k \in \{1, \dots, n\}.$$

Note that for the case  $n = 2$  and  $m = 1$ , it reduces to the Heisenberg groups case. Let us introduce the canonical sub-Laplacian operator in the Heisenberg-type groups

$$\Delta := \sum_{j=1}^n X_j^2 = \Delta_x + \frac{1}{4}|x|^2 \Delta_t + \sum_{k=1}^m \langle x, U^{(k)} \nabla_x \rangle \frac{\partial}{\partial t_k},$$

where the  $U^{(k)}$  are as (3.1). Here, we use the notation  $\Delta_x = \sum_{j=1}^n (\frac{\partial}{\partial x_j})^2$ ,  $\Delta_t = \sum_{k=1}^m (\frac{\partial}{\partial t_k})^2$  and  $\nabla_x = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})^T$ . In this model, for the Carnot-Carathéodory distance  $d(g)$  between 0 and  $g = (x, t) \in \mathbb{H}_{n,m}$ , we have  $d(g) \geq |x|$ .

Moreover, for any radial function  $f(r, t)$ ,  $r^2 := \sum_{i=1}^n x_i^2$ , as done in the non-isotropic case,  $\Delta$  has the following form

$$\Delta f = \Delta_x f + \frac{1}{4}|x|^2 \Delta_t f = \frac{\partial^2 f}{\partial r^2} + \frac{n-1}{r} \frac{\partial f}{\partial r} + \frac{1}{4} r^2 \Delta_t f. \tag{3.2}$$

Hence for any radial functions  $f = f(r, t)$  and  $g = g(r, t)$ , we have

$$\Gamma(f, g) = \frac{\partial f}{\partial r} \frac{\partial g}{\partial r} + \frac{1}{4} r^2 \nabla_t f \nabla_t g \tag{3.3}$$



and

$$\begin{aligned}
 \Gamma_2(f, f) &= \left(\frac{\partial^2 f}{\partial r^2}\right)^2 + \frac{n-1}{4}|\nabla_t f|^2 + \frac{1}{2}\left|\nabla_t f + r\nabla_t \frac{\partial f}{\partial r}\right|^2 + \frac{r^4}{16}\left(\sum_{i,j=1}^m \frac{\partial^2 f}{\partial t_i \partial t_j}\right)^2 \\
 &\quad - \frac{r}{2} \frac{\partial f}{\partial r} \Delta_t f + \frac{n-1}{r^2} \left(\frac{\partial f}{\partial r}\right)^2 \\
 &\geq \left(\frac{\partial^2 f}{\partial r^2}\right)^2 + \frac{n-1}{4}|\nabla_t f|^2 + \frac{1}{2}\left|\nabla_t f + r\nabla_t \frac{\partial f}{\partial r}\right|^2 + \frac{r^4}{16m}(\Delta_t f)^2 \\
 &\quad - \frac{r}{2} \frac{\partial f}{\partial r} \Delta_t f + \frac{n-1}{r^2} \left(\frac{\partial f}{\partial r}\right)^2 \\
 &= \left(\frac{\partial^2 f}{\partial r^2}\right)^2 + \frac{n-1}{4}|\nabla_t f|^2 + \frac{1}{2}\left|\nabla_t f + r\nabla_t \frac{\partial f}{\partial r}\right|^2 \\
 &\quad + \left(\frac{r^2}{4\sqrt{m}}\Delta_t f - \frac{\sqrt{m}}{r} \frac{\partial f}{\partial r}\right)^2 + \frac{n-1-m}{r^2} \left(\frac{\partial f}{\partial r}\right)^2 \\
 &\geq 0,
 \end{aligned}$$

where the last inequality follows from the fact that  $n \geq 1 + m$ .

Let  $p_s$  be the heat kernel of the semi-group  $P_s$  starting from the origin  $o$ . Then we have the following form (see [15, 17]):

$$\begin{aligned}
 p_s(x, t) &= (2\pi)^{-m}(4\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^m} \left(\frac{|\lambda|}{\sinh s|\lambda|}\right)^{\frac{n}{2}} \exp\left(-t \cdot \lambda i - \frac{|x|^2|\lambda|}{4} \coth s|\lambda|\right) d\lambda \\
 &= (2\pi)^{-m}(4\pi)^{-\frac{n}{2}} s^{-\frac{n}{2}-m} \int_{\mathbb{R}^m} \left(\frac{|\lambda|}{\sinh |\lambda|}\right)^{\frac{n}{2}} \exp\frac{1}{s}\left(-t \cdot \lambda i - \frac{|x|^2|\lambda|}{4} \coth |\lambda|\right) d\lambda. \quad (3.4)
 \end{aligned}$$

As done above, we can also get the Li-Yau type inequality holds in the Heisenberg type group  $\mathbb{H}_{n,m}$ , hence the Harnack type inequality follows. Also, we have the following sharp gradient estimates for the heat kernel. This proposition was obtained by Li [15] by a different method.

**Proposition 3.1** *There exists a constant  $C > 0$  such that for  $t > 0$  and  $g = (x, t) \in \mathbb{H}_{n,m}$ ,*

$$\sqrt{\Gamma(\log p_t)(g)} \leq \frac{Cd(g)}{t}, \quad (3.5)$$

where  $d(g)$  denotes the Carnot-Carathéodory distance between 0 and  $g$ .

**Acknowledgements** The author would like to express sincere thanks to Professor H.-Q. Li for his helpful discussions, especially for the improvement of Proposition 2.2. The author is also grateful to the anonymous referee for his/her helpful comments and suggestions, which improve the presentation of the paper. The author is greatly indebted to Professor L. M. Wu, Professor D. Bakry and Professor X. D. Li for their constant encouragement and support.

## References

- [1] Bakry, D., On Sobolev and logarithmic Sobolev inequalities for Markov semigroups, *New Trends in Stochastic Analysis*, Charingworth, 1994, World Scientific Publishing, River Edge, 1997, 43–75.
- [2] Bakry, D., Baudoin, F., Bonnefont, M., et al, On gradient bounds for the heat kernel on the Heisenberg group, *J. Funct. Anal.*, **255**, 2008, 1905–1938.

- [3] Bakry, D., Baudoin, F., Bonnefont, M., et al, Subelliptic Li-Yau estimates on three dimensional model spaces, *Potential Theory and Stochastics in Albac*, Theta Series in Advanced Mathematics, 2009.
- [4] Baudoin, F. and Bonnefont, M., The subelliptic heat kernel on  $SU(2)$ : representations, asymptotics and gradient bounds, *Math. Z.*, **263**(3), 2009, 647–672.
- [5] Beals, R., Gaveau, B. and Greiner, P. C., Hamilton-Jacobi theory and the heat kernel on Heisenberg groups, *J. Math. Pures Appl.* (9), **79**(7), 2000, 633–689.
- [6] Bonfiglioli, A. and Uguzzoni, F., Nonlinear Liouville theorems for some critical problems on H-type groups, *J. Funct. Anal.*, **207**(1), 2004, 161–215.
- [7] Davies, E. B., *Heat Kernels and Spectral Theory*, Cambridge Tracts in Mathematics, Vol. 92, Cambridge University Press, New York, 1989.
- [8] Engoulatov, A., A universal bound on the gradient of logarithm of the heat kernel for manifolds with bounded Ricci curvature, *J. Funct. Anal.*, **238**, 2006, 518–529.
- [9] Gaveau, B., Principe de moindre action, propagation de la chaleur et estimées sous-elliptiques sur certains groupes nilpotents, *Acta Math.*, **139**, 1977, 95–153.
- [10] Kaplan, A., Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms, *Trans. Amer. Math. Soc.*, **258**, 1980, 147–153.
- [11] Juillet, N., Geometric inequalities and generalized Ricci bounds on the Heisenberg group, *Int. Math. Res. Not.*, **2009**, 2009, 2347–2373.
- [12] Ledoux, M., The geometry of Markov diffusion generators, *Ann. Fac. Sci. Toulouse Math.* (6), **9**(2), 2000, 305–366.
- [13] Li, H.-Q., Estimation optimale du gradient du semi-groupe de la chaleur sur le groupe de Heisenberg, *J. Funct. Anal.*, **236**(2), 2006, 369–394.
- [14] Li, H.-Q., Estimations asymptotiques du noyau de la chaleur sur les groupes de Heisenberg, *C. R. Math. Acad. Sci. Paris*, **344** 2007, 497–502.
- [15] Li, H.-Q., Estimations optimales du noyau de la chaleur sur les groupes de type Heisenberg, *J. Reine Angew. Math.*, in press, 2010.
- [16] Li, P. and Yau, S. T., On the parabolic kernel of the Schrödinger operator, *Acta Math.*, **156**, 1986, 153–201.
- [17] Randall, J., The heat kernel for generalized Heisenberg groups, *J. Geom. Anal.*, **6**, 1996, 287–316.
- [18] Varopolous, N. Th., Saloff-Coste, L. and Coulhon, T., *Analysis and Geometry on Groups*, Cambridge Tracts in Mathematics, Vol. 100, Cambridge University Press, Cambridge, 1992.