

## Operations on 3-Dimensional Small Covers\*\*

Shintarô KUROKI\*

**Abstract** The purpose of this paper is to study relations among equivariant operations on 3-dimensional small covers. The author gets three formulas for these operations. As an application, the Nishimura's theorem on the construction of oriented 3-dimensional small covers and the Lü-Yu's theorem on the construction of all 3-dimensional small covers are improved. Moreover, for a construction of 3-dimensional 2-torus manifolds, it is shown that all operations can be obtained by using the equivariant surgeries.

**Keywords** Equivariant surgery, Finite group action, Small cover, 3-dimensional manifold, 3-dimensional simple polytope

**2000 MR Subject Classification** 57M50, 57M60, 57S17, 52B10

### 1 Introduction

Small covers were introduced by Davis and Januszkiewicz as a real version of quasitoric manifolds in 1991. In their paper [2], it was shown that there are strong links between a small cover  $M$  with the orbit projection map  $\pi : M \rightarrow P$  and a combinatorial structure of its orbit polytope  $P$ . For example, some topological invariants (e.g., equivariant cohomologies or  $\mathbb{Z}_2$ -Betti numbers) of small covers  $\pi : M \rightarrow P$  are decided by the combinatorial invariants of their orbit polytopes  $P$  (e.g., Stanley-Reisner rings or  $h$ -vectors).

Not only topological invariants but also topological operations on  $M$  (e.g., equivariant connected sums or equivariant surgeries) correspond with combinatorial operations on the orbit polytope  $P$  (see Section 3). Making use of this correspondence, constructions of 3-dimensional small covers  $M^3$  from basic small covers have been studied by Izmistiev, Nishimura, Lü and Yu. In [3], Izmistiev studied the class of small covers  $M^3$  called a linear model, i.e., small covers over 3-colored polytopes. He proved that 3-dimensional linear models are constructed from one basic small cover (the 3-dimensional torus  $T^3$ ) by using two operations  $\sharp$  and  $\natural$  (see Section 3 and Theorem 4.1). In [11], Nishimura generalized Izmistiev's result to oriented small covers  $M^3$ , i.e., small covers over 3 or 4-colored polytopes. He proved that 3-dimensional oriented small covers are constructed from two basic small covers ( $T^3$  and the real projective space  $\mathbb{R}P(3)$ ) by using three operations  $\sharp$ ,  $\natural$  and  $\flat$  (see Section 3 and Theorem 4.2). In [7], Lü and

---

Manuscript received October 23, 2008. Revised March 31, 2009. Published online April 20, 2010.

\*Department of Mathematical Science, KAIST, 335 Gwahangno (373-1 Guseong-dong), Yuseong-gu, Daejeon 305-701, Korea; School of Mathematical Sciences, Fudan University, Shanghai 200433, China.  
E-mail: kuroki@kaist.ac.kr

\*\*Project supported by Fudan University, the Fujiyukai Foundation and Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2009-0063179).

Yu proved that all 3-dimensional small covers are constructed from five basic small covers by using six operations  $\sharp$ ,  $\sharp^e$ ,  $\sharp^{\text{eve}}$ ,  $\natural$ ,  $\sharp^\Delta$  and  $\sharp^C$  (see Section 3 and Theorem 4.3).

The operation  $\flat$  appearing in Theorem 4.2 is not used in Theorem 4.3. On the other hand, the operations  $\sharp^e$ ,  $\sharp^{\text{eve}}$ ,  $\sharp^\Delta$  and  $\sharp^C$  appearing in Theorem 4.3 are not used in Theorem 4.2. So we can naturally ask what relations exist between Theorem 4.2 and Theorem 4.3. Motivated by this question, in this paper, we prove the following theorem.

**Theorem 1.1** *The operations  $\flat$ ,  $\sharp^e$ ,  $\sharp^{\text{eve}}$  can be obtained by using  $\sharp$ ,  $\natural$  as follows:*

- (1)  $\flat = \natural \circ (\sharp^\Delta)^3$ ;
- (2)  $\sharp^e = \natural \circ (\sharp^{P^3(3)})$ ;
- (3)  $\sharp^{\text{eve}} = \natural^2 \circ (\sharp^{P_-^3(3)})$ ,

where  $\natural^2$  denotes two times Dehn surgery  $\natural \circ \natural$ .

Using Theorems 1.1 and 4.2, we have the following corollary.

**Corollary 1.1** *Each 3-dimensional oriented small cover can be (equivariantly) constructed from the real projective space  $\mathbb{RP}(3)$  and the 3-dimensional torus  $T^3$  by using finite times the following two operations: the equivariant connected sum  $\sharp$ , and the equivariant Dehn surgery  $\natural$ .*

Moreover, we prove the following corollary by making use of Theorem 1.1 and Theorem 4.3.

**Corollary 1.2** *Each 3-dimensional small cover can be (equivariantly) constructed from  $\mathbb{RP}(3)$ ,  $M(P^3(3), \lambda_2)$  and  $M(P^3(3), \lambda_3)$  (up to weakly  $\mathbb{Z}_2^3$ -equivariant diffeomorphism) by using finite times the following four operations: the equivariant connected sum  $\sharp$ , the equivariant Dehn surgery  $\natural$ , the operation  $\sharp^\Delta$  and the coloring change  $\sharp^C$ , where  $(M(P^3(3)), \lambda_i)$  denotes  $S^1 \times \mathbb{RP}(2)$  with the  $\mathbb{Z}_2^3$ -action induced by  $\lambda_i$  for  $i = 2, 3$ .*

For 2-torus manifolds, the following theorem holds.

**Theorem 1.2** *The operations  $\sharp^\Delta$ ,  $\sharp^C$  can be obtained by using  $\natural$ ,  $\natural_0$  and  $\sharp$  as follows:*

- (1)  $\sharp^\Delta = \natural_0 \circ \natural \circ \sharp$ ;
- (2)  $\sharp^C P^3(l) = \natural_0 \circ \natural^{l-2} \circ (\sharp^{P^3(l)})$ ,

where  $P^3(l)$  is the  $l$ -sided prism,  $l \geq 3$  and  $\natural^{l-2}$  denotes  $(l-2)$ -times  $\natural \circ \cdots \circ \natural$ .

The organization of this paper is as follows. In Section 2, we recall the basic facts for small covers. In Section 3, we introduce the seven operations on 3-dimensional small covers. In Section 4, we prove the main theorem (i.e. Theorem 1.1). As an application of Theorem 1.1, we also prove Corollaries 1.1 and 1.2 which improve Theorems 4.2 and 4.3. In Section 5, we remark a relation between Theorems 4.2 and 4.3. In Section 6, we introduce a new operation and prove Theorem 1.2 for 2-torus manifolds.

## 2 Basics of Small Covers

In this section, we recall some basic facts for small covers (see [2] for details). We describe the quotient additive group  $\mathbb{Z}/2\mathbb{Z}$  as  $\mathbb{Z}_2$  throughout this paper.

## 2.1 Definition of a small cover

First, we shall recall the terminology: 2-torus manifolds in [5, 6]. A 2-torus manifold  $M^n$  is an  $n$ -dimensional, closed smooth manifold with a non-free effective smooth  $\mathbb{Z}_2^n$ -action.

Let  $P^n$  be a simple convex  $n$ -polytope, i.e., precisely  $n$  facets of  $P^n$  meet at each vertex. A small cover is a 2-torus manifold  $M^n$  which satisfies the following two conditions:

- (a) the  $\mathbb{Z}_2^n$ -action is locally standard, i.e., locally the same as the  $\mathbb{Z}_2^n$ -action on  $\mathbb{R}^n$ ;
- (b) there is the orbit projection map  $\pi : M^n \rightarrow P^n$  constant on  $\mathbb{Z}_2^n$ -orbits which maps every rank  $k$  orbit (i.e., orbits which isomorphic to  $\mathbb{Z}_2^k$ ) to a point in the interior of  $k$ -dimensional face of  $P^n$ ,  $k = 0, \dots, n$ .

We can easily show that  $\pi$  sends  $\mathbb{Z}_2^n$ -fixed points in  $M^n$  to vertices of  $P^n$  by using the above condition (b). We often call  $P^n$  an orbit polytope of  $M$ .

## 2.2 Characteristic function and constructions of small covers

On the other hand, for given  $P^n$ , small covers  $M^n$  with orbit projection  $\pi : M^n \rightarrow P^n$  can be reconstructed by using a characteristic function  $\lambda : \mathcal{F} \rightarrow \mathbb{Z}_2^n$ , where  $\mathcal{F}$  is the set of facets in  $P$ . In this subsection, we recall the characteristic function  $\lambda$  and the construction of small covers by using  $P$  and  $\lambda$ .

Due to the definition of a small cover  $\pi : M \rightarrow P$ , we have that  $\pi^{-1}(\text{int } F^{n-1})$  consists of  $(n-1)$ -rank orbits. In other words, the isotropy subgroup at  $x \in \pi^{-1}(\text{int } F^{n-1})$  is  $K \subset \mathbb{Z}_2^n$  such that  $K \simeq \mathbb{Z}_2$ , where  $\text{int } F^{n-1}$  is an interior of a facet  $F^{n-1}$ . Hence, the isotropy subgroup at  $x$  is determined by a primitive vector  $v \in \mathbb{Z}_2^n$  generating a subgroup  $K$ . In this way, we obtain a function  $\lambda$  from the set of facets of  $P$ , denoted by  $\mathcal{F}$ , to primitive vectors in  $\mathbb{Z}_2^n$ . We call such  $\lambda : \mathcal{F} \rightarrow \mathbb{Z}_2^n$  a characteristic function or coloring on  $P$ . By the locally standard property, a characteristic function satisfies the following property (called the property  $(\star)$ ):

$$\text{if } F_1 \cap \dots \cap F_n \neq \emptyset \text{ for } F_i \in \mathcal{F} \ (i = 1, \dots, n), \text{ then } \{\lambda(F_1), \dots, \lambda(F_n)\} \text{ spans } \mathbb{Z}_2^n. \quad (\star)$$

Next, we mention the construction of small covers by using  $P^n$  and  $\lambda$ . Let  $P^n$  be a simple convex polytope. Suppose that the characteristic function  $\lambda : \mathcal{F} \rightarrow \mathbb{Z}_2^n$  which satisfies the above property  $(\star)$  is defined on  $P^n$ . Small covers can be constructed from  $P$  and  $\lambda$  as follows:

$$\mathbb{Z}_2^n \times P / \sim,$$

where  $(t, x) \sim (t', y)$  is defined as  $x = y \in P$  and

$$\begin{aligned} t &= t', & \text{if } x \in \text{int } P; \\ t^{-1}t' &\in \langle \lambda(F_1), \dots, \lambda(F_l) \rangle \simeq \mathbb{Z}_2^l, & \text{if } x \in \text{int}(F_1 \cap \dots \cap F_l), \end{aligned}$$

where  $\langle \lambda(F_1), \dots, \lambda(F_l) \rangle \subset \mathbb{Z}_2^n$  denotes the subgroup generated by  $\lambda(F_i)$  for  $i = 1, \dots, l$ . We describe such small cover as  $M(P, \lambda)$ .

Before we show examples of small covers, we define the equivalence relation on small covers. Let  $(M_1, \mathbb{Z}_2^n)$  and  $(M_2, \mathbb{Z}_2^n)$  be small covers. We denote their  $\mathbb{Z}_2^n$ -actions as  $\varphi_1$  and  $\varphi_2$ , respectively. We call that  $(M_1, \mathbb{Z}_2^n)$  and  $(M_2, \mathbb{Z}_2^n)$  are weakly equivariantly homeomorphic, if there is a homeomorphism  $f : M_1 \rightarrow M_2$  such that  $f(\varphi_1(t, x)) = \varphi_2(g(t), f(x))$ , where  $t \in \mathbb{Z}_2^n$ ,  $x \in M_1$  and  $g : \mathbb{Z}_2^n \rightarrow \mathbb{Z}_2^n$  is an isomorphism. If  $g$  is the identity map, we call that  $(M_1, \mathbb{Z}_2^n)$  and  $(M_2, \mathbb{Z}_2^n)$  are equivariantly homeomorphic.

### 2.3 Examples

Let  $\{e_1, \dots, e_n\}$  be the standard basis in  $\mathbb{Z}_2^n$ . We call a pair of a polytope and its characteristic function  $(P^n, \lambda)$  a polytope with  $m$ -coloring or an  $m$ -coloring polytope, if the image of  $\lambda$  is a set of  $m$ -independent vectors in  $\mathbb{Z}_2^n$ , i.e.,  $\lambda(\mathcal{F}) = \{f_1, \dots, f_m\}$ , where  $f_j$  ( $j = 1, \dots, m$ ) is a linear combination of  $e_1, \dots, e_n$  and  $f_1, \dots, f_m$  are linearly independent. Now Figure 1 shows two examples of characteristic functions on polytopes. The polytope of the left example is the 3-simplex  $\Delta^3$  with 4-coloring, i.e.,  $\lambda_0(\mathcal{F}) = \{e_1, e_2, e_3, e_1 + e_2 + e_3\}$ . The polytope of the right example is the 3-cube  $I^3$  with 3-coloring, i.e.,  $\lambda_0^I(\mathcal{F}) = \{e_1, e_2, e_3\}$ .




Figure 1 The left figure is  $(\Delta^3, \lambda_0)$  and the right figure is  $(I^3, \lambda_0^I)$ , where the bottom  $e_1 + e_2 + e_3$  and  $e_1$  are the colors of facets on the back. We see that  $M(\Delta^3, \lambda_0) = \mathbb{R}P(3)$  and  $M(I^3, \lambda_0^I) = T^3$ , where  $\mathbb{R}P(3)$  and  $T^3$  have the standard  $\mathbb{Z}_2^3$ -actions.

We can easily show that  $\Delta^3$  has the unique characteristic function  $\lambda_0$  up to  $\text{GL}(3, \mathbb{Z}_2)$ , i.e., for all characteristic function  $\lambda$  on  $\Delta^3$ , there is an element  $\sigma \in \text{GL}(3, \mathbb{Z}_2)$  such that  $\sigma \circ \lambda = \lambda_0$ . In other words, small covers over  $\Delta^3$  are unique up to weakly equivariant diffeomorphisms. We can assume that there is at least one vertex  $v = F_1 \cap \dots \cap F_n$  in  $P$  such that  $\lambda(F_i) = e_i$  for all  $i = 1, \dots, n$  up to  $\text{GL}(3, \mathbb{Z}_2)$  because of the property  $(\star)$ .

A small cover over  $P^n$  with  $n$ -coloring, i.e.,  $\lambda(\mathcal{F}) = \{e_1, \dots, e_n\}$  (up to  $\text{GL}(n, \mathbb{Z}_2)$ ) is called a linear model. So the right example in Figure 1 is a linear model. Nakayama and Nishimura show that a 3-dimensional small cover is orientable if and only if the characteristic function on the orbit polytope  $P^3$  is 3 or 4 colored in [10]. Therefore, small covers which are constructed from the two examples in Figure 1 are orientable.

### 2.4 $\mathbb{Z}_2^3$ -invariant normal bundle over an invariant $S^1$

Let  $\pi : M \rightarrow P$  be a 3-dimensional small cover. In this subsection, we study the equivariant normal bundle of the inverse of edges in  $P$ .

Fix  $\{e_1, e_2, e_3\}$  as a basis (not necessary standard) of  $\mathbb{Z}_2^3$ . We can easily show, by using the property  $(\star)$ , that the characteristic functions on neighboring facets around one edge in a 3-dimensional polytope have only the following four cases in Figure 2.




Figure 2 The characteristic functions around edges

The small cover over a 1-simplex  $\Delta^1 (= I^1)$  is identified with  $\mathbb{R}P(1) (= S^1)$ . Hence, for the small cover  $\pi : M \rightarrow P$ , the inverse  $\pi^{-1}(I^1)$  of any edge  $I^1$  in  $P$  is an invariant submanifold diffeomorphic to  $S^1$ . Moreover, we know an equivariant normal bundle of  $\pi^{-1}(I^1)$  from a characteristic function around  $I^1$ . Let  $(t_1, t_2, t_3) \in \mathbb{Z}_2^3$  act on  $((x_1, x_2), (y_1, y_2)) \in S^1 \times D^2$  by  $((x_1, x_2), (y_1, y_2)) \mapsto ((t_1 x_1, x_2), (t_2 y_1, t_3 y_2))$ . We can easily show the following proposition by computing a characteristic function of  $S^1 \times_{\mathbb{Z}_2} D^2$ 's and using Figure 2 and the construction of small covers in Subsection 2.2.

**Proposition 2.1** *Let  $I$  be an edge in  $P$  and  $N$  be a small neighborhood of  $I$ . For an invariant submanifold  $S^1 = \pi^{-1}(I)$  in the 3-dimensional small cover, the normal bundles  $\pi^{-1}(N)$  are weakly  $\mathbb{Z}_2^3$ -equivariantly isomorphic to one of the following four disk bundles:*

- (1) *if  $N$  satisfies (1) in Figure 2, then  $\pi^{-1}(N) \simeq S^1 \times_{\mathbb{Z}_2} D(\underline{\mathbb{R}} \oplus \underline{\mathbb{R}})$ ;*
- (2) *if  $N$  satisfies (2) in Figure 2, then  $\pi^{-1}(N) \simeq S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R})$ ;*
- (3) *if  $N$  satisfies (3) in Figure 2, then  $\pi^{-1}(N) \simeq S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \underline{\mathbb{R}})$ ;*
- (4) *if  $N$  satisfies (4) in Figure 2, then  $\pi^{-1}(N) \simeq S^1 \times_{\mathbb{Z}_2} D(\underline{\mathbb{R}} \oplus \mathbb{R})$ ,*

*where the non-trivial element in  $\mathbb{Z}_2$  acts on  $S^1$  by the antipodal involution,  $D(V \oplus V') = D^2$  denotes a closed disk in  $V \oplus V'$  ( $V$  and  $V'$  are 1-dimensional real vector spaces), and  $\mathbb{Z}_2$  acts on  $\mathbb{R}$  canonically and on  $\underline{\mathbb{R}}$  trivially.*

**Remark 2.1** In the above Proposition 2.1, we have that  $S^1 \times_{\mathbb{Z}_2} (\underline{\mathbb{R}} \oplus \underline{\mathbb{R}}) \cong \epsilon^1 \oplus \epsilon^1$ ,  $S^1 \times_{\mathbb{Z}_2} (\mathbb{R} \oplus \mathbb{R}) \cong \gamma_1^1 \oplus \gamma_1^1$ ,  $S^1 \times_{\mathbb{Z}_2} (\mathbb{R} \oplus \underline{\mathbb{R}}) \cong \gamma_1^1 \oplus \epsilon^1$  and  $S^1 \times_{\mathbb{Z}_2} (\underline{\mathbb{R}} \oplus \mathbb{R}) \cong \epsilon^1 \oplus \gamma_1^1$ , where  $\epsilon^1$  is the trivial bundle and  $\gamma_1^1$  is the canonical bundle over  $\mathbb{R}P(1) (= S^1)$ . Hence, we have that the bundle (1) (resp. (3)) is isomorphic to the bundle (2) (resp. (4)) in Proposition 2.1, by making use of the basic facts of the vector bundle over  $\mathbb{R}P(1)$  (see [9]). However, these four bundles are different as the  $\mathbb{Z}_2^3$ -equivariant bundle because their characteristic functions are different (see Figure 2). We also remark that, up to weakly  $\mathbb{Z}_2^3$ -equivariant diffeomorphism, (3) and (4) are the same.

### 3 Operations for 3-Dimensional Small Covers

Henceforth, we assume that  $M(P, \lambda)$  is a 3-dimensional small cover over a 3-dimensional simple convex polytope with coloring  $(P, \lambda)$ , and  $\{e_1, e_2, e_3\}$  is a basis (not necessary standard) of  $\mathbb{Z}_2^3$ . In this section, we introduce operations on small covers and orbit polytopes (see also [3, 7, 11]). From this section, we often call the set of colorings included in  $\{f_1, f_2, f_1 + f_2\}$  a 2-independent coloring, where  $f_i$  ( $i = 1, 2$ ) is a linear combination of  $\{e_1, e_2, e_3\}$ .

#### 3.1 The equivariant connected sum $\sharp$

The operation in Figure 3 is called the equivariant connected sum  $\sharp$  (from left to right) and its inverse  $\sharp^{-1}$  (from right to left). The left figure shows two neighborhoods of two same colored vertices in  $(P_1, \lambda_1)$  and  $(P_2, \lambda_2)$ . We can do the connected sum of the two polytopes at these vertices (see [3, Definition 3]). Then we get a new polytope with coloring  $(P_1 \sharp P_2, \lambda)$ , and vice versa. Remark that  $P_1 \sharp P_2$  is a combinatorial simple convex polytope by using the Steinitz's theorem: the graph  $\Gamma$  is a graph of the 3-dimensional polytope  $P$  if and only if  $\Gamma$  is 3-connected and planar (see [13, Chapter 4]).

From the geometric point of view, this operation  $\sharp$  corresponds with the equivariant connected sum  $M(P_1, \lambda_1) \sharp M(P_2, \lambda_2)$  for two fixed points in  $M(P_1, \lambda_1)$  and  $M(P_2, \lambda_2)$ . We can

easily check that  $M(P_1, \lambda_1) \sharp M(P_2, \lambda_2)$  is a small cover and its orbit polytope with coloring is  $(P_1 \sharp P_2, \lambda)$ , i.e.,  $M(P_1, \lambda_1) \sharp M(P_2, \lambda_2) = M(P_1 \sharp P_2, \lambda)$  (see also [3, Lemma 2]).




Figure 3 The equivariant connected sum  $\sharp$  (from left to right) and its inverse  $\sharp^{-1}$  (from right to left). Here,  $e_3$  is the coloring of the facet on the back.

### 3.2 The cutting edge operation $\sharp^e$

Before we mention the cutting edge operation, we introduce the connected sum along edges. Let  $P_1$  and  $P_2$  be 3-dimensional, simple, convex polytopes. Suppose that the edges  $I_1 \subset P_1$  and  $I_2 \subset P_2$  are chosen, and a one-to-one correspondence  $F_i \mapsto F'_i$  ( $i = 1, \dots, 4$ ) is established, where this correspondence is from the facets  $\{F_1, F_2, F_3, F_4\}$  containing  $I_1$  as  $I_1 = F_2 \cap F_3$  to the facets  $\{F'_1, F'_2, F'_3, F'_4\}$  containing  $I_2$  as  $I_2 = F'_2 \cap F'_3$ . The connected sum along edges with respect to these data is a polytope combinatorially equivalent to the result of the gluing  $P_1$  and  $P_2$  with small neighborhoods of  $I_1$  and  $I_2$  removed. The corresponding facets must be glued together. The operation in Figure 4 is the special case of this operation.

The operation in Figure 4 is called the cutting edge operation  $\sharp^e$  (from left to right) and its inverse  $(\sharp^e)^{-1}$  (from right to left). The left figure shows two neighborhoods of two edges whose neighboring facets have a same coloring in  $(P_1, \lambda_1)$  and  $(P_2, \lambda_2)$ , where  $P_2$  is the 3-sided prism, i.e.,  $P_2 = P^3(3) = I^1 \times \Delta^2$ . We can do the connected sum along these edges, then we get a new polytope (remark that this is combinatorially equivalent to a simple polytope because of the Steinitz' theorem) with coloring  $(P_1 \sharp^e P_2, \lambda)$ , and vice versa (see also [7, Section 2]).




Figure 4 The cutting edge operation  $\sharp^e$  (from left to right) and its inverse  $(\sharp^e)^{-1}$  (from right to left). Here,  $e_3$  is the coloring of the facet on the back, and  $w$  is an element in  $\mathbb{Z}_2^3$  such that the property  $(\star)$  holds around vertices (see Figure 2).

From the geometric point of view, the operation  $\sharp^e$  corresponds with the following operation. Let  $\pi_i : M_i \rightarrow P_i$  be a small cover and  $I_i$  an edge in  $P_i$  for  $i = 1, 2$ . Suppose that  $P_2 = P^3(3)$  and colorings of facets around  $I_1$  and  $I_2$  are the same as in Figure 4. Then we see that a closed invariant tubular neighborhood  $N_1$  of  $\pi_1^{-1}(I_1)$  is equivariantly isomorphic to a closed invariant tubular neighborhood  $N_2$  of  $\pi_2^{-1}(I_2)$  (see Subsection 2.4). Therefore, two boundaries of  $M(P_1, \lambda_1) \setminus \text{int } N_1$  and  $M(P_2, \lambda_2) \setminus \text{int } N_2$  are equivariantly diffeomorphic, where  $\text{int } N_i$  is the interior of  $N_i$  for  $i = 1, 2$ . Hence, we can glue equivariantly these two boundaries, and get the

$\mathbb{Z}_2^3$ -manifold  $M(P_1, \lambda_1) \setminus \text{int } N_1 \cup_{\partial} M(P_2, \lambda_2) \setminus \text{int } N_2$ ; we denote it by  $M(P_1, \lambda_1) \#^e M(P_2, \lambda_2)$ . We can easily show that  $M(P_1, \lambda_1) \#^e M(P_2, \lambda_2)$  is small cover and its orbit polytope with coloring is  $(P_1 \#^e P_2, \lambda)$  (see also [7, Section 5.2]).

### 3.3 The cutting edge-vertex-edge operation $\#^{\text{eve}}$

In this paper, the operation in Figure 5 is called the cutting e-v-e operation  $\#^{\text{eve}}$  (from left to right) and its inverse  $(\#^{\text{eve}})^{-1}$  (from right to left), where e-v-e means edge-vertex-edge. Let 2-edges in a simple polytope  $P$  be a union of two different edges with the common vertex. The left figure shows two neighborhoods of two same colored 2-edges in  $(P_1, \lambda_1)$  and  $(P_2, \lambda_2)$ , where  $P_2$  is the truncated prism, i.e.,  $P_2$  is the polytope constructed by the connected sum of the 3-simplex  $\Delta^3$  and the 3-sided prism  $P_3^3$ :  $P_2 = P_-^3(3) = P^3(3) \# \Delta^3$ . We denote the 2-edges in  $(P_1, \lambda_1)$  as  $I_1 \vee I_2$  and that in  $(P_2, \lambda_2)$  as  $I'_1 \vee I'_2$ . Then we can establish a one-to-one correspondence  $F_i \mapsto F'_i$  ( $i = 1, \dots, 5$ ), from the facets  $\{F_1, \dots, F_5\}$  containing  $I_1 \vee I_2$  as  $I_1 = F_2 \cap F_5$  and  $I_2 = F_3 \cap F_5$  to the facets  $\{F'_1, \dots, F'_5\}$  containing  $I'_1 \vee I'_2$  as  $I'_1 = F'_2 \cap F'_5$  and  $I'_2 = F'_3 \cap F'_5$ , such that  $\lambda_1(F_i) = \lambda_2(F'_i)$ . The cutting e-v-e operation with respect to these data is a polytope combinatorially equivalent to the result of the gluing  $P_1$  and  $P_2$  with small neighborhoods of  $I_1 \vee I_2$  and  $I'_1 \vee I'_2$  removed. The corresponding facets must be glued together. Remark that the cutting e-v-e operation of  $P_1$  and combinatorially equivalent to a simple polytope because of the Steinitz's theorem. Therefore, we have the new polytope with coloring  $(P_1 \#^{\text{eve}} P_2, \lambda)$  (see also [7, Section 2]).




Figure 5 The cutting e-v-e operation  $\#^{\text{eve}}$  (from left to right) and its inverse  $(\#^{\text{eve}})^{-1}$  (from right to left). Here,  $e_3$  is the coloring of the facet on the back, and  $w_1, w_2$  are elements in  $\mathbb{Z}_2^3$  such that the property  $(\star)$  holds around vertices.

From the geometric point of view, this operation  $\#^{\text{eve}}$  corresponds with the following operation. Let  $\pi_i : M_i \rightarrow P_i$  be a small cover and  $I_i$  an edge in  $P_i$  for  $i = 1, 2$ . Assume that  $P_2 = P_-^3(3)$  and colorings of facets around  $I_1 \vee I_2$  in  $P_1$  and  $I'_1 \vee I'_2$  in  $P_2$  are the same as in Figure 5. Now we see that  $\pi_1^{-1}(I_1 \vee I_2) \cong S^1 \vee S^1$  and  $\pi_2^{-1}(I'_1 \vee I'_2) \cong S^1 \vee S^1$ . Here,  $S^1 \vee S^1$  is the  $\mathbb{Z}_2^3$ -invariant bouquet of two  $S^1$ 's on the fixed points, i.e.,  $S^1 \vee S^1 = (S^1 \sqcup S^1)/v_1 \sim v_2$ , where  $S^1 \sqcup S^1$  is a disjoint union of two  $S^1$ 's and  $v_i$  is a  $\mathbb{Z}_2^3$ -fixed point in the  $i$ -th  $S^1$  for  $i = 1, 2$ . Moreover, by our assumption, there are  $\mathbb{Z}_2^3$ -invariant closed neighborhoods  $N_1$  and  $N_2$  of  $\pi_1^{-1}(I_1 \vee I_2)$  and  $\pi_2^{-1}(I'_1 \vee I'_2)$ , respectively, and they are equivariantly diffeomorphic. Therefore, with a method similar to that demonstrated in Subsection 3.3, we have the manifold  $M(P_1, \lambda_1) \setminus \text{int } N_1 \cup_{\partial} M(P_2, \lambda_2) \setminus \text{int } N_2$  with a  $\mathbb{Z}_2^3$ -action; we denote it by  $M(P_1, \lambda_1) \#^{\text{eve}} M(P_2, \lambda_2)$ . We can easily show that  $M(P_1, \lambda_1) \#^{\text{eve}} M(P_2, \lambda_2)$  is a small cover and its orbit polytope with coloring is  $(P_1 \#^{\text{eve}} P_2, \lambda)$  (see [7, Section 5.3] for detail).



### 3.4 The equivariant Dehn surgery $\natural$

The operation described in Figure 6 is called the equivariant Dehn surgery  $\natural$  (from left to right) and its inverse  $\natural^{-1}$  (from right to left). If a coloring around an edge is 3-coloring (i.e., from geometric point of view, an equivariant normal bundle which is weakly equivariantly isomorphic to  $S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R})$  by Proposition 2.1), then we can do this operation  $\natural$  for  $(P, \lambda)$ . Remark that the obtained object by  $\natural$  might not be a convex polytope (see also [3]); however, we get the coloring on this object which satisfies the property  $(\star)$  around a vertex. We denote the object obtained by the operation  $\natural$  as  $\natural(P, \lambda)$ .




Figure 6 The equivariant Dehn surgery  $\natural$  (from left to right) and its inverse  $\natural^{-1}$  (from right to left)

From the geometric point of view, this operation  $\natural$  corresponds with the following operation. Let  $\mathbb{Z}_2^3$  act on the last three coordinates in  $S^3 \subset \mathbb{R} \oplus \mathbb{R}^3$ , naturally. Then there is the  $\mathbb{Z}_2^3$ -invariant submanifold  $S^1 \subset \mathbb{R}^2 \oplus \{0\} \oplus \{0\}$  (remark that there are two fixed points in this submanifold), and we can take its tubular neighborhood  $S^1 \times D^2 \cong S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R})$ . We next consider  $S^3 = S^1 \times D^2 \cup D^2 \times S^1$ , and prepare the  $\mathbb{Z}_2^3$ -invariant part  $S^3 \setminus S^1 \times D^2 = D^2 \times S^1$  (remark that there is no fixed points in this manifold). Then we remove the invariant neighborhood  $S^1 \times D^2 (\simeq S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R}))$  around  $S^1$  from  $M(P, \lambda)$ , i.e., we take  $M(P, \lambda) \setminus S^1 \times D^2$ . Finally, we glue these two invariant manifold  $M(P, \lambda) \setminus S^1 \times D^2 \cup_{\partial} S^3 \setminus S^1 \times D^2$ . This operation is identically the equivariant Dehn surgery of three dimensional manifold (see also [3, Definition 5] or [7, Section 5.4]). We denote the manifold obtained by this equivariant surgery as  $\natural(M(P, \lambda))$ . If  $\natural(P, \lambda)$  is a convex polytope, then  $\natural(M(P, \lambda)) = M(\natural(P, \lambda))$ .

### 3.5 The equivariant Dehn surgery $\flat$

The operation described in Figure 7 is called the equivariant Dehn surgery  $\flat$  (from left to right) and its inverse  $\flat^{-1}$  (from right to left). This move is also called the bistellar 1-move (see [1, Chapter 2]). Remark that the operation in Figure 7 corresponds with the dual of the bistellar 1-move in [1]). If a coloring around an edge is 4-coloring as that in Figure 7 (i.e., from geometric point of view, an equivariant normal bundle which is weakly equivariantly isomorphic to  $S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R})$  by Proposition 2.1), then we can do this operation  $\flat$ . We describe the object with coloring obtained by this operation as  $\flat(P, \lambda)$ . Remark that if the polytope  $P$  is prime, i.e,  $P$  is not decomposed into connected sum of two different polytopes, then this operation  $\flat$  does not destroy the convex property (see [11]). We also remark that this move is the same as the connected sum along edges between  $(P, \lambda)$  and  $(\Delta^3, \sigma \circ \lambda_0)$  for some  $\sigma \in \text{GL}(3; \mathbb{Z}_2)$ .

Now we may explain what happens from the geometric point of view. We can regard  $\mathbb{R}P(3)$  as  $S^3/\mathbb{Z}_2$  by the antipodal involution  $\mathbb{Z}_2$ . Then we can consider  $\mathbb{R}P(3) = S^1 \times_{\mathbb{Z}_2} D^2 \cup D^2 \times_{\mathbb{Z}_2} S^1$ , where  $S^1 \times_{\mathbb{Z}_2} D^2 \simeq S^1 \times_{\mathbb{Z}_2} D(\mathbb{R} \oplus \mathbb{R})$ . We first prepare the  $\mathbb{Z}_2^3$ -invariant part  $\mathbb{R}P(3) \setminus S^1 \times_{\mathbb{Z}_2} D^2 = D^2 \times_{\mathbb{Z}_2} S^1$  (remark that there are two fixed points in this manifold). Next, we remove the invariant neighborhood  $S^1 \times_{\mathbb{Z}_2} D^2$  around  $S^1$  from  $M(P, \lambda)$ , i.e., we take  $M(P, \lambda) \setminus S^1 \times_{\mathbb{Z}_2} D^2$ .



Finally, we glue these two invariant manifold  $M(P, \lambda) \setminus S^1 \times_{\mathbb{Z}_2} D^2 \cup_{\partial} \mathbb{R}P(3) \setminus S^1 \times_{\mathbb{Z}_2} D^2$ . We denote the manifold obtained by this operation as  $\flat(M(P, \lambda))$ . If  $\flat(P, \lambda)$  is a convex polytope, then  $\flat(M(P, \lambda)) = M(\flat(P, \lambda))$ .




Figure 7 The equivariant surgery  $\flat$  (from left to right) and its inverse  $\flat^{-1}$  (from right to left)

### 3.6 The operation $\sharp^\Delta$

The operation described in Figure 8 is called the operation  $\sharp^\Delta$  (from left to right) and its inverse  $(\sharp^\Delta)^{-1}$  (from right to left). If a coloring around a triangle facet is 2-independent coloring (i.e., their colorings can be chosen from one of  $\{e_1, e_2, e_1 + e_2\}$ ), then we can do this operation  $\sharp^\Delta$ . Using the Steinitz's theorem,  $P_1 \sharp^\Delta P_2$  is a convex, simple polytope (see also [7, Section 2.2]).




Figure 8 The operation  $\sharp^\Delta$  (from left to right) and its inverse  $(\sharp^\Delta)^{-1}$  (from right to left)

From the geometric point of view, a neighborhood of a triangle facet whose neighboring facets are 2-independent coloring corresponds with an invariant normal bundle which is weakly equivariantly isomorphic to  $\mathbb{R}P(2) \times I^1$  with the standard  $\mathbb{Z}_2^3$ -action (i.e., the first  $\mathbb{Z}_2^2$  acts on  $\mathbb{R}P(2)$  and the last  $\mathbb{Z}_2$  acts on  $I^1 = [-1, 1] \subset \mathbb{R}$  naturally), by computing its characteristic function. Therefore, we have that the operation  $\sharp^\Delta$  corresponds with the following operation. We first remove an open invariant neighborhood  $\mathbb{R}P(2) \times \text{int } I^1$  from  $M(P_i, \lambda_i)$ , i.e.,  $M(P_i, \lambda_i) \setminus \mathbb{R}P(2) \times \text{int } I^1$  for  $i = 1, 2$ . Next we glue these two manifolds along boundaries, i.e.,  $M(P_1, \lambda_1) \setminus \mathbb{R}P(2) \times \text{int } I^1 \cup_{\partial} M(P_2, \lambda_2) \setminus \mathbb{R}P(2) \times \text{int } I^1$ ; we denote it by  $M(P_1, \lambda_1) \sharp^\Delta M(P_2, \lambda_2)$ . We can easily show that  $M(P_1, \lambda_1) \sharp^\Delta M(P_2, \lambda_2)$  is a small cover and its orbit polytope with coloring is  $(P_1 \sharp^\Delta P_2, \lambda)$  (see also [7, Section 5.5]).

### 3.7 The coloring change $\sharp^C$

We explain the operation in Figure 9. Let  $F$  be an  $l$ -gon facet whose neighboring facets are 2-independent coloring in  $(P, \lambda)$  (see the left bottom polytope in Figure 9). Then we can construct an  $l$ -sided prism  $P^3(l) = F \times I^1$  (see the left above polytope in Figure 9), which naturally admits a coloring such that the coloring of the neighboring facets around the bottom facet (or top facet) is the same as that of  $F$  in  $(P, \lambda)$ . Next, we glue these two polytopes along the facet  $F$ , and we have the new polytope; we denote this new polytope by  $P \sharp^C P^3(l)$ . Remark that two polytopes  $P$  and  $P \sharp^C P^3(l)$  are combinatorially equivalent. However, the colorings of the  $l$ -gon facet  $F$  in  $P$  and  $P \sharp^C P^3(l)$  are different (see also [7, Section 2.3]).

From the geometric point of view, this operation corresponds with a (geometric) operation similar to that described in Subsection 3.6 (see [7, Section 5.6] for detail).




Figure 9 The coloring change  $\sharp^C$  (from left to right) and its inverse  $(\sharp^C)^{-1}$  (from right to left), where  $w = e_1$  or  $e_1 + e_2$  and the left  $e_3$  means the coloring of the bottom of prism and the top of polytope. We also have  $x = e_1 + e_3$ ,  $e_2 + e_3$  or  $e_1 + e_2 + e_3$ .

## 4 Main Theorem and Corollaries

In this section, we prove the main theorem of this paper.

### 4.1 Proof of the main theorem

**Proof of Theorem 1.1** The relations (1)–(3) are shown by the Figures 10–12, respectively.






Figure 10  $\flat = \natural \circ (\sharp \Delta)$

In these figures, the upper map from left to right means the connected sum  $\sharp$  on two black vertices with  $\Delta$ ,  $P^3(3)$  and  $P_-^3(3)$ , respectively, and the right map from top to bottom means the equivariant Dehn surgery  $\natural$  along the black edge. Remark that in Figure 12, we do the operation  $\natural$  two times along two black edges. As the result, we get the formulas (1)–(3) in the statement.

Figure 11  $\#^e = \# \circ (\#P^3(3))$ Figure 12  $\#^{\text{eve}} = \#^2 \circ (\#P^3(3))$ 

## 4.2 Constructions of oriented small covers

In this and next Subsections, we apply our main theorem (i.e., Theorem 1.1) to constructions of the 3-dimensional small covers. First, we recall the following Izmistiev's theorem (see [3, Theorem 3]).

**Theorem 4.1** (Izmestiev) *Suppose that the characteristic function  $\lambda$  of a 3-dimensional small cover  $M(P, \lambda)$  satisfies that  $\lambda(\mathcal{F}) = \{e_1, e_2, e_3\}$ , i.e.,  $M(P, \lambda)$  is a 3-dimensional linear model. Then  $M(P, \lambda)$  can be (equivariantly) constructed from the 3-dimensional torus  $T^3$  by using finite times the following two operations: the equivariant connected sum  $\#$ , and the equivariant Dehn surgery  $\natural$ .*

Nishimura generalizes the above theorem to the following theorem (see [11, Theorem 1.10]).

**Theorem 4.2** (Nishimura) *Suppose that the characteristic function  $\lambda$  of a 3-dimensional small cover  $M(P, \lambda)$  satisfies that  $\lambda(\mathcal{F}) \subset \{e_1, e_2, e_3, e_1 + e_2 + e_3\}$ , i.e.,  $M(P, \lambda)$  is an oriented small cover. Then  $M(P, \lambda)$  can be (equivariantly) constructed from  $T^3$  and the 3-dimensional real projective space  $\mathbb{R}P(3)$  by using finite times the following three operations: the equivariant connected sum  $\#$ , and the equivariant Dehn surgeries  $\natural$  and  $\flat$ .*

By Subsection 2.3, we see that the small cover over  $\Delta^3$  is  $\mathbb{R}P(3)$ . Therefore Corollary 1.1 can be proved by applying Theorem 1.1(1) to the above Nishimura's theorem.

### 4.3 Constructions of all small covers

For all 3-dimensional small covers, the following L  -Yu's theorem are known (see [7, Theorems 1.1 and 1.2]).

**Theorem 4.3** (L  -Yu) *Each 3-dimensional small cover can be (equivariantly) constructed from  $\mathbb{R}P(3)$  and  $S^1 \times \mathbb{R}P(2)$  with certain four type  $\mathbb{Z}_2^3$ -actions (up to weakly equivariant diffeomorphism) by using the following six operations:  $\sharp$ ,  $\sharp^e$ ,  $\sharp^{\text{eve}}$ ,  $\natural$ ,  $\sharp^\Delta$  and  $\sharp^C$ .*

**Remark 4.1** In this paper, we abuse the notations of the operations on small covers and on polytopes.

We shall explain the four types  $\mathbb{Z}_2^3$ -actions on  $S^1 \times \mathbb{R}P(2)$  in Theorem 4.3 (see [7, Lemmas 4.2 and 4.3] for detail). The manifold  $S^1 \times \mathbb{R}P(2)$  is the small cover over the three sided prism  $P^3(3) = I^1 \times \Delta^2$ . The four types  $\mathbb{Z}_2^3$ -actions are defined by using the coloring in Figure 13. We call them (from left-hand side)  $M(P^3(3), \lambda_1)$ ,  $M(P^3(3), \lambda_2)$ ,  $M(P^3(3), \lambda_3)$ ,  $M(P^3(3), \lambda_4)$ , respectively.




Figure 13 Basic four  $\mathbb{Z}_2^3$ -actions on  $S^1 \times \mathbb{R}P(2)$  in Theorem 4.3;  $M(P^3(3), \lambda_1)$ ,  $M(P^3(3), \lambda_2)$ ,  $M(P^3(3), \lambda_3)$ ,  $M(P^3(3), \lambda_4)$ . Here,  $e_2 + e_3$  is the coloring of facets on the back.

Remark that colorings of neighboring facets of triangle facets in Figure 13 are 2-independent. Therefore, we can do  $\sharp^\Delta$  (see Subsection 3.6) for these manifolds as Figure 14.




Figure 14  $M(P, \lambda_1)$  and  $M(P, \sigma \circ \lambda_4)$  can be constructed from  $M(P, \lambda_2)$  and  $M(P, \lambda_3)$  by using  $\sharp^\Delta$ . This figure shows that we do  $\sharp^\Delta$  along black facets.

In Figure 14, we remark that the middle two  $P^3(3)$ 's are equivariantly diffeomorphic to  $M(P, \lambda_2)$  because there is the combinatorially equivalent to the left-second figure (i.e.,  $M(P^3(3), \lambda_2)$ ) in Figure 13 which preserves the colorings (from the geometric point of view, this equivalence on polytopes corresponds with the equivariant diffeomorphisms on small covers). We can easily show that the coloring of the right bottom figure in Figure 14 is the same as the right figure (i.e.,  $M(P^3(3), \lambda_4)$ ) in Figure 13 up to  $\text{GL}(3, \mathbb{Z}_2)$ -equivariant isomorphism, by using the following  $\sigma \in \text{GL}(3, \mathbb{Z}_2)$ :

$$\sigma = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

By the above argument, we have the following lemma.

**Lemma 4.1** *For the basic small covers in Theorem 4.3 (see Figure 13), the following relations hold:*

$$\begin{aligned} (P^3(3), \lambda_1) &= (P^3(3), \lambda_2) \sharp^\Delta (P^3(3), \lambda_2), \\ (P^3(3), \sigma \circ \lambda_4) &= (P^3(3), \lambda_2) \sharp^\Delta (P^3(3), \lambda_3). \end{aligned}$$

Next we remark that the operation  $\sharp^e$  (resp.  $\sharp^{\text{eve}}$ ) itself decomposed as the connected sum of  $P^3(3)$  (resp.  $P^3(3)$ ) with Dehn surgery. However, the 3-sided prism with coloring  $(P^3(3), \lambda)$  which is not included in Figure 13 can be constructed as  $(\Delta, \lambda_0) \sharp (\Delta, \lambda_0)$  by using the argument in [7, Section 3]. Moreover, all truncated prisms with colorings  $(P^3_-(3), \lambda)$  can be constructed as  $(\Delta, \lambda_0) \sharp (P^3(3), \lambda)$  by using the argument in [7, Section 3(a)]. Hence, we get Corollary 1.2 by applying Theorem 1.1(2), (3) and Lemma 4.1 to Theorem 4.3.

## 5 Relation between the Nishimura's Theorem and the Lü-Yu's Theorem

In this section, we give the relations among the Izmitiev's, Nishimura's and Lü-Yu's theorems.

### 5.1 Relation of Corollary 1.1 to the operation $\sharp^C$

In this subsection, we apply the operation  $\sharp^C$  to Corollary 1.1 for oriented small covers.

The torus with standard  $\mathbb{Z}_2^3$ -action  $T^3 = M(I^3, \lambda_0^I)$  can be constructed from  $(\Delta^3, \lambda_0)$  by using  $\sharp$ ,  $\natural$  and  $\sharp^C$  as Figure 15.

In Figure 15, the first figure shows the connected sum on the black vertices, then we get  $((\Delta^3, \lambda_0) \sharp (\Delta^3, \lambda_0) = (P^3(3), \lambda)$ . The second figure shows the connected sum along the black edges: this operation is identical with  $\sharp^e = \natural \circ (\sharp P^3(3))$ , and we get  $(P^3(3), \lambda) \sharp^e (P^3(3), \lambda) = (I^3, \lambda')$ . In  $(I^3, \lambda')$  (the second right figure  $I^3$ ), the colorings around the square facets with coloring  $w$  are 2-independent; therefore, we can do  $\sharp^C$  along this facets. The third figure shows this coloring change, i.e., the coloring change along the black square facets and we get  $(I^3, \lambda') \sharp^C (I^3, \lambda') = (I^3, \lambda_0^I)$ .

Therefore, we have the following proposition by applying the above argument to Corollary 1.1.

**Proposition 5.1** *Each 3-dimensional oriented small cover can be (equivariantly) constructed from  $\mathbb{RP}(3)$  by using finite times the following three operations: the equivariant connected sum  $\sharp$ , the equivariant Dehn surgery  $\natural$ , and the coloring change  $\sharp^C$ .*




Figure 15  $(I^3, \lambda_0^I)$  can be constructed from  $(\Delta^3, \lambda_0)$  by using  $\sharp$ ,  $\natural$  and  $\sharp^C$ , where  $w = e_1 + e_2 + e_3$ .

## 5.2 Problem

The above Proposition 5.1 shows that Corollary 1.2 is not the (direct) generalization of Corollary 1.1. Now we may give the relations among Theorem 4.1, Corollaries 1.1, 1.2 and Proposition 5.1 as the following list.

Table 1 Relations among Theorem 4.1, Corollaries 1.1, 1.2 and Proposition 5.1

Linear model	Oriented	3-dimensional small covers
Izmestiev (Theorem 4.1)	Nishimura (Corollary 1.1) Proposition 5.1	? Lü-Yu (Corollary 1.2)

Here, the column in the list means the category of 3-dimensional small covers, and this list means that Corollary 1.1 is the generalization of Theorem 4.1 and Corollary 1.2 is the generalization of Proposition 5.1. So we can ask the following problem.

**Problem 5.1** *What is the generalization of Corollary 1.1 for 3-dimensional small covers? In other words, what are basic small covers which construct all 3-dimensional small covers by using operations  $\sharp$ ,  $\natural$  (or  $\sharp^\Delta$ )?*

## 6 On 2-Torus Manifolds

In this final section, we shall give some remark for 2-torus manifolds. A 2-torus manifold  $M^n$  is an  $n$ -dimensional, closed smooth manifold with a non-free effective smooth  $\mathbb{Z}_2^n$ -action

(see Subsection 2.1). In this paper, we are interested in the  $n = 3$  case. First we give some basic facts for 2-torus manifolds (see [5, 6] for detail).

### 6.1 Basics of 2-torus manifolds

Let  $\pi : M^3 \rightarrow M^3/\mathbb{Z}_2^3$  be the orbit projection of the 2-torus manifold  $M$ . We see that the orbit space  $M^3/\mathbb{Z}_2^3$  is a 3-dimensional closed space. If  $\mathbb{Z}_2^3$ -action is locally standard, then the orbit space is a 3-dimensional manifold with corner. Moreover, we can define the cell decomposition on  $\partial M^3/\mathbb{Z}_2^3$  (we call it the orbit cell decomposition) induced from the information of  $\mathbb{Z}_2^3$ -orbits as follows. The information of fixed points as 0-cells (vertices), the information of rank one-orbits (i.e., orbits  $\mathbb{Z}_2^3/K \simeq \mathbb{Z}_2$ ) as 1-cells (edges), the information of rank two-orbits (i.e., orbits  $\mathbb{Z}_2^3/K \simeq \mathbb{Z}_2^2$ ) as 2-cell (facets). We remark that the free orbits correspond with the interior of  $M^3/\mathbb{Z}_2^3$ . With a method similar to that defines the characteristic function on the small cover, we can define the characteristic function from facets in  $\partial M^3/\mathbb{Z}_2^3$  to  $\mathbb{Z}_2^3$ . For example,  $\mathbb{Z}_2^3$  acts canonically on the last three coordinates of  $S^3 \subset \mathbb{R} \oplus \mathbb{R}^3$ . Then  $S^3$  is a 2-torus manifold (not a small cover), and its orbit space with characteristic function is as that in Figure 16 by computing isotropy subgroups on rank two-orbits.




Figure 16 The orbit cell decomposition on  $S^3/\mathbb{Z}_2^3$ . We describe this orbit cell decomposition with coloring as  $(D^3, \rho)$ .

Figure 16 shows that  $S^3/\mathbb{Z}_2^3$  becomes the 3-disk  $D^3$ , and its orbit cell decomposition is as follows: two vertices, three edges, and three facets. We remark that, in Figure 16,  $e_3$  is the coloring of the facet on the back.

We denote the orbit cell decomposition with coloring in Figure 16 as  $(D^3, \rho)$ .

### 6.2 The equivariant surgery $\natural_0$

For the 2-torus manifold, we can define the new operation  $\natural_0$  introduced from the (geometric) equivariant surgery. First, we explain this operation  $\natural_0$ .

The equivariant surgery  $\natural_0$  is the operation described in Figure 17. As we see in Figure 16, generally in the orbit cell decomposition of 2-torus manifolds, there is the multi-edge (i.e., two vertices connected by more than two edges, also see the left figures in Figure 17). The left figures in Figure 17 show the neighborhood around the multi-edge with coloring in the orbit cell decomposition  $(Q, \lambda)$  and  $(D^3, \rho)$ . First we take two facets  $D_1 \subset Q$  and  $D_2 \subset D^3$  as the facets surrounded by black edges in Figure 17. The equivariant surgery  $\natural_0$  is the gluing  $Q$  and  $D^3$  with small neighborhoods of  $D_1$  and  $D_2$  removed. As a result, we get the new orbit cell decomposition with coloring  $(Q', \lambda')$  as in the right figure in Figure 17, and vice versa.  $(Q', \lambda')$  is denoted by  $\natural_0(Q, \lambda)$ .






Figure 17 The equivariant surgery  $\mathfrak{h}_0$  (from left to right) and its inverse  $(\mathfrak{h}_0)^{-1}$  (from right to left)

Let  $D$  be a facet in the orbit cell decomposition  $Q$  whose boundary consists of 2 vertices and 2 edges, and  $\pi : M \rightarrow Q$  be an orbit projection. We can easily show that this facet  $D$  corresponds with the invariant submanifold diffeomorphic to  $S^2$  in  $M$ , that is,  $\pi^{-1}(D) = S^2$ . So we can understand the geometric meaning of the operation  $\mathfrak{h}_0$  as follows. Because  $\mathbb{Z}_2^3$  acts on the last three coordinates of  $S^3 \subset \mathbb{R} \oplus \mathbb{R}^3$ , there is the  $\mathbb{Z}_2^3$ -invariant submanifold  $S^2 \subset \mathbb{R}^3 \oplus \{0\}$  (remark that there are two fixed points in this submanifold). Then its invariant, closed, tubular neighborhood is equivariantly isomorphic to  $S^2 \times D^1$  with the standard  $\mathbb{Z}_2^3$ -action, where  $D^1 = I^1$ . We next consider  $S^3 = S^2 \times D^1 \cup D^3 \times S^0$ , and prepare the  $\mathbb{Z}_2^3$ -invariant part  $S^3 \setminus S^2 \times D^1 = D^3 \times S^0$  (remark that there is no fixed points in this disconnected manifold). Next we remove the invariant neighborhood  $S^2 \times D^1$  around  $S^2 = \pi^{-1}(D)$  from the 2-torus manifold  $M$ , i.e., we take  $M \setminus S^2 \times D^1$ . Finally we glue these two invariant manifold  $M \setminus S^2 \times D^1 \cup_\partial S^3 \setminus S^2 \times D^1$ . This operation is identically the equivariant surgery of 3-dimensional manifold which is different from the equivariant surgery explained in Subsection 3.4.

### 6.3 Remark on the operations $\sharp^\Delta$ and $\sharp^C$

Because we can easily regard the equivariant connected sum as the equivariant surgery, the three type different operations  $\sharp$ ,  $\mathfrak{h}$  and  $\mathfrak{h}_0$  in this paper are introduced by the (geometric) equivariant surgeries. Finally in this paper, we prove that  $\sharp^\Delta$  and  $\sharp^C$  can be constructed by these equivariant surgeries, i.e., we prove Theorem 1.2.

**Proof of Theorem 1.2** These relations (1) and (2) are shown by Figures 18 and 19, respectively.




Figure 18  $\sharp^\Delta = \mathfrak{h}_0 \circ \mathfrak{h} \circ \sharp$

In Figure 18, the figure on the top from left to right means the connected sum  $\sharp$  on two

black vertices. Next we can do  $\natural$  along the black edge, then we have the 2-gon facet as in the left bottom figure. Finally, we can do  $\natural_0$  for this 2-gon facet. As the result, we have the operation  $\sharp^\Delta = \natural_0 \circ \natural \circ \sharp$ .




Figure 19  $\sharp^C P^3(l) = \natural_0 \circ \natural^{l-2} \circ \sharp$

In Figure 19, we explain the  $l = 5$  case only. However, we can easily apply the same argument for all  $l \geq 3$ . In Figure 19, the first map means the connected sum  $\sharp$  on two black vertices. Then we have the  $(2l - 2)$ -gon facet (in Figure 19, this facet is the 8-gon). Next we can do  $\natural$  along the black edge as in the second figure, then, this facet becomes the  $(2l - 4)$ -gon facet. Iterating this argument  $((l - 2)$ -times), finally, we have the 2-gon facet. Then we can do  $\natural_0$  along this 2-gon facet. As the result, we have the operation  $\sharp^C P^3(l) = \natural_0 \circ \natural^{l-2} \circ \sharp$ .

**Acknowledgement** The author would like to thank Professors Zhi Lü and Taras Panov for invaluable discussions and advices. He also would like to thank Professors Zhi Lü and DongYoup Suh for the excellent circumstances to do research, and the referee for invaluable comments.

## References

- [1] Buchstaber, V. M. and Panov, T. E., Torus actions and their applications in topology and combinatorics, Univ. Lect. Ser., Vol. 24, A. M. S., Providence, RI, 2002.
- [2] Davis, M. and Januszkiewicz, T., Convex polytopes, coxeter orbifolds and torus action, *Duke Math. J.*, **62**(2), 1991, 417–451.
- [3] Izmitiev, I. V., Three-dimensional manifolds defined by coloring a simple polytope, *Math. Notes*, **69**, 2001, 340–346.
- [4] Kuroki, S., Remarks on McGavran’s paper and Nishimura’s results, Trends in Mathematics, Vol. 10, No. 1, Proceedings of Toric Topology Workshop KAIST 2008, Suh, D. Y., et al (eds.), ICMS in KAIST, 2008, 77–79.
- [5] Lü, Z., 2-torus manifolds, cobordism and small covers, *Pacific J. Math.*, **241**, 2009, 285–308.
- [6] Lü, Z. and Masuda, M., Equivariant classification of 2-torus manifolds, *Colloq. Math.*, **115**, 2009, 171–188.
- [7] Lü, Z. and Yu, L., Topological types of 3-dimensional small covers, *Forum Math.*, to appear. arXiv: 0710.4496
- [8] McGavran, D.,  $T^3$ -actions on simply connected 6-manifolds, I, *Trans. Amer. Math. Soc.*, **220**, 1976, 59–85.
- [9] Milnor, J. W. and Stasheff, J. D., Characteristic Classes, Princeton University Press, Princeton, 1974.
- [10] Nakayama, H. and Nishimura, Y., The orientability of small covers and coloring simple polytopes, *Osaka J. Math.*, **42**(1), 2005, 243–256.
- [11] Nishimura, Y., Equivariant surgeries of small covers (in Japanese), *RIMS Kokyuroku*, **1393**, 2004, 44–47.
- [12] Orlik, P. and Raymond, F., Actions of the torus on 4-manifolds, I, *Trans. Amer. Math. Soc.*, **152**, 1970, 531–559.
- [13] Ziegler, G., Lecture on Polytopes, Graduate Texts in Math., Vol. 152, Springer-Verlag, New York, 1995.