# The Cauchy Integral Operator on Weighted Hardy Space\*\*\*

Jianmiao RUAN\* Jiecheng CHEN\*\*

**Abstract** The authors show that the Cauchy integral operator is bounded from  $H^p_{\omega}(R^1)$  to  $h^p_{\omega}(R^1)$  (the weighted local Hardy space). To prove the results, a kind of generalized atoms is introduced and a variant of weighted "Tb theorem" is considered.

 Keywords Cauchy integral, Calderón-Zygmund operator, Weighted Hardy space, Weighted local Hardy space
 2000 MR Subject Classification 42B20, 42B25

# 1 Introduction

Let  $\mathbb{R}^n$  be the *n*-dimensional Euclidean space. The Cauchy integral operator is defined by

$$C_A f(x) = \text{p.v.} \int_{\mathbb{R}^1} \frac{1}{x - y + i(A(x) - A(y))} f(y) dy,$$
 (1.1)

where A(x) is a real valued function. This operator is very important in real and complex analysis, and has attracted many mathematicians to investigate it (see, for example, [2–4, 11]).

It is well-known that  $C_A$  is bounded on  $L^p(\mathbb{R}^1)$ , but few results are known on the Hardy space  $H^p(\mathbb{R}^1)$ . Recently, Komori [14] showed that  $C_A$  is bounded from  $H^p(\mathbb{R}^1)$  to  $h^p(\mathbb{R}^1)$  (the local Hardy space). In this paper, we consider the weighted version of Hardy space and show that  $C_A$  is bounded from  $H^p_w(\mathbb{R}^1)$  to  $h^p_w(\mathbb{R}^1)$ . To prove the theorem, we introduce a kind of generalized atoms and consider a variant of weighted "Tb theorem".

# 2 Definitions and Notations

Throughout this paper, we always use the letter C to denote positive constants that may vary at each occurrence, but is independent of the essential variables. And we assume that, unless otherwise stated, all given functions are complex valued.

We denote the Euclidean ball with center x of radius r by B(x, r), and the Lebesgue measure of a measurable set E by |E|.

Manuscript received February 11, 2009, Revised January 19, 2010. Published online June 21, 2010.

<sup>\*</sup>Department of Mathematics, Zhejiang University, Hangzhou 310027, China; Department of Mathematics, Zhejiang Education Institute, Hangzhou 310012, China. E-mail: rjmath@163.com

<sup>\*\*</sup>Corresponding author. Department of Mathematics, Zhejiang University, Hangzhou 310027, China. E-mail: jcchen@zju.edu.cn

<sup>\*\*\*</sup>Project supported by the National Natural Science Foundation of China (Nos. 10571156, 10871173, 10931001), the Zhejiang Provincial Natural Science Foundation of China (No. Y606117) and the Science Foundation of Education Department of Zhejiang Province (No. Y200803879).

Let  $\varphi$  be a fixed real valued Schwartz function in  $S(\mathbb{R}^n)$  such that  $\operatorname{supp}(\varphi) \subset B(0,1)$  and  $\int \varphi(x) dx = 1$ . We denote

$$f^{++}(x) = \sup_{t>0} |\varphi_t * f(x)|, \quad f^+(x) = \sup_{0 < t < 1} |\varphi_t * f(x)|,$$

where  $\varphi_t(x) = \frac{1}{t^n} \varphi(\frac{x}{t})$ .

**Definition 2.1** Let  $0 , the Hardy space <math>H^p(\mathbb{R}^n)$  (see [6]) and the local Hardy space  $h^p(\mathbb{R}^n)$  (see [9]) are defined respectively by

$$H^{p}(\mathbb{R}^{n}) = \{ f \in S'(\mathbb{R}^{n}) : \|f\|_{H^{p}} = \|f^{++}\|_{L^{p}} < \infty \}$$

and

$$h^{p}(R^{n}) = \{ f \in S'(R^{n}) : ||f||_{h^{p}} = ||f^{+}||_{L^{p}} < \infty \}$$

We remark that  $H^p(\mathbb{R}^n) \subset h^p(\mathbb{R}^n), h^1(\mathbb{R}^n) \subset L^1(\mathbb{R}^n)$  and all the inclusions are proper.

**Definition 2.2** For  $0 < \alpha \leq 1$ , the Lipschitz space  $\Lambda^{\alpha}(\mathbb{R}^n)$  and the local Lipschitz space  $\Lambda^{\alpha}_{loc}(\mathbb{R}^n)$  are the sets of all functions f satisfying the following conditions respectively

$$\|f\|_{\Lambda^{\alpha}} = \sup_{0 < |x-y|} \frac{|f(x) - f(y)|}{|x-y|} < \infty,$$
$$\|f\|_{\Lambda^{\alpha}_{\text{loc}}} = \sup_{0 < |x-y| < 2} \frac{|f(x) - f(y)|}{|x-y|} < \infty.$$

It is easy to see that  $\Lambda^1(\mathbb{R}^n) = \Lambda^1_{\text{loc}}(\mathbb{R}^n)$  and  $\Lambda^{\alpha}(\mathbb{R}^n) \subset \Lambda^{\alpha}_{\text{loc}}(\mathbb{R}^n)$   $(0 < \alpha < 1)$ , where the inclusion is proper. Furthermore, we know that the dual space of  $H^p(\mathbb{R}^n)$  is  $\Lambda^{n(\frac{1}{p}-1)}(\mathbb{R}^n)$ , i.e.  $(H^p(\mathbb{R}^n))^* = \Lambda^{n(\frac{1}{p}-1)}(\mathbb{R}^n)$ , where  $\frac{n}{n+1} (see [6]).$ 

**Definition 2.3** A locally integrable function f is in BMO if

$$\sup_{B} \frac{1}{|B|} \int_{B} |f - m_B f| \mathrm{d}x < \infty,$$

where  $m_B f = \frac{1}{|B|} \int_B f(x) dx$  and the supremum is taken over all balls B. We denote the supremum by  $\|f\|_{BMO}$ .

A weight is a nonnegative, locally integrable function. We consider weights satisfying the following conditions.

**Definition 2.4** Let  $1 < q < \infty$ . We say that a weight w satisfies the  $A_q$  condition if there exists a positive constant C such that for all balls B,

$$\left(\frac{1}{|B|}\int_B w(x)\mathrm{d}x\right)\left(\frac{1}{|B|}\int_B w(x)^{-\frac{1}{q-1}}\mathrm{d}x\right)^{q-1} \le C.$$

We also say that a weight w satisfies the  $A_1$  condition if there is a constant C > 0 such that for all balls B,

$$\frac{1}{|B|} \int_B w(x) \mathrm{d}x \le C \operatorname{essinf}_{x \in B} w(x).$$

Finally, we define  $A_{\infty} = \bigcup_{q \ge 1} A_q$ .

462

Observe that  $A_p \subset A_q$  if  $1 \le p < q$ .

**Definition 2.5** (see [17]) Let  $w \in A_{\infty}$  and  $0 . We define the weighted Hardy space <math>H^p_w(\mathbb{R}^n)$  and the weighted local Hardy space  $h^p_w(\mathbb{R}^n)$  as follows:

$$H_w^p(R^n) = \{ f \in S'(R^n) : \|f\|_{H_w^p} = \|f^{++}\|_{L_w^p} < \infty \}$$

and

$$h_w^p(R^n) = \{ f \in S'(R^n) : \|f\|_{h_w^p} = \|f^+\|_{L_w^p} < \infty \}$$

Next we define Calderón-Zygmund operators. One may refere to [10, 16]. But since we are interested in the Cauchy integral operator, our definitions will be presented as follows (see [14]).

**Definition 2.6** Let  $0 < \delta \leq 1$ . A locally integrable function K(x, y) defined on  $\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n : x \neq y\}$  is called a Calderón-Zygmund kernel if it satisfies the following conditions:

$$|K(x,y)| \le \frac{C}{|x-y|^n},$$
(2.1)

$$|K(x,y) - K(x,z)| + |K(y,x) - K(z,x)| \le C \frac{|y-z|^{\delta}}{|x-z|^{n+\delta}}, \quad 2|y-z| < |x-z|.$$
(2.2)

**Definition 2.7** We say that an operator T is a  $\delta$ -Calderón-Zygmund operator associated with a Calderón-Zygmund kernel K(x, y) if for every  $f \in L^2(\mathbb{R}^n)$ ,

$$Tf(x) = \lim_{\epsilon \to 0} \int_{|x-y| > \epsilon} K(x,y) f(y) dy$$

exists almost everywhere in  $\mathbb{R}^n$  and T is bounded on  $L^2(\mathbb{R}^n)$ , i.e.  $\|Tf\|_{L^2} \leq C \|f\|_{L^2}$ .

**Remark 2.1** If T is a  $\delta$ -Calderón-Zygmund operator and  $w \in A_q$ , then T is bounded on  $L^q_w(\mathbb{R}^n), q > 1$  (see [7], [11, p. 52] and [15]).

**Definition 2.8** The transpose of an operator T is denoted by

$${}^{\mathrm{t}}Tf(x) = \lim_{\epsilon \to 0} \int_{|x-y| > \epsilon} K(y,x)f(y) \mathrm{d}y.$$

**Definition 2.9** For a bounded function b, we define

$$\widetilde{^{\mathbf{t}}T}b(x) = \lim_{\epsilon \to 0} \int_{|x-y| > \epsilon} \{ K(y,x) - K(y,0)\chi_{|y| \ge 1}(y) \} b(y) \mathrm{d}y.$$

Note that if  $b \in L^2(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ , then  $\widetilde{tT}b(x) = tTb(x) + C_b$  a.e., where  $C_b$  is a constant.

**Definition 2.10** Let  $\beta > 0$ . A bounded function b is said to be  $\beta$ -accretive if  $\operatorname{Re} b(x) \ge \beta$  for almost all x.

#### 3 Theorems

First we recall some known results. The  $L^p$  boundedness of  $C_A$  is well-known, and the following theorem is the most essential (see [3] and [10, p. 647]).

**Theorem 3.1** If  $A' \in L^{\infty}(\mathbb{R}^1)$ , then the Cauchy integral operator  $C_A$  defined by (1.1) is a 1-Calderón-Zygmund operator.

Recently, Komori [14] showed that  $C_A$  is bounded from  $H^p(\mathbb{R}^1)$  to  $h^p(\mathbb{R}^1)$ , i.e., the following theorem.

**Theorem 3.2** Let  $0 < \alpha < 1$  and  $\frac{1}{1+\alpha} \le p \le 1$ . If  $A' \in L^{\infty}(\mathbb{R}^1) \cap \wedge^{\alpha}(\mathbb{R}^1)$ , then  $C_A$  is bounded from  $H^p(\mathbb{R}^1)$  to  $h^p(\mathbb{R}^1)$ .

In order to prove Theorem 3.2, he considered a variant of "Tb theorem" as follows.

**Theorem 3.3** Let  $0 < \alpha < 1$ ,  $\frac{n}{n+\delta} , <math>\frac{n}{n+\alpha} \leq p$  and T be a  $\delta$ -Calderón-Zygmund operator. If there exists a  $\beta$ -accretive function b such that  $b, \widetilde{tT}b \in \wedge^{\alpha}(\mathbb{R}^n)$ , then T is a bounded operator from  $H^p(\mathbb{R}^n)$  to  $h^p(\mathbb{R}^n)$  and

$$||Tf||_{h^p} \leq C ||f||_{H^p}.$$

As a corollary, he got Theorem 3.2 by Theorem 3.3.

**Remark 3.1** In [14], the author obtained the  $h^p(\mathbb{R}^n)$   $(0 estimate by <math>(H^p(\mathbb{R}^n))^* = \Lambda^{n(\frac{1}{p}-1)}(\mathbb{R}^n)$  and the  $h^1(\mathbb{R}^n)$  estimate by interpolation. In this paper, we use a different method to prove our theorems. The details are presented in Section 5.

Now we turn to the weighted Hardy space case. Our main results are the following.

**Theorem 3.4** Let  $0 < \alpha \leq 1 \leq q$ ,  $\frac{nq}{n+\delta} and <math>\frac{nq}{n+\alpha} \leq p < q$ . Assume that  $w \in A_q$ and T is a  $\delta$ -Calderón-Zygmund operator. If there exists a  $\beta$ -accretive function b such that b,  $\widetilde{^{t}Tb} \in \wedge^{\alpha}_{loc}(\mathbb{R}^n)$ , then T is a bounded operator from  $H^p_w(\mathbb{R}^n)$  to  $h^p_w(\mathbb{R}^n)$  and

$$||Tf||_{h_w^p} \le C ||f||_{H_w^p}$$

At the end point p = 1, if we strengthen the weight condition, we have

**Theorem 3.5** If  $w \in A_1$  and T is a  $\delta$ -Calderón-Zygmund operator, then T is a bounded operator from  $H^1_w(\mathbb{R}^n)$  to  $h^1_w(\mathbb{R}^n)$  and

$$||Tf||_{h^1_w} \leq C ||f||_{H^1_w}.$$

**Remark 3.2** For a  $\delta$ -Calderón-Zygmund operator T and  $w \in A_1$ , it is easy to check that  $\|Tf\|_{L^1_w} \leq C \|f\|_{H^1_w}$  by some standard argument. Furthermore, Quek and Yang [15] obtained that  $\|Tf\|_{H^1_w} \leq C \|f\|_{H^1_w}$  if  $\widetilde{tT}1 = C$  and Komori [13] obtained that  $\|Tf\|_{h^1_w} \leq C \|f\|_{H^1_w}$  if  $\widetilde{tT}1 \in \wedge^{\alpha}(\mathbb{R}^n)$ . Since  $H^1_w(\mathbb{R}^n) \subset h^1_w(\mathbb{R}^n) \subset L^1_w(\mathbb{R}^n)$ , Theorem 3.5 extends these results. Especially, taking w = 1, our results are also new.

As a corollary of these theorems, we obtain the boundedness of the Cauchy integral operator.

**Theorem 3.6** Let  $0 < \alpha \leq 1 \leq q$ ,  $\frac{q}{2} and <math>\frac{q}{1+\alpha} \leq p < q$ . If  $w \in A_q$  and  $A' \in L^{\infty}(\mathbb{R}^1) \cap \wedge_{\text{loc}}^{\alpha}(\mathbb{R}^1)$ , then  $C_A$  is bounded from  $H^p_w(\mathbb{R}^1)$  to  $h^p_w(\mathbb{R}^1)$ .

**Theorem 3.7** If  $A' \in L^{\infty}(\mathbb{R}^1)$  and  $w \in A_1$ , then  $C_A$  is bounded from  $H^1_w(\mathbb{R}^1)$  to  $h^1_w(\mathbb{R}^1)$ .

### 4 Some Lemmas

First we present two elementary lemmas on weight functions without proof (see [8] or [19, p. 226]).

**Lemma 4.1** If  $w \in A_q$ ,  $q \ge 1$ , then there exists a positive constant C such that

$$\frac{w(B(x_0, r))}{w(B(x_0, s))} \le C\Big(\frac{|B(x_0, r)|}{|B(x_0, s)|}\Big)^{\frac{1}{2}}$$

for all r > s and  $x_0 \in \mathbb{R}^n$ . Especially

$$w(B(x_0, 2^j r)) \le C 2^{nqj} w(B(x_0, r)).$$

**Lemma 4.2** Let f be a nonnegative locally integrable function. If  $w \in A_q$ ,  $q \ge 1$ , then

$$\frac{1}{|B(x_0,r)|} \int_{B(x_0,r)} f(x) \mathrm{d}x \le C \Big( \frac{1}{w(B(x_0,r))} \int_{B(x_0,r)} f(x)^q w(x) \mathrm{d}x \Big)^{\frac{1}{q}}.$$

Next we define atoms and molecules in  $H^p_w(\mathbb{R}^n)$  and  $h^p_w(\mathbb{R}^n)$  and obtain some properties of Hardy spaces.

**Definition 4.1** Let  $1 \le q \le \infty$ . A function a(x) is called an  $(H_w^p, q)$ -atom centered at  $x_0$  if there exists a ball  $B(x_0, r)$  such that the following conditions are satisfied

$$\operatorname{supp}(a) \subset B(x_0, r),\tag{4.1}$$

$$\|a\|_{L^q_w} \le w(B(x_0, r))^{\frac{1}{q} - \frac{1}{p}},\tag{4.2}$$

$$\int_{R^n} a(x) \mathrm{d}x = 0. \tag{4.3}$$

**Lemma 4.3** (see [7] and [17, p. 111]) Let  $\frac{nq}{n+1} and <math>p < q$ . If  $w \in A_q$  and a(x) is an  $(H_w^p, q)$ -atom, then  $a \in H_w^p(\mathbb{R}^n)$  and

 $||a(x)||_{H^p_w} \le C.$ 

Furthermore, we have the atomic decomposition of  $H^p_w(\mathbb{R}^n)$  (see [7, 17]) as follows.

**Lemma 4.4** Let  $\frac{nq}{n+1} and <math>p < q$ . If  $w \in A_q$  and  $f \in H^p_w(\mathbb{R}^n)$ , then f can be written as

$$f = \sum_{j=1}^{\infty} \lambda_j a_j,$$

where  $a_j$ 's are  $(H^p_w, q)$ -atoms and

$$\sum_{j=1}^{\infty} |\lambda_j|^p \approx \|f\|_{H^p_w}^p$$

**Definition 4.2** Let  $1 \le q \le \infty$ . A function a(x) is called a large  $(h_w^p, q)$ -atom centered at  $x_0$  if there exists a ball  $B(x_0, r)$  of radius  $r \ge 1$  such that conditions (4.1) and (4.2) are satisfied.

**Lemma 4.5** (see [13]) Let  $\frac{n}{n+1} and <math>p < q$ . If  $w \in A_q$  and a(x) is a large  $(h_w^p, q)$ -atom, then  $a \in h_w^p(\mathbb{R}^n)$  and

$$\|a\|_{h^p_w} \le C.$$

**Definition 4.3** Let  $1 \le q \le \infty$ . A function a(x) is a small  $(h_w^p, q)$ -atom centered at  $x_0$  if there exists a ball  $B(x_0, r)$  of radius r < 1, which satisfies conditions (4.1), (4.2) and

$$\left|\int_{\mathbb{R}^n} a(x) \mathrm{d}x\right| \le r^{n\frac{q-1}{p}} \left(\frac{|B(x_0, r)|}{w(B(x_0, r))}\right)^{\frac{1}{p}}$$

**Lemma 4.6** (see [13]) Let  $\frac{nq}{n+1} and <math>p < q$ . If  $w \in A_q$  and a(x) is a small  $(h_w^p, q)$ -atom, then  $a \in h_w^p(\mathbb{R}^n)$  and

$$\|a\|_{h^p_m} \le C.$$

**Definition 4.4** Let b be  $\beta$ -accretive and  $1 \le q \le \infty$ . A function a(x) is a small  $(h_w^p, q, b)$ atom centered at  $x_0$  if there exists a ball  $B(x_0, r)$  of radius r < 1, which satisfies conditions (4.1), (4.2) and

$$\left| \int_{R^{n}} a(x)b(x) \mathrm{d}x \right| \le r^{n\frac{q-1}{p}} \left( \frac{|B(x_{0}, r)|}{w(B(x_{0}, r))} \right)^{\frac{1}{p}}.$$
(4.4)

**Lemma 4.7** Let b be  $\beta$ -accretive and  $b \in \wedge_{\text{loc}}^{\alpha}(\mathbb{R}^n)$   $(0 < \alpha \leq 1)$ . Assume that  $\frac{nq}{n+1} and <math>\frac{nq}{n+\alpha} \leq p < q$ . If  $w \in A_q$  and a(x) is a small  $(h_w^p, q, b)$ -atom, then  $a \in h_w^p(\mathbb{R}^n)$  and

 $||a||_{h^p_w} \le C.$ 

**Proof** According to Lemma 4.6, we only need to show that a is a small  $(h_w^p, q)$ -atom.

$$\begin{split} \left| \int_{B(x_0,r)} a(x) \mathrm{d}x \right| &\leq \left| \frac{1}{b(x_0)} \int_{B(x_0,r)} a(x)(b(x) - b(x_0)) \mathrm{d}x \right| + \left| \frac{1}{b(x_0)} \int_{B(x_0,r)} a(x)b(x) \mathrm{d}x \right| \\ &\leq C \frac{r^{\alpha}}{\beta} \Big( \int_{B(x_0,r)} |a(x)|^q w(x) \mathrm{d}x \Big)^{\frac{1}{q}} \frac{|B(x_0,r)|}{w(B(x_0,r))^{\frac{1}{q}}} + \frac{1}{\beta} r^{n\frac{q-1}{p}} \Big( \frac{|B(x_0,r)|}{w(B(x_0,r))} \Big)^{\frac{1}{p}} \\ &\leq C r^{\alpha+n-\frac{n}{p}} \Big( \frac{|B(x_0,r)|}{w(B(x_0,r))} \Big)^{\frac{1}{p}} + \frac{1}{\beta} r^{n\frac{q-1}{p}} \Big( \frac{|B(x_0,r)|}{w(B(x_0,r))} \Big)^{\frac{1}{p}} \\ &\leq C r^{n\frac{q-1}{p}} \Big( \frac{|B(x_0,r)|}{w(B(x_0,r))} \Big)^{\frac{1}{p}}. \end{split}$$

Note that we have used the fact r < 1 and  $\frac{nq}{n+\alpha} \leq p$  in the last inequality.

**Definition 4.5** Let  $w \in A_q$ . A function M(x) is called a large  $(h_w^p, q, \delta)$ -molecule centered at  $x_0$  if there exists  $r \ge 1$  such that the following conditions are satisfied:

$$\left(\int_{|x-x_0|\leq 2r} |M(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \leq Cw(B(x_0,r))^{\frac{1}{q}-\frac{1}{p}},\tag{4.5}$$

$$|M(x)| \le \frac{r^{n+\delta} w(B(x_0, r))^{-\frac{1}{p}}}{|x - x_0|^{n+\delta}}, \quad |x - x_0| \ge 2r.$$
(4.6)

**Definition 4.6** Let  $w \in A_q$  and b be  $\beta$ -accretive. A function M(x) is called a small  $(h_w^p, q, \delta, b)$ -molecule centered at  $x_0$  if there exists r < 1 such that (4.5), (4.6) and the following condition

$$\left| \int_{\mathbb{R}^{n}} M(x)b(x) \mathrm{d}x \right| \le Cr^{n\frac{q-1}{p}} \left( \frac{|B(x_{0}, r)|}{w(B(x_{0}, r))} \right)^{\frac{1}{p}}.$$
(4.7)

466

**Lemma 4.8** Let  $\frac{nq}{n+\delta} and <math>p < q$ . If  $w \in A_q$  and a function M(x) is a large  $(h_w^p, q, \delta)$ -molecule, then  $M \in h_w^p(\mathbb{R}^n)$  and

$$\|M\|_{h^p_w} \le C.$$

**Lemma 4.9** Let  $0 < \alpha \leq 1 \leq q$ ,  $\frac{nq}{n+\delta} and <math>\frac{nq}{n+\alpha} \leq p \leq 1$ . Suppose that b is  $\beta$ -accretive and  $b \in \wedge_{\text{loc}}^{\alpha}(\mathbb{R}^n)$ . If a function M(x) is a small  $(h_w^p, q, \delta, b)$ -molecule, then  $M \in h_w^p(\mathbb{R}^n)$  and

$$\|M\|_{h^p_w} \le C.$$

Lemmas 4.8 and 4.9 are the key lemmas to prove our theorems. The proofs of the two lemmas are similar in nature. So we shall only prove Lemma 4.9 below. The idea of our proof comes from [12–14, 18].

**Proof of Lemma 4.9** Let  $E_0 = \{x : |x - x_0| < 2r\}$  and  $E_i = \{x : 2^i r \le |x - x_0| < 2^{i+1}r\},$  $i = 1, 2, 3, \dots, b(E_i) = \int_{E_i} b(x) dx$ . Since  $b(E_i) \ne 0$ , we denote  $\chi_i = \chi_{E_i}(x), \ \widetilde{\chi}_i = \frac{\chi_i}{b(E_i)},$  $m_i = \frac{1}{b(E_i)} \int_{E_i} b(x) M(x) dx$  and  $\widetilde{m}_i = \int_{E_i} b(x) M(x) dx$ .

We write

$$M(x) = \sum_{i=0}^{\infty} (M(x) - m_i)\chi_i(x) + \sum_{i=0}^{\infty} m_i\chi_i(x) = \sum_{i=0}^{\infty} M_i(x) + \sum_{i=0}^{\infty} \tilde{m}_i\tilde{\chi}_i(x),$$

where  $M_i(x) = (M(x) - m_i)\chi_i(x)$ . Letting  $N_j = \sum_{i=j}^{\infty} \widetilde{m}_i$ , we have

$$M(x) = \sum_{i=0}^{\infty} M_i(x) + \sum_{i=1}^{\infty} N_i(\widetilde{\chi}_i(x) - \widetilde{\chi}_{i-1}(x)) + N_0\widetilde{\chi}_0(x)$$
  
= I + II + III.

Next we shall estimate the above three terms.

(a) It is clear that  $\operatorname{supp}(M_i) \subset B(x_0, 2^{i+1}r)$  and  $\int M_i(x)b(x)dx = 0$ . So

$$\left(\int |M_0(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \le \left(\int_{E_0} |M(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} + |m_0| \left|\int_{E_0} w(x) \mathrm{d}x\right|^{\frac{1}{q}} \\ \le Cw(B(x_0, r))^{\frac{1}{q} - \frac{1}{p}} + |m_0|w(E_0)^{\frac{1}{q}}.$$

By the definition of  $m_0$  and Lemma 4.2, we have

$$|m_{0}| = \frac{1}{|b(E_{0})|} \left| \int_{E_{0}} M(y)b(y)dy \right|$$
  

$$\leq C \frac{||b||_{L^{\infty}}}{\beta|E_{0}|} \int_{E_{0}} |M(y)|dy$$
  

$$\leq C \left(\frac{1}{w(E_{0})} \int_{E_{0}} |M(y)|^{q}w(y)dy\right)^{\frac{1}{q}}$$
  

$$\leq Cw(B(x_{0},r))^{\frac{1}{q}-\frac{1}{p}}w(E_{0})^{-\frac{1}{q}}.$$

Therefore, we get

$$\left(\int |M_0(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \le Cw(B(x_0,r))^{\frac{1}{q}-\frac{1}{p}}.$$

So by Lemma 4.5 or 4.7 we have

$$\|M_0\|_{h^p_w} \le C.$$

When  $i \geq 1$ , we have

$$\left(\int |M_i(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \le \left(\int_{E_i} |M(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} + |m_i| w(E_i)^{\frac{1}{q}}$$
$$= \widetilde{\mathbf{I}} + \widetilde{\mathbf{I}}.$$

By condition (4.6), we have

$$\widetilde{\mathbf{I}} \leq r^{n+\delta} w(B(x_0, r))^{-\frac{1}{p}} \Big( \int_{E_i} \frac{w(x)}{|x - x_0|^{q(n+\delta)}} \mathrm{d}x \Big)^{\frac{1}{q}} \\ \leq \frac{r^{n+\delta}}{(2^i r)^{n+\delta}} w(B(x_0, r))^{-\frac{1}{p}} w(B(x_0, 2^{i+1} r))^{\frac{1}{q}} \\ \leq C 2^{-i\delta} w(B(x_0, r))^{\frac{1}{q} - \frac{1}{p}}.$$

The last inequality was obtained by Lemma 4.1.

Using the condition (4.6) again and the fact  $|E_i| \approx |B(x_0, 2^{i+1}r)|$ , we have

$$\begin{split} \widetilde{\Pi} &\leq w(E_i)^{\frac{1}{q}} \frac{\|b\|_{L^{\infty}}}{\beta |E_i|} \int_{E_i} |M(y)| \mathrm{d}y \\ &\leq C \frac{w(E_i)^{\frac{1}{q}}}{|E_i|} \int_{E_i} \frac{r^{n+\delta} w(B(x_0,r))^{-\frac{1}{p}}}{|y-x_0|^{n+\delta}} \mathrm{d}y \\ &\leq C 2^{-i\delta} w(B(x_0,r))^{\frac{1}{q}-\frac{1}{p}}. \end{split}$$

According to Lemma 4.1, we reach

$$\left(\int |M_i|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \le C 2^{-i\delta} w(B(x_0, r))^{\frac{1}{q} - \frac{1}{p}}$$
$$\le C 2^{-i(\delta + nq(\frac{1}{q} - \frac{1}{p}))} w(B(x_0, 2^{i+1}r))^{\frac{1}{q} - \frac{1}{p}}.$$

By Lemma 4.5 or 4.7, we get

$$||M_i||_{h^p_m} \le C2^{-i(\delta + nq(\frac{1}{q} - \frac{1}{p}))}.$$

Since  $\frac{nq}{n+\delta} < p$ , we finally conclude

$$\sum_{i=0}^{\infty} \|M_i\|_{h^p_w}^p \le C \quad \text{and} \quad \|\mathbf{I}\|_{h^p_w} \le C.$$

(b) Let  $A_i = N_i(\tilde{\chi}_i(x) - \tilde{\chi}_{i-1}(x))$ . It is easy to see that  $\operatorname{supp}(A_i) \subset B(x_0, 2^{i+1}r)$  and  $\int_{\mathbb{R}^n} A_i(x)b(x)dx = 0, i = 1, 2, 3 \cdots$ .

Using condition (4.6) and Lemma 4.1, we have

$$\begin{split} \left( \int |A_{i}(x)|^{q} w(x) \mathrm{d}x \right)^{\frac{1}{q}} &\leq |N_{i}| \left( \int |\tilde{\chi}_{i}(x) - \tilde{\chi}_{i-1}(x)|^{q} w(x) \mathrm{d}x \right)^{\frac{1}{q}} \\ &\leq \left( \frac{|w(E_{i})|^{\frac{1}{q}}}{|b(E_{i})|} + \frac{|w(E_{i-1})|^{\frac{1}{q}}}{|b(E_{i-1})|} \right) \sum_{k=i}^{\infty} \left| \int_{E_{k}} M(y) b(y) \mathrm{d}y \right| \\ &\leq C \frac{w(B(x_{0}, 2^{i+1}r))^{\frac{1}{q}}}{\beta(2^{i}r)^{n}} \|b\|_{L^{\infty}} \sum_{k=i}^{\infty} \int_{E_{k}} |M(y)| \mathrm{d}y \\ &\leq C \frac{w(B(x_{0}, r))^{\frac{1}{q}}}{r^{n}} \sum_{k=i}^{\infty} \int_{E_{k}} \frac{r^{n+\delta}w(B(x_{0}, r))^{-\frac{1}{p}}}{|y-x_{0}|^{n+\delta}} \mathrm{d}y \\ &\leq Cr^{\delta}w(B(x_{0}, r))^{\frac{1}{q} - \frac{1}{p}} \int_{|y-x_{0}| \geq 2^{i}r} \frac{1}{|y-x_{0}|^{n+\delta}} \mathrm{d}y \\ &\leq C2^{-i\delta}w(B(x_{0}, r))^{\frac{1}{q} - \frac{1}{p}} \\ &\leq C2^{-i(\delta+nq(\frac{1}{q} - \frac{1}{p}))}w(B(x_{0}, 2^{i+1}r))^{\frac{1}{q} - \frac{1}{p}}. \end{split}$$

So by Lemma 4.5 or 4.7, we have

$$\sum_{i=0}^{\infty} \|A_i\|_{h^p_w}^p \le C \text{ and } \|\Pi\|_{h^p_w} \le C.$$

(c) Conditions (4.5), (4.6) together with the fact  $\operatorname{supp}(N_0 \widetilde{\chi}_0(x)) \subset B(x_0, 2r)$  imply

$$\left(\int |N_0 \tilde{\chi}_0(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \le |N_0| \frac{1}{|b(E_0)|} \left(\int_{E_0} w(x) \mathrm{d}x\right)^{\frac{1}{q}}$$
$$\le C \frac{w(E_0)^{\frac{1}{q}}}{\beta r^n} \left|\int M(x) b(x) \mathrm{d}x\right|$$
$$\le C \frac{w(E_0)^{\frac{1}{q}}}{r^n} \|b\|_{L^{\infty}} \left(\int_{|x-x_0|<2r} |M(x)| \mathrm{d}x + \int_{|x-x_0|\ge 2r} |M(x)| \mathrm{d}x\right)$$
$$\le C w(B(x_0, 2r))^{\frac{1}{q} - \frac{1}{p}}.$$

By condition (4.7), we have

$$\left|\int_{R^n} N_0 \widetilde{\chi}_0(x) b(x) \mathrm{d}x\right| = \left|\int_{R^n} M(x) b(x) \mathrm{d}x\right| \le Cr^{n\frac{q-1}{p}} \left(\frac{|B(x_0, r)|}{w(B(x_0, r))}\right)^{\frac{1}{p}}$$

By Lemma 4.5 or 4.7, we have

$$\|N_0\widetilde{\chi}_0\|_{h^p_w} \le C.$$

Finally combining (a), (b) and (c), we complete the proof of Lemma 4.9.

**Lemma 4.10** Let  $w \in A_1$  and M(x) be a function on  $\mathbb{R}^n$ . If there exists a ball  $B(x_0, r)$ , r > 0, satisfies the following conditions

$$\int_{|x-x_0| \le 2r} |M(x)| w(x) \mathrm{d}x \le C,\tag{4.8}$$

$$|M(x)| \le C \frac{r^{n+\delta} w(B(x_0, r))^{-1}}{|x - x_0|^{n+\delta}}, \quad |x - x_0| \ge 2r;$$
(4.9)

J. M. Ruan and J. C. Chen

and furthermore, if 0 < r < 1, M(x) also satisfies

$$\left| \int_{\mathbb{R}^n} M(x) \mathrm{d}x \right| \le C \frac{|B(x_0, r)|}{w(B(x_0, r))},$$
(4.10)

then  $M(x) \in h^1_w(\mathbb{R}^n)$  and

 $\|M(x)\|_{h^1_w} \le C.$ 

The proof of Lemma 4.10 is almost similar to that of Lemma 7 in [13], so we omit the details.

## 5 Proof of the Theorems

**Proof of Theorem 3.4** By Lemma 4.4 and Theorem 7.2 in [1], it suffices to show that there is a constant C > 0 such that  $||Ta||_{h_w^p} \leq C$  for every  $(H_w^p, 2q)$ -atom a(x).

Assuming  $(H_w^p, \infty)$ -atom a(x) supported in  $B(x_0, r)$ , we show that Ta(x) is a constant multiple of a large  $(h_w^p, q, \delta)$ -molecule with  $r \ge 1$  or a constant multiple of a small  $(h_w^p, q, \delta, b)$ -molecule with r < 1.

Since T is bounded on  $L^{2q}_w(\mathbb{R}^n)$  (see [11, p. 52]), we have

$$\left(\int_{|x-x_0|\leq 2r} |Ta(x)|^q w(x) \mathrm{d}x\right)^{\frac{1}{q}} \leq C \left(\int_{|x-x_0|\leq 2r} |Ta(x)|^{2q} w(x) \mathrm{d}x\right)^{\frac{1}{2q}} \left(\int_{|x-x_0|\leq 2r} w(x) \mathrm{d}x\right)^{\frac{1}{2q}}$$
$$\leq C ||a||_{L^{2q}_w} w(B(x_0,r))^{\frac{1}{2q}}$$
$$\leq C w(B(x_0,r))^{\frac{1}{q}-\frac{1}{p}}.$$

If  $|x - x_0| \ge 2r$ , then

$$|Ta(x)| = \left| \int_{B(x_0,r)} [K(x,y) - K(x,x_0)]a(y)dy \right|$$
$$\leq C \frac{r^{\delta}}{|x - x_0|^{n+\delta}} \int_{B(x_0,r)} |a(y)|dy$$
$$\leq C \frac{r^{n+\delta}}{|x - x_0|^{n+\delta}} w(B(x_0,r))^{-\frac{1}{p}}.$$

If  $r \ge 1$ , by Lemma 4.8, we have

$$|Ta(x)||_{h^p_w} \le C.$$

If r < 1, by (4.2) and (4.3), we have

$$\left| \int Ta(x)b(x)dx \right| = \left| \langle a, \widetilde{{}^{t}Tb} \rangle \right| = \left| \int_{B(x_{0},r)} a(x)[\widetilde{{}^{t}Tb}(x) - \widetilde{{}^{t}Tb}(x_{0})]dx \right|$$
$$\leq \|\widetilde{{}^{t}Tb}\|_{\Lambda^{\alpha}_{loc}} r^{\alpha} \int_{B(x_{0},r)} |a(x)|dx$$
$$\leq Cr^{\alpha}w(B(x_{0},r))^{-\frac{1}{p}}|B(x_{0},r)|$$
$$\leq Cr^{\alpha+n\frac{p-1}{p}} \left(\frac{|B(x_{0},r)|}{w(B(x_{0},r))}\right)^{\frac{1}{p}}$$
$$\leq Cr^{n\frac{q-1}{p}} \left(\frac{|B(x_{0},r)|}{w(B(x_{0},r))}\right)^{\frac{1}{p}},$$

470

where the last inequality is obtained for r < 1 and  $\frac{nq}{n+\alpha} \leq p$ .

By Lemma 4.9 and the above argument, we obtain the desired result

$$||Ta||_{h^p_w} \le C.$$

**Proof of Theorem 3.5** Similar to the argument of Theorem 3.4, it suffices to show that there is a constant C > 0 such that  $||Ta||_{h_w^1} \leq C$  for all  $(H_w^1, 2)$ -atoms a(x) supported in  $B(x_0, r)$ . It is easy to see

$$\int_{|x-x_0| \le 2r} |Ta(x)| w(x) \mathrm{d}x \le C.$$

And if  $|x - x_0| \ge 2r$ ,

$$Ta(x)| \le C \frac{r^{n+\delta} w(B(x_0,r))^{-1}}{|x-x_0|^{n+\delta}}.$$

If T is a  $\delta$ -Calderón-Zygmund operator, then  $\widetilde{tT}1 \in BMO$  and  $\|\widetilde{tT}1\|_{BMO} \leq C$  by the famous "T1 theorem" (see [5]). So we have

$$\left| \int_{\mathbb{R}^n} Ta(x) \mathrm{d}x \right| = \left| \langle a, \widetilde{^{\mathrm{t}}T1} \rangle \right| = \left| \int_{B(x_0, r)} a(x) [\widetilde{^{\mathrm{t}}T1}(x) - m_B \widetilde{^{\mathrm{t}T1}}] \mathrm{d}x \right|$$
$$\leq \|\widetilde{^{\mathrm{t}}T1}\|_{\mathrm{BMO}} |B(x_0, r)| w(B(x_0, r))^{-1}$$
$$\leq C \frac{|B(x_0, r)|}{w(B(x_0, r))},$$

where  $m_B f = \frac{1}{|B(x_0,r)|} \int_{B(x_0,r)} f(x) dx$ . According to Lemma 4.10 and the above argument, we get the desired conclusion immediately.

**Proof of Theorem 3.6** Note that  $C_A$  is a 1-Calderón-Zygmund operator by Theorem 3.1. Let b(x) = 1 + iA'(x). Then b is a 1-accretive and  $b \in \wedge^{\alpha}_{loc}(R^1)$ . By the calculus of complex analysis (refer to calculation in [14] or [19, p. 407]),

$$\begin{split} \widetilde{t}_{C_A} b(x) &= \lim_{\epsilon \to 0} \int_{|x-y| > \epsilon} \Big\{ \frac{1 + iA'(y)}{y - x + i(A(y) - A(x))} - \frac{1 + iA'(y)}{y + i(A(y) - A(0))} \chi_{|y| \ge 1}(y) \Big\} dy \\ &= \text{constant}, \end{split}$$

which implies  $\widetilde{{}^{t}C_{A}}b(x) \in \wedge_{\text{loc}}^{\alpha}(\mathbb{R}^{1})$ . Therefore the theorem is proved by Theorem 3.4.

Acknowledgements The authors would like to thank the referee for his/her several valuable suggestions. The first author also would like to thank Professor Komori for sending his paper [12] to him.

## References

- Bownik, M., Li, B., Yang, D. and Zhou, Y., Weighted anisotropic Hardy spaces and their applications in boundedness of sublinear operators, *Indiana Univ. Math. J.*, 57, 2008, 3065–3100.
- [2] Calderón, A. P., Commutators singular integrals on Lipschitz curves and applications, Proc. Inter. Congress of Math. Helsinki, 1978, 85–96.
- [3] Coifman, C., McIntosh, A. and Meyer, Y., L'integrale de Cauchy définit un opérateur borné sur L<sup>2</sup> pour les courbes Lipschitziennes, Ann. of Math., 116, 1982, 361–388.

- [4] David, G., Wavelets and Singular Integrals on Curves, Lecture Notes in Math., 1465, Springer-Verlag, Berlin, 1991.
- [5] David, G. and Journé, J. L., A boundedness criterion for generalized Calderón-Zygmund operators, Ann. of Math., 120, 1984, 371–397.
- [6] Fefferman, C. and Stein, E. M., H<sup>p</sup> spaces of several variables, Acta. Math., 129, 1972, 137–193.
- [7] García-Cuerva, J. and Kazarian, K. S., Weighted Norm Inequalities and Related Topics, North Holland, Amsterdam, 1985.
- [8] García-Cuerva, J. and Rubio de Francia, J., Calderón-Zygmund operators and unconditional bases of weighted hardy spaces, *Stud. Math.*, 109, 1994, 255–276.
- [9] Goldberg, D., A local version of real Hardy spaces, Duck Math. J., 46, 1979, 27–42.
- [10] Grafakos, L., Classical and Modern Fourier Analysis, China Machine Press, Beijing, 2006.
- [11] Journé, J. L., Calderón-Zygmund-Type Operators, Pseudo-differential Operators and the Cauchy Integral of Calderón, Lecture Notes in Math., 994, Springer-Verlag, New York, 1983.
- [12] Komori, Y., Calderón-Zygmund operators on  $H^p(\mathbb{R}^n)$ , Sci. Math. Japonicae, 53, 2001, 65–73.
- [13] Komori, Y., Calderón-Zygmund operator on weighted H<sup>p</sup>(R<sup>n</sup>), Hokk. Math. J., **32**, 2003, 673–684.
- [14] Komori, Y., The Cauchy integral on Hardy space, Hokk. Math. J., 37, 2008, 389–398.
- [15] Quek, T. and Yang, D., Calderón-Zygmund-type operators on weighted weak Hardy spaces over R<sup>n</sup>, Acta Math. Sinica (Engl. Ser.), 16, 2000, 141–160.
- [16] Stein, E. M., Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals, Princeton University Press, Princeton, 1993.
- [17] Strömberg, J. and Torchinsky, A., Weighted Hardy Spaces, Lecture Notes in Math., 1381, Springer-Verlag, New York, 1989.
- [18] Taibleson, M. H. and Weiss, G., The molecular characterization of certain Hardy spaces, Astérisgne, 77, 1980, 67–149.
- [19] Torchinsky, A., Real Variable Method in Harmonic Analysis, Academic Press, New York, 1986.