

Exact Controllability for the Fourth Order Schrödinger Equation*

Chuang ZHENG¹ Zhongcheng ZHOU²

Abstract The boundary controllability of the fourth order Schrödinger equation in a bounded domain is studied. By means of an L^2 -Neumann boundary control, the authors prove that the solution is exactly controllable in $H^{-2}(\Omega)$ for an arbitrarily small time. The method of proof combines both the HUM (Hilbert Uniqueness Method) and multiplier techniques.

Keywords Fourth order Schrödinger equation, HUM method, Controllability, Multiplier

2000 MR Subject Classification 93B05, 35Q40, 93C20

1 Introduction

Let Ω be a nonempty open bounded domain in \mathbb{R}^n ($n \in \mathbb{N}$) with C^3 boundary Γ , Γ_0 be a nonempty open subset of Γ , and $T > 0$ be a given time duration. Fix some $x_0 \in \mathbb{R}^n$, and put

$$\Gamma_0 \triangleq \{x \in \Gamma \mid (x - x_0) \cdot \nu(x) > 0\}, \quad (1.1)$$

where $\nu(x)$ is the unit outward normal vector of Ω at $x \in \Gamma$. We consider the following controlled fourth order linear Schrödinger equation with a controller acting on the subset of the boundary

$$\begin{cases} iy_t + \Delta^2 y = 0, & \text{in } \Omega \times (0, T), \\ y = 0, \quad \frac{\partial y}{\partial \nu} = v \chi_{\Gamma_0}, & \text{on } \partial\Omega \times (0, T), \\ y(x, 0) = y_0(x), & \text{in } \Omega. \end{cases} \quad (1.2)$$

Here and henceforth, χ_{Γ_0} is the characteristic function of the set Γ_0 and Δ is the Laplacian in the space variable $x \in \Omega$. In (1.2), $y(\cdot, t)$ can be considered as the probability amplitude of the state and $v(\cdot, t)$ is the control. Both are complex valued functions. The control space of system (1.2) is chosen to be $L^2((0, T) \times \Gamma_0)$.

As we will show later in Section 4, the well-posedness of the system is given as follows. For any initial data $y_0 \in H^{-2}(\Omega)$ and $v \in L^2(\Gamma_0 \times (0, T))$, there exists a unique solution $y \in C([0, T]; H^{-2}(\Omega))$ to (1.2), in the transposition sense (see [13]).

Manuscript received May 24, 2011. Revised January 8, 2012.

¹Laboratory of Mathematics and Complex Systems, School of Mathematical Sciences, Beijing Normal University, Beijing 100875, China. E-mail: chuang.zheng@bnu.edu.cn

²Department of Mathematics and Statistics, Southwest University, Chongqing 400715, China.
E-mail: zhouzc@swu.edu.cn

*Project supported by the Fundamental Research Funds for the Central Universities (No. XDJK2009C099), the National Natural Science Foundation of China (Nos. 11001018, 11026111) and the Specialized Research Fund for the Doctoral Program of Higher Education (No. 201000032006).

In this paper, we are interested in the exact (boundary) controllability problem of (1.2), which is stated as follows. Let y_0 be a given function in $H^{-2}(\Omega)$ and let $T > 0$ be given. Whether there exists a boundary function v on $\Gamma_0 \times (0, T)$ such that the solution to the equation (1.2) satisfies $y(0) = y_0$ and $y(T) = 0$ in Ω ? If such a control v exists, we say that the system (1.2) is exactly controllable from y_0 to the rest at the time T by the boundary control v .

The fourth order Schrödinger equation arises in many scientific fields such as quantum mechanics, nonlinear optics and plasma physics, and has been intensively studied with fruitful references. For instance, the well-posedness and existence of the solutions has been shown in [7–8, 16–17] by means of the energy method and harmonic analysis. However, it is still unknown for the corresponding controllability properties.

As far as we know, there are plenty of references concerning the controllability properties of the second order Schrödinger equations (see [14]). For the higher order operators, these control problems are mostly studied for parabolic cases, such as the approximate controllability of the nonlinear equation (see [4]), the null boundary controllability of the $1 - d$ and $N - d$ cases (see [3, 10]), etc. Recent results (see [2, 15, 18]) considered the exact observability and some equivalent assertions for the skew-adjoint operators, which can be seen as an abstract model for the higher order Schrödinger equations.

By establishing the control theory for the linear fourth order model (1.2), we hope it would be helpful to understand the phenomena of the high dimensional higher order nonlinear systems. In this paper, we attempt to establish the boundary controllability properties of system (1.2) by means of the Hilbert Uniqueness Method (HUM) and the multiplier techniques. More precisely, by classical duality arguments (see [12]), the above controllability property is equivalent to a (boundary) observability estimate of the following uncontrolled Schrödinger equation:

$$\begin{cases} i\varphi_t + \Delta^2 \varphi = 0, & \text{in } \Omega \times (0, T), \\ \varphi = 0, \quad \frac{\partial \varphi}{\partial \nu} = 0, & \text{on } \partial\Omega \times (0, T), \\ \varphi(x, 0) = \varphi^0, & \text{in } \Omega. \end{cases} \quad (1.3)$$

Our first result is the observability inequality of (1.3), which reads as follows.

Theorem 1.1 *For equation (1.3), the solution to (1.3) satisfies*

$$\|\varphi^0\|_{H_0^2(\Omega)}^2 \leq C \int_0^T \int_{\Gamma_0} |\Delta \varphi|^2 d\sigma dt, \quad \forall \varphi^0 \in H_0^2(\Omega). \quad (1.4)$$

Here and thereafter, we use C to denote a generic positive constant (depending only on T , Ω and Γ_0) which may vary from line to line.

As a direct consequence of Theorem 1.1, the controllability property of (1.2) is stated as follows.

Theorem 1.2 *Let $T > 0$, Γ_0 be defined by (1.1) and $\Sigma_0 = \Gamma_0 \times (0, T)$. Then, for any $y_0 \in H^{-2}(\Omega)$, there exists a $v \in L^2(\Gamma_0 \times (0, T))$ such that the unique solution $y \in C([0, T]; H^{-2}(\Omega))$ to (1.2) satisfies $y(T) = 0$.*

Remark 1.1 Without loss of generality, the final state $y(T)$ is driven to the rest. This is due to the fact that system (1.2) is linear and time reversible. This phenomenon happens in

finite-dimensional linear controlled systems, and the situation is completely different from the case of the time irreversible one, such as the heat equation.

The rest of the paper is organized as follows. An identity of the fourth order Schrödinger operator is given in Section 2 by choosing a suitable multiplier and playing carefully with the boundary terms. In Section 3, we show the observability estimate (1.4). The well-posedness and the exact controllability of the system (1.2) are both given in Section 4. Finally, we state some open problems and further comments in the last section.

2 Identity via Multipliers

This section is dedicated to establishing two fundamental identities by multipliers. Letting $f \in L^2(0, T; H_0^2(\Omega))$, we consider the system

$$\begin{cases} i\theta_t + \Delta^2 \theta = f, & \text{in } \Omega \times (0, T), \\ \theta = 0, \quad \frac{\partial \theta}{\partial \nu} = 0, & \text{on } \partial\Omega \times (0, T), \\ \theta(x, 0) = \theta^0, & \text{in } \Omega. \end{cases} \quad (2.1)$$

First, we show the following lemma.

Lemma 2.1 *Let $q = q(x, t) \in C^3(\overline{Q}, \mathbb{R}^n)$ with \overline{Q} being the closed set of Q . For every solution to (2.1) with $f \in \mathcal{D}(Q)$ and $\varphi^0 \in \mathcal{D}(\Omega)$, the following identity holds:*

$$\begin{aligned} 0 &= \frac{i}{2} \int_{\Omega} \theta \nabla \bar{\theta} \cdot q|_0^T - \frac{i}{2} \int_Q (\theta \nabla \bar{\theta}_t + \bar{\theta} \nabla \theta_t) \cdot q - \frac{i}{2} \int_Q \theta \nabla \bar{\theta} \cdot q_t \\ &\quad - \frac{1}{2} \int_{\Sigma} |\Delta \theta|^2 q \cdot \nu - \frac{1}{2} \int_Q (\nabla \theta H(\Delta \bar{\theta}) - \nabla \bar{\theta} H(\Delta \theta)) \cdot q \\ &\quad + \frac{1}{2} \int_Q \sum_{i,j} (\theta_{x_i x_j} \Delta \bar{\theta} q_{x_i}^j + \theta_{x_i} \Delta \bar{\theta} q_{x_i x_j}^j + 3 \Delta \theta \bar{\theta}_{x_i x_j} q_{x_i}^j + 2 \Delta \theta \bar{\theta}_{x_j} q_{x_i x_i}^j) \\ &\quad + \frac{1}{2} \int_Q (\Delta \theta \nabla \bar{\theta} \cdot \nabla \operatorname{div}_x q + \Delta \bar{\theta} \nabla \theta \cdot \nabla \operatorname{div}_x q) - \int_Q f \left(\nabla \bar{\theta} \cdot q + \frac{1}{2} \bar{\theta} \operatorname{div}_x q \right), \end{aligned} \quad (2.2)$$

where $H(f)$ is the Hessian Matrix of f .

Remark 2.1 For convenience, we drop all $dx, d\sigma, dt$ terms in all integrals here and thereafter. More precisely, we write $\int_{\Omega}(\cdot)$, $\int_Q(\cdot)$, $\int_{\Sigma}(\cdot)$ and $\int_{\Sigma_0}(\cdot)$ instead of $\int_{\Omega}(\cdot)dx$, $\int_{\Omega \times (0, T)}(\cdot)dxdt$, $\int_{\Gamma \times (0, T)}(\cdot)d\sigma dt$ and $\int_{\Gamma_0 \times (0, T)}(\cdot)d\sigma dt$, respectively.

Proof of Lemma 2.1 Step 1 Multiplying (1.3) by $\nabla \bar{\theta} \cdot q + \frac{1}{2} \bar{\theta} \operatorname{div}_x q$, integrating on Q of the left-hand side of (1.3) (abbreviated by ILHS), we get

$$\begin{aligned} \text{ILHS} &= \frac{i}{2} \int_{\Omega} \theta \nabla \bar{\theta} \cdot q|_0^T - \frac{i}{2} \int_Q (\theta \nabla \bar{\theta}_t + \bar{\theta} \nabla \theta_t) \cdot q - \frac{1}{2} \int_Q \nabla(\Delta \theta) \bar{\theta} \cdot \nabla \operatorname{div}_x q \\ &\quad - \frac{i}{2} \int_Q \theta \nabla \bar{\theta} \cdot q_t - \frac{1}{2} \int_{\Sigma} \Delta \theta H(\bar{\theta}) q \cdot \nu - \frac{1}{2} \int_Q (\nabla \theta H(\Delta \bar{\theta}) - \nabla \bar{\theta} H(\Delta \theta)) \cdot q \\ &\quad + \int_Q \sum_{i,j} \left(\frac{1}{2} \Delta \theta \bar{\theta}_{x_i x_j} q_{x_i}^j - \frac{1}{2} \theta_{x_i} \Delta \bar{\theta}_{x_j} q_{x_i}^j - \Delta \theta_{x_i} \bar{\theta}_{x_j} q_{x_i}^j \right). \end{aligned} \quad (2.3)$$

In fact, $\text{ILHS} = \int_Q (i\theta_t + \Delta^2\theta)(\nabla\bar{\theta} \cdot q + \frac{1}{2}\bar{\theta}\text{div}_x q)$ equals $(A+B) + C + D$ with the notation

$$A+B = \frac{i}{2} \int_{\Omega} \theta \nabla \bar{\theta} \cdot q|_0^T - \frac{i}{2} \int_Q (\theta \nabla \bar{\theta}_t \cdot q + \bar{\theta} \nabla \theta_t \cdot q) - \frac{i}{2} \int_Q \theta \nabla \bar{\theta} \cdot q_t, \quad (2.4)$$

$$C = \int_Q \Delta^2 \theta \nabla \bar{\theta} \cdot q, \quad D = \frac{1}{2} \int_Q \Delta^2 \theta \bar{\theta} \text{div}_x q. \quad (2.5)$$

Taking into account the boundary conditions, we arrive at

$$C = - \int_Q \sum_{i,j} \Delta \theta_{x_i} \bar{\theta}_{x_j} q_{x_i}^j - I, \quad (2.6)$$

$$D = \frac{1}{2} \int_Q \nabla \bar{\theta} H(\Delta \theta) q + \frac{1}{2} I - \frac{1}{2} \int_Q \nabla(\Delta \theta) \cdot \bar{\theta} \nabla \text{div}_x q, \quad (2.7)$$

with

$$I = \int_Q \nabla(\Delta \theta) H(\bar{\theta}) q.$$

Moreover,

$$\begin{aligned} I &= \int_{\Sigma} \Delta \theta H(\bar{\theta}) q \cdot \nu - \int_Q \Delta \theta \left(\nabla(\Delta \bar{\theta}) \cdot q + \sum_{i,j} \bar{\theta}_{x_i x_j} q_{x_i}^j \right) \\ &= \int_{\Sigma} \Delta \theta H(\bar{\theta}) q \cdot \nu + \int_Q \nabla \theta H(\Delta \bar{\theta}) q + \int_Q \sum_{i,j} (\theta_{x_i} \Delta \bar{\theta}_{x_j} q_{x_i}^j - \Delta \theta \bar{\theta}_{x_i x_j} q_{x_i}^j). \end{aligned} \quad (2.8)$$

Combining (2.6)–(2.8), we get

$$\begin{aligned} C+D &= -\frac{1}{2} \int_{\Sigma} \Delta \theta H(\bar{\theta}) q \cdot \nu - \int_Q \sum_{i,j} \left(\Delta \theta_{x_i} \bar{\theta}_{x_j} q_{x_i}^j + \frac{1}{2} \theta_{x_i} \Delta \bar{\theta}_{x_j} q_{x_i}^j - \frac{1}{2} \Delta \theta \bar{\theta}_{x_i x_j} q_{x_i}^j \right) \\ &\quad - \frac{1}{2} \int_Q (\nabla \theta H(\Delta \bar{\theta}) q - \nabla \bar{\theta} H(\Delta \theta) q) - \frac{1}{2} \int_Q \nabla(\Delta \theta) \bar{\theta} \cdot \nabla \text{div}_x q. \end{aligned} \quad (2.9)$$

Finally, from (2.4) and (2.9), we obtain the desired identity (2.3).

Step 2 Integrating by parts with respect to x , we get

$$\int_Q \sum_{i,j} \theta_{x_i} \Delta \bar{\theta}_{x_j} q_{x_i}^j = - \int_Q \sum_{i,j} (\theta_{x_i x_j} \Delta \bar{\theta} q_{x_i}^j + \theta_{x_i} \Delta \bar{\theta} q_{x_i x_j}^j), \quad (2.10)$$

$$\int_Q \sum_{i,j} \Delta \theta_{x_i} \bar{\theta}_{x_j} q_{x_i}^j = - \int_Q \sum_{i,j} \Delta \theta (\bar{\theta}_{x_i x_j} q_{x_i}^j + \bar{\theta}_{x_j} q_{x_i x_i}^j), \quad (2.11)$$

$$\int_Q \nabla(\Delta \theta) \bar{\theta} \cdot \nabla \text{div}_x q = - \int_Q \Delta \theta (\nabla \bar{\theta} \cdot \nabla \text{div}_x q + \bar{\theta} \Delta(\text{div}_x q)). \quad (2.12)$$

On the other side, since $\theta = 0$ and $\frac{\partial \theta}{\partial \nu} = 0$ on the boundary Σ , we have $\theta_{x_i} = 0$, $i = 1, \dots, n$ and $\theta_{x_i, x_j} = \frac{\partial \theta_{x_i}}{\partial \nu} \nu_j$, $i, j = 1, \dots, n$ for any $x \in \Gamma$. Consequently, for any $x \in \Gamma$, it holds

$$\begin{aligned} \sum_{i,j} q_i \theta_{x_i x_j} \nu_j &= \sum_{i,j,k} q_i \theta_{x_j, x_k} \nu_k \nu_i \nu_j = \left(\sum_i q_i \nu_i \right) \sum_{j,k} (\theta_{x_j, x_k} \nu_k \nu_j) \\ &= \left(\sum_i q_i \nu_i \right) \sum_k \frac{\partial \theta_{x_k}}{\partial \nu} \nu_k = \left(\sum_i q_i \nu_i \right) \sum_k \theta_{x_k, x_k} = \Delta \theta (q \cdot \nu). \end{aligned}$$

Hence,

$$\int_{\Sigma} \Delta \theta H(\bar{\theta}) q \cdot \nu = \int_{\Sigma} \Delta \theta \sum_{i,j} q_i \bar{\theta}_{x_i x_j} \nu_j = \int_{\Sigma} |\Delta \theta|^2 q \cdot \nu. \quad (2.13)$$

Taking (2.10)–(2.13) into (2.3) and putting the right-hand side of (2.1) into account, we finish the proof of (2.2).

The conservation laws hold for the solutions to (1.3).

Lemma 2.2 *For any positive time t , the solution φ to (1.3) satisfies*

$$\|\varphi(t)\|_{L^2(\Omega)} = \|\varphi(0)\|_{L^2(\Omega)}, \quad (2.14)$$

$$\|\nabla \varphi(t)\|_{L^2(\Omega)} = \|\nabla \varphi(0)\|_{L^2(\Omega)}, \quad (2.15)$$

$$\|\Delta \varphi(t)\|_{L^2(\Omega)} = \|\Delta \varphi(0)\|_{L^2(\Omega)}. \quad (2.16)$$

Remark 2.2 Note that in quantum mechanics, the conservation of the norms validates the Born's statistical interpretation of the probability amplitude function $\varphi(x, t)$. More precisely, $\int_{\Omega} |\varphi(x, t)|^2 dx$ represents the probability of finding the particle in domain Ω at the time t and the conservation law provides the particle which will not disappear in Ω .

Proof of Lemma 2.2 We use multipliers $\bar{\varphi}$, $\Delta \bar{\varphi}$ and $\bar{\varphi}_t$ on (1.3) and we achieve the above identities (2.14), (2.15) and (2.16), respectively.

3 Observability

Proposition 3.1 *For every $T > 0$, there exist $c_i = c_i(T, \Omega) > 0$ ($i = 1, 2$), such that*

$$\int_0^T \int_{\Gamma_0} |\Delta \varphi|^2 \leq c_1 \|\varphi^0\|_{H_0^2(\Omega)}^2 \quad (3.1)$$

and

$$\|\varphi^0\|_{H_0^2(\Omega)}^2 \leq c_2 \int_0^T \int_{\Gamma_0} |\Delta \varphi|^2 \quad (3.2)$$

for every solution $\varphi = \varphi(x, t)$ to the problem (1.3) with $\varphi^0 \in H_0^2(\Omega)$.

Proof For the admissibility inequality (3.1), we choose $q = q(x) \in C^3(\bar{Q}, \mathbb{R}^n)$ such that $q = \nu$ on Γ (see [13] for the construction of this vector field). Taking the real part of the identity (2.2) with $f = 0$, we obtain

$$\begin{aligned} \frac{1}{2} \int_{\Sigma} |\Delta \varphi|^2 q \cdot \nu &= -\frac{1}{2} \operatorname{Im} \int_{\Omega} \varphi \nabla \bar{\varphi} \cdot q \Big|_0^T + \frac{1}{2} \operatorname{Re} \int_Q (\Delta \varphi \nabla \bar{\varphi} \cdot \nabla \operatorname{div}_x q + \Delta \varphi \bar{\varphi} \Delta \operatorname{div}_x q) \\ &\quad + \frac{1}{2} \int_Q \sum_{i,j} (\varphi_{x_i x_j} \Delta \bar{\varphi} q_{x_i}^j + \varphi_{x_i} \Delta \bar{\varphi} q_{x_i x_j}^j + 3 \Delta \varphi \bar{\varphi}_{x_i x_j} q_{x_i}^j + 2 \Delta \varphi \bar{\varphi}_{x_j} q_{x_i x_i}^j). \end{aligned}$$

Consequently,

$$\begin{aligned}
& \frac{1}{2} \int_{\Sigma} |\Delta \varphi|^2 \\
& \leq k_1 \|q\|_{L^\infty(\Omega)} (\|\varphi(T)\|_{L^2(\Omega)}^2 + \|\nabla \varphi(T)\|_{L^2(\Omega)}^2 + \|\varphi(0)\|_{L^2(\Omega)}^2 + \|\nabla \varphi(0)\|_{L^2(\Omega)}^2) \\
& \quad + k_2 \|q\|_{W^{2,\infty}(\Omega)} \int_0^T ((\|H(\varphi)\|_{L^2(\Omega)} + \|\nabla \varphi\|_{L^2(\Omega)}) \|\Delta \varphi\|_{L^2(\Omega)} + \|\varphi\|_{L^2(\Omega)} \|\nabla \varphi\|_{L^2(\Omega)}) \\
& \quad + k_3 \|q\|_{W^{3,\infty}(\Omega)} \int_0^T (\|\Delta \varphi\|_{L^2(\Omega)} \|\nabla \varphi\|_{L^2(\Omega)} + \|\Delta \varphi\|_{L^2(\Omega)} \|\varphi\|_{L^2(\Omega)}).
\end{aligned}$$

Combining with the conservation law in Lemma 2.2, we obtain

$$\int_0^T \int_{\Gamma_0} |\Delta \varphi|^2 \leq c_1 \|\varphi^0\|_{H_0^2(\Omega)}^2, \quad \forall \varphi^0 \in \mathcal{D}(\Omega).$$

Since $\mathcal{D}(\Omega)$ is dense in $H_0^2(\Omega)$, the estimate (3.1) holds for every solution to the problem (1.3) with the initial data $\varphi^0 \in H_0^2(\Omega)$.

Now we prove (1.4). We choose $q(x, t) = m(x) = x - x_0$. By using (2.2), we obtain

$$\int_{\Sigma} m \cdot \nu |\Delta \varphi|^2 = -\text{Im} \int_{\Omega} \varphi \nabla \overline{\varphi} \cdot m|_0^T + 4T \int_{\Omega} |\Delta \varphi|^2.$$

Furthermore, there exists an $\varepsilon > 0$ such that

$$\left| \text{Im} \int_{\Omega} \varphi \nabla \overline{\varphi} \cdot m|_0^T \right| \leq c_\varepsilon \|\varphi^0\|_{L^2(\Omega)}^2 + \varepsilon \|\varphi^0\|_{H_0^1(\Omega)}^2.$$

Thus

$$4T \|\varphi^0\|_{H_0^2(\Omega)}^2 \leq C \left(\int_{\Sigma_0} m \cdot \nu |\Delta \varphi|^2 + c_\varepsilon \|\varphi^0\|_{L^2(\Omega)}^2 + \varepsilon \|\varphi^0\|_{H_0^1(\Omega)}^2 \right). \quad (3.3)$$

To conclude the proof of (1.4), it is enough to prove the following estimates:

$$\|\varphi^0\|_{L^2(\Omega)}^2 \leq C \int_{\Sigma_0} m \cdot \nu |\Delta \varphi|^2, \quad (3.4)$$

$$\|\varphi^0\|_{H_0^1(\Omega)}^2 \leq C \int_{\Sigma_0} m \cdot \nu |\Delta \varphi|^2. \quad (3.5)$$

We argue by contradiction. We only state the proof of (3.5) and the one for (3.4) can be obtained directly with the Poincaré inequality. If (3.5) is not satisfied, then for any $C > 0$, there exists a sequence $\{\varphi_n\}$ of the solutions to (1.3) such that

$$\|\varphi_n(0)\|_{H_0^1(\Omega)} = 1, \quad \forall n \in \mathbb{N} \quad (3.6)$$

and

$$\int_{\Sigma_0} m \cdot \nu |\Delta \varphi_n|^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.7)$$

Obviously, $\{\varphi_n(0)\}$ is bounded in $H_0^1(\Omega)$ and from (3.3), it is also bounded in $H_0^2(\Omega)$. Then

$$\{\varphi_n\} \text{ is bounded in } L^\infty(0, T; H_0^2(\Omega)) \cap W^{1,\infty}(0, T; H^{-2}(\Omega)).$$

Thus, by extracting a subsequence (still denoted by $\{\varphi_n\}$), we have

- (i) $\varphi_n \rightarrow \varphi$ in $L^\infty(0, T; H_0^2(\Omega))$ weak *;
- (ii) $(\varphi_n)_t \rightarrow \varphi_t$ in $L^\infty(0, T; L^2(\Omega))$ weak *.

The function $\varphi \in L^\infty(0, T; H_0^2(\Omega)) \cap W^{1,\infty}(0, T; H^{-2}(\Omega))$ is clearly a solution to (1.3), and from the compactness of the embedding (see [19])

$$L^\infty(0, T; H_0^2(\Omega)) \cap W^{1,\infty}(0, T; H^{-2}(\Omega)) \rightarrow C([0, T]; H_0^1(\Omega))$$

and (3.6), we deduce

$$\|\varphi(0)\|_{H_0^1(\Omega)} = 1. \quad (3.8)$$

On the other hand, (3.7) implies

$$\Delta\varphi = 0 \quad \text{on } \Sigma_0,$$

which combined with (1.3) implies $\varphi \equiv 0$, from Holmgren's Uniqueness Theorem (see [9, Chapter V, Theorem 5.3.3]). This is in contradiction with (3.8). This ends the proof of (3.5).

Taking (3.4) and (3.5) into account, (1.4) is a direct consequence of (3.3).

4 Well-posedness and Exact Controllability

We say that $y \in L^\infty(0, T; H^{-2}(\Omega))$ is a solution to (1.2) in the transposition sense if and only if

$$\int_0^T \langle y(t), \bar{f}(t) \rangle_{(H^{-2}(\Omega), H_0^2(\Omega))} dt + i \langle y(0), \bar{\theta}(0) \rangle_{(H^{-2}(\Omega), H_0^2(\Omega))} + \int_\Sigma v \Delta \bar{\theta} d\Sigma = 0 \quad (4.1)$$

for every $f \in L^2(0, T; H_0^2(\Omega))$, where $\theta = \theta(x, t)$ is the solution to the problem (2.1) with $\theta(T) = 0$.

The following proposition claims the existence of a unique solution to system (1.2) in the sense of transposition.

Proposition 4.1 *Let $v \in L^2(\Sigma)$. Then there exists a unique solution $y \in C([0, T]; H^{-2}(\Omega))$ in the transposition sense, to the problem (1.2) with the initial data $y_0 \in H^{-2}(\Omega)$. Furthermore, the map $v \mapsto y$ is linear and continuous from $L^2(\Sigma)$ into $C([0, T]; H^{-2}(\Omega))$.*

Proof Without loss of generality, we assume that $y_0 = 0$, which is due to the time reversibility of system (1.2). It is not hard to prove that

$$\|\theta(t)\|_{H_0^2(\Omega)} \leq \|f\|_{L^1(0, T; H_0^2(\Omega))}, \quad \forall t \in [0, T].$$

Applying the identity (2.2) with a vector field $q = \nu$ on Γ and using the above estimate, we obtain

$$\|\Delta\theta\|_{L^2(\Sigma)} \leq c\|f\|_{L^1(0, T; H_0^2(\Omega))}.$$

Hence, we have

$$\left| \operatorname{Re} \int_\Sigma v \Delta \bar{\theta} d\Sigma \right| \leq \|v\|_{L^2(\Sigma)} \|\Delta\theta\|_{L^2(\Sigma)} \leq c\|v\|_{L^2(\Sigma)} \|f\|_{L^1(0, T; H_0^2(\Omega))}. \quad (4.2)$$

It means that the map from f into $\operatorname{Re} \int_{\Sigma} v \Delta \bar{\theta} d\Sigma$ is linear and continuous from $L^1(0, T; H_0^2(\Omega))$ into \mathbb{R} .

Hence, there exists a unique $y \in L^\infty(0, T; H^{-2}(\Omega))$ that satisfies (4.1) for every $f \in L^1(0, T; H_0^2(\Omega))$.

From (4.1) and (4.2), we have

$$\|y\|_{L^\infty(0, T; H^{-2}(\Omega))} \leq c\|v\|_{L^2(\Sigma)}. \quad (4.3)$$

Thus, the map $v \mapsto y$ is continuous from $L^2(\Sigma)$ into $L^\infty(0, T; H^{-2}(\Omega))$.

Moreover, $y \in C([0, T]; H^{-2}(\Omega))$. Indeed, we consider $\{v_n\}_{n \in \mathbb{N}} \subset \mathcal{D}(0, T; C^2(\Gamma))$ such that

$$v_n \rightarrow v \quad \text{strongly in } L^2(\Sigma). \quad (4.4)$$

Let y_n be the solution to (1.2) with the boundary condition v_n . Since v_n is regular, in particular, we have $y_n \in C([0, T]; H^{-2}(\Omega))$.

From (4.3) and (4.4), we have

$$y_n \rightarrow y \quad \text{in } L^\infty(0, T; H^{-2}(\Omega)).$$

Since $C([0, T]; H^{-2}(\Omega))$ is a closed subspace of $L^\infty(0, T; H^{-2}(\Omega))$, we have $y \in C([0, T]; H^{-2}(\Omega))$.

Proof of Theorem 1.2 We consider the problem

$$\begin{cases} iy_t + \Delta^2 y = 0, & \text{in } \Omega \times (0, T), \\ y = 0, \quad \frac{\partial y}{\partial \nu} = v \chi_{\Gamma_0}, & \text{on } \partial\Omega \times (0, T), \\ y(T) = 0, & \text{in } \Omega. \end{cases} \quad (4.5)$$

It is easy to see that, by multiplying (4.5) by $\bar{\varphi}$, taking the real part, and integrating it by parts, we have the following identity:

$$\langle -iy(0), \varphi^0 \rangle = \int_{\Sigma_0} |\Delta \varphi|^2 d\Sigma, \quad \forall \varphi^0 \in \mathcal{D}(\Omega),$$

where φ is the corresponding solution to system (1.3) with an initial data φ^0 . Let Λ be a linear continuous operator from $H_0^2(\Omega)$ into $H^{-2}(\Omega)$ defined by $\Lambda \varphi^0 = -iy(0)$, where $y = y(x, t)$ is the solution to the problem (4.5).

From Proposition 3.1, we have $\langle \Lambda \varphi^0, \varphi^0 \rangle \geq c \|\varphi^0\|_{H_0^2(\Omega)}^2$. Hence Λ is an isomorphism from $H_0^2(\Omega)$ to $H^{-2}(\Omega)$ and the theorem is proved. The control v is chosen by $v = \Delta \varphi$ on Σ_0 where φ is the solution to (1.2) with the initial data $\varphi^0 = \Lambda^{-1}(-iy(0))$.

5 Further Comments and Open Problems

(1) Transmutation method We derived Theorem 1.2 by means of the multiplier techniques. One could expect a different proof by means of the transmutation method. Roughly speaking, the controllability for system (1.2) can be seen as a combination of the exact controllability for the Schrödinger equation on a segment (see [14]) and a plate equation (see [11]),

following the instruction in [15]. However, both methods cannot tell us whether the control domain is sharp. It is still an open problem.

(2) Internal controllability In this paper, we have only dealt with the L^2 -Neumann boundary control. On the other hand, one can expect the same result with L^2 controls supported in a neighborhood of the boundary, by following the same methodology in [14]. Furthermore, for the controlled wave equation, the sharp control domain is the one satisfying GCC condition (see [1]) instead of the one in (1.1). It is still an open problem whether the same happens for system (1.2).

(3) Carleman estimate There are several different methods to derive observability inequalities. The Carleman estimate (see [5–6, 20]) is developed to derive the observability inequalities in a bounded domain with potentials. One may expect to solve the control problem for the fourth order Schrödinger with potentials by means of the corresponding global Carleman estimate.

Acknowledgment The first author wishes to thank Institut Henri Poincaré (Paris, France) for providing a very stimulating environment during the “Control of Partial and Differential Equations and Applications” program in the Fall 2010.

References

- [1] Bardos, C., Lebeau, G. and Rauch, J., Sharp sufficient conditions for the observation, control and stabilization of waves from the boundary, *SIAM J. Control Optim.*, **30**, 1992, 1024–1065.
- [2] Burq, N. and Zworski, M., Geometric control in the presence of a black box, *J. Amer. Math. Soc.*, **17**, 2004, 443–471.
- [3] Chou, H. and Guo, Y., Null boundary controllability for a fourth order semilinear equation, *Taiwanese J. Math.*, **10**, 2006, 251–263.
- [4] Díaz, J. I. and Ramos, A. M., On the approximate controllability for higher order parabolic nonlinear equations of Cahn-Hilliard type, *Internat. Ser. Numer. Math.*, **126**, 1998, 111–127.
- [5] Fu, X., Yong, J. and Zhang, X., Controllability and observability of a heat equation with hyperbolic memory kernel, *J. Differential Equations*, **247**, 2009, 2395–2439.
- [6] Fursikov, A. and Imanuvilov, O., Controllability of Evolution Equations, Lecture Notes Series, Vol. 34, Seoul National University, Seoul, 1996.
- [7] Hao, C., Hsiao, L. and Wang, B., Wellposedness for the fourth order nonlinear Schrödinger equations, *J. Math. Anal. Appl.*, **320**, 2006, 246–265.
- [8] Hao, C., Hsiao, L. and Wang, B., Well-posedness of Cauchy problem for the fourth order nonlinear Schrödinger equations in multi-dimensional spaces, *J. Math. Anal. Appl.*, **328**, 2007, 58–83.
- [9] Hörmander, L., Linear Partial Differential Operators, Springer-Verlag, Berlin, 1976.
- [10] Lin, P. and Zhou, Z., Observability estimate for a one-dimensional fourth order parabolic equation, Proceedings of the 29th Chinese Control Conference, China, 2010, 830–832.
- [11] Lions, J. L., Contrôlabilité Exacte, Perturbations et Stabilisation de Systèmes Distribués, Tome 1, Recherches en Mathématiques Appliquées, Vol. 8, Masson, Paris, 1988, 1–68.
- [12] Lions, J. L., Exact controllability, stabilization and perturbations for distributed systems, *SIAM Rev.*, **30**, 1988, 1–68.
- [13] Lions, J. L. and Magenes, E., Problèmes aux Limites Non Homogènes et Applications, Dunod, Paris, 1968.
- [14] Machtyngier, E., Exact controllability for the Schrödinger equation, *SIAM J. Control Optim.*, **32**, 1994, 24–34.

- [15] Miller, L., Controllability cost of conservative systems: resolvent condition and transmutation, *J. Funct. Anal.*, **218**, 2005, 425–444.
- [16] Pausader, B., Global well-posedness for energy critical fourth-order Schrödinger equations in the radial case, *Dyn. Partial Differ. Equ.*, **4**, 2007, 197–225.
- [17] Pausader, B., The cubic fourth-order Schrödinger equation, *J. Funct. Anal.*, **256**, 2009, 2473–2517.
- [18] Ramdani, K., Takahashi, T., Tenenbaum, G. and Tucsnak, M., A spectral approach for the exact observability of infinite-dimensional systems with skew-adjoint generator, *J. Funct. Anal.*, **226**, 2005, 193–229.
- [19] Simon, J., Compact sets in the space $L^p(0, T; B)$, *Ann. Mat. Pura Appl.*, **146**(4), 1987, 65–96.
- [20] Zhang, X., Exact controllability of semilinear plate equations, *Asympt. Anal.*, **27**, 2001, 95–125.